

Lähdeluettelo luku 1

Abidin NZ, Adam MB. Prediction of vertical jump height from anthropometric factors in male and female martial arts athletes. *Malays J Med Sci.* 2013 Jan;20(1):39–45.

Arrese AL, Ostáriz ES. Skinfold thicknesses associated with distance running performance in highly trained runners. *J Sports Sci.* 2006 Jan;24(1):69–76.

Balaguera-Cortes L, Wallman KE, Fairchild TJ, Guelfi KJ. Energy intake and appetite-related hormones following acute aerobic and resistance exercise. *Appl Physiol Nutr Metab.* 2011 Dec;36(6):958–66.

Becker GF, Macedo RC, Cunha Gdos S, Martins JB, Laitano O, Reischak-Oliveira A. Combined effects of aerobic exercise and high-carbohydrate meal on plasma acylated ghrelin and levels of hunger. *Appl Physiol Nutr Metab.* 2012 Feb;37(1):184–92.

Berglund L, Sundgot-Borgen J, Berglund B. Adipositas athletica: a group of neglected conditions associated with medical risks. *Scand J Med Sci Sports.* 2011 Oct;21(5):617–24.

Boström P, Wu J, Jedrychowski MP, Korde A, Ye L, Lo JC, Rasbach KA, Boström EA, Choi JH, Long JZ, Kajimura S, Zingaretti MC, ym. A PGC1- α -dependent myokine that drives brown-fat-like development of white fat and thermogenesis. *Nature.* 2012 Jan 11;481(7382):463–8.

Bratland-Sanda S, Sundgot-Borgen J. Eating disorders in athletes: overview of prevalence, risk factors and recommendations for prevention and treatment. *Eur J Sport Sci.* 2013;13(5):499–508.

Carbone JW, McClung JP, Pasiakos SM. Skeletal muscle responses to negative energy balance: effects of dietary protein. *Adv Nutr.* 2012 Mar 1;3(2):119–26.

Chaston TB, Dixon JB, O'Brien PE. Changes in fat-free mass during significant weight loss: a systematic review. *Int J Obes.* 2007;11:743–750

Churchward-Venne TA, Murphy CH, Longland TM, Phillips SM. Role of protein and amino acids in promoting lean mass accretion with resistance exercise and attenuating lean mass loss during energy deficit in humans. *Amino Acids.* 2013 Aug;45(2):231–40.

Dangour AD, Lock K, Hayter A, Aikenhead A, Allen E, Uauy R. Nutrition-related health effects of organic foods: a systematic review. *Am J Clin Nutr.* 2010 Jul;92(1):203–10.

Deighton K, Barry R, Connon CE, Stensel DJ. Appetite, gut hormone and energy intake responses to low volume sprint interval and traditional endurance exercise. *Eur J Appl Physiol.* 2013 May;113(5):1147–56.

De Lorenzo A, Bertini I, Candeloro N, Piccinelli R, Innocente I, Brancati A. A new predictive equation to calculate resting metabolic rate in athletes. *J Sports Med Phys Fitness.* 1999 Sep;39(3):213–9.

De Maddalena C, Vodo S, Petroni A, Aloisi AM. Impact of testosterone on body fat composition. *J Cell Physiol.* 2012;11:3744–3748.

Deutz RC, Benardot D, Martin DE, Cody MM. Relationship between energy deficits and body composition in elite female gymnasts and runners. *Med Sci Sports Exerc.* 2000 Mar;32(3):659–68.

Dirlewanger M, di Vetta V, Guenat E, Battilana P, Seematter G, Schneiter P, Jequier E, Tappy L. Effects of short-term carbohydrate or fat overfeeding on energy expenditure and plasma leptin concentrations in healthy female subjects. *Int J Obes Relat Metab Disord.* 2000;11:1413–1418.

Dolan E, Crabtree N, McGoldrick A, Ashley DT, McCaffrey N, Warrington GD. Weight regulation and bone mass: a comparison between professional jockeys, elite amateur boxers, and age, gender and BMI matched controls. *J Bone Miner Metab.* 2012 Mar;30(2):164–70.

Doucet E, Imbeault P, St-Pierre S, Almeras N, Mauriege P, Despres JP, Bouchard C, Tremblay A. Greater than predicted decrease in energy expenditure during exercise after body weight loss in obese men. *Clin Sci.* 2003;11:89–95.

Dulloo AG, Jacquet J, Montani JP. How dieting makes some fatter: from a perspective of human body composition autoregulation. *Proc Nutr Soc.* 2012;11:379–389.

El Ghoch M, Soave F, Calugi S, Dalle Grave R. Eating disorders, physical fitness and sport performance: a systematic review. *Nutrients.* 2013 Dec 16;5(12):5140–60.

Frankenfield DC. Bias and accuracy of resting metabolic rate equations in non-obese and obese adults. *Clin Nutr.* 2013 Dec;32(6):976–82.

Frankenfield D, Roth-Yousey L, Compher C. Comparison of predictive equations for resting metabolic rate in healthy nonobese and obese adults: a systematic review. *J Am Diet Assoc.* 2005 May;105(5):775–89.

Garthe I, Raastad T, Refsnes PE, Koivisto A, Sundgot-Borgen J. Effect of two different weight-loss rates on body composition and strength and power-related performance in elite athletes. *Int J Sport Nutr Exerc Metab.* 2011 Apr;21(2):97–104.

Garthe I, Raastad T, Refsnes PE, Sundgot-Borgen J. Effect of nutritional intervention on body composition and performance in elite athletes. *Eur J Sport Sci.* 2013;13(3):295–303.

Gibson JH, Mitchell A, Harries MG, Reeve J. Nutritional and exercise-related determinants of bone density in elite female runners. *Osteoporos Int.* 2004 Aug;15(8):611–8. Epub 2004 Mar 26.

Gleeson M. Nutritional support to maintain proper immune status during intense training. *Nestle Nutr Inst Workshop Ser.* 2013;75:85–97.

Guelfi KJ, Donges CE, Duffield R. Beneficial effects of 12 weeks of aerobic compared with resistance exercise training on perceived appetite in previously sedentary overweight and obese men. *Metabolism.* 2013 Feb;62(2):235–43.

Halson SL. Sleep in elite athletes and nutritional interventions to enhance sleep. *Sports Med.* 2014 May;44 Suppl 1:13–23.

Hasson RE, Howe CA, Jones BL, Freedson PS. Accuracy of four resting metabolic rate prediction equations: effects of sex, body mass index, age, and race/ethnicity. *J Sci Med Sport.* 2011 Jul;14(4):344–51.

Hawley JA, Burke LM. Effect of meal frequency and timing on physical performance. *Br J Nutr.* 1997 Apr;77 Suppl 1:S91–103.

Henry CJ. Basal metabolic rate studies in humans: measurement and development of new equations. *Public Health Nutr.* 2005 Oct;8(7A):1133–52.

Horie LM, Gonzalez MC, Torrinhas RS, Cecconello I, Waitzberg DL. New specific equation to estimate resting energy expenditure in severely obese patients. *Obesity (Silver Spring).* 2011 May;19(5):1090–4.

Horvath PJ, Eagen CK, Fisher NM, Leddy JJ, Pendergast DR. The effects of varying dietary fat on performance and metabolism in trained male and female runners. *J Am Coll Nutr.* 2000 Feb;19(1):52–60.

Ihle R, Loucks AB. Dose-response relationships between energy availability and bone turnover in young exercising women. *J Bone Miner Res.* 2004 Aug;19(8):1231–40.

IOC. Nutrition Working Group of the International Olympic Committee. Nutrition for Athletes. A practical guide to eating for health and performance. Revised and updated 2012.

Jackman MR, Steig A, Higgins JA, Johnson GC, Fleming-Elder BK, Bessesen DH, MacLean PS. Weight regain after sustained weight reduction is accompanied by suppressed oxidation of dietary fat and adipocyte hyperplasia. *Am J Physiol Regul Integr Comp Physiol.* 2008;11:R1117–1129.

Janssens PL, Hursel R, Martens EA, Westerterp-Plantenga MS. Acute effects of capsaicin on energy expenditure and fat oxidation in negative energy balance. *PLoS One.* 2013 Jul 2;8(7):e67786.

Janssens PL, Hursel R, Westerterp-Plantenga MS. Capsaicin increases sensation of fullness in energy balance, and decreases desire to eat after dinner in negative energy balance. *Appetite.* 2014 Jun;77:44–9.

Hagmar M, Berglund B, Brismar K, Hirschberg AL. Body composition and endocrine profile of male Olympic athletes striving for leanness. *Clin J Sport Med.* 2013;11:197–201.

Harper ME, Dent R, Monemdjou S, Bezaire V, Van Wyck L, Wells G, Kavaslar GN, Gauthier A, Tesson F ym. Decreased mitochondrial proton leak and reduced expression of uncoupling protein 3 in skeletal muscle of obese diet-resistant women. *Diabetes.* 2002;11:2459–2466.

Hunter D, Foster M, McArthur JO, Ojha R, Petocz P, Samman S. Evaluation of the micronutrient composition of plant foods produced by organic and conventional agricultural methods. *Crit Rev Food Sci Nutr.* 2011 Jul;51(6):571–82.

Jastroch M, Divakaruni AS, Mookerjee S, Treberg JR, Brand MD. Mitochondrial proton and electron leaks. Essays Biochem. 2010;11:53–67.

Júdice PB, Silva AM, Magalhães JP, Matias CN, Sardinha LB. Sedentary behaviour and adiposity in elite athletes. J Sports Sci. 2014 Jun 10:1–8

Kim B. Thyroid hormone as a determinant of energy expenditure and the basal metabolic rate. Thyroid. 2008;11:141–144.

King NA, Blundell JE. High-fat foods overcome the energy expenditure induced by high-intensity cycling or running. Eur J Clin Nutr. 1995 Feb;49(2):114–23.

Knechtle B, Wirth A, Knechtle P, Rosemann T. Training volume and personal best time in marathon, not anthropometric parameters are associated with performance in male 100-km ultrarunners. J Strength Cond Res. 2010;24:604–609.

Knechtle B, Knechtle P, Barandun U, Rosemann T. Anthropometric and training variables related to half-marathon running performance in recreational female runners. Phys Sportsmed. 2011 May;39(2):158–66.

Knowles SB, Marshall SW, Bowling JM, Loomis D, Millikan R, Yang J, Weaver NL, Kalsbeek W, Mueller FO. A prospective study of injury incidence among North Carolina high school athletes. Am J Epidemiol. 2006 Dec 15;164(12):1209–21.

Legaz-Arrese A, Kinfu H, Munguía-Izquierdo D, Carranza-Garcia LE, Calderón FJ. Basic physiological measures determine fitness and are associated with running performance in elite young male and female Ethiopian runners. J Sports Med Phys Fitness. 2009 Dec;49(4):358–63.

Leibel RL, Rosenbaum M, Hirsch J. Changes in energy expenditure resulting from altered body weight. N Engl J Med. 1995;11:621–628

Levine JA. Non-exercise activity thermogenesis (NEAT) Best Pract Res Clin Endocrinol Metab. 2002;11:679–702.

Loucks AB. Energy balance and body composition in sports and exercise. J Sports Sci. 2004 Jan;22(1):1–14.

Loucks AB. Low energy availability in the marathon and other endurance sports. Sports Med. 2007;37(4–5):348–52.

Loucks AB. Energy availability in athletes. IOC Consensus In Sports Nutrition Conference Lausanne, Switzerland 25th–27th October 2010

Maclean PS, Bergouignan A, Cornier MA, Jackman MR. Biology's response to dieting: the impetus for weight regain. Am J Physiol Regul Integr Comp Physiol. 2011 Sep;301(3):R581–600.

Maestu J, Jurimae J, Valter I, Jurimae T. Increases in ghrelin and decreases in leptin without altering adiponectin during extreme weight loss in male competitive bodybuilders. Metabolism. 2008;11:221–225

Martins C, Morgan LM, Bloom SR, Robertson MD. Effects of exercise on gut peptides, energy intake and appetite. J Endocrinol. 2007 May;193(2):251–8.

Melin A, Tornberg AB, Skouby S, Møller SS, Sundgot-Borgen J, Faber J, Sidelmann JJ, Aziz M, Sjödin A. Energy availability and the female athlete triad in elite endurance athletes. Scand J Med Sci Sports. 2014 May 30. (a)

Melin A, Torstveit MK, Burke L, Marks S, Sundgot-Borgen J. Disordered Eating and Eating Disorders in Aquatic Sports. Int J Sport Nutr Exerc Metab. 2014 Mar 25. (b)

Mifflin MD, St Jeor ST, Hill LA, Scott BJ, Daugherty SA, Koh YO. A new predictive equation for resting energy expenditure in healthy individuals. Am J Clin Nutr. 1990 Feb;51(2):241–7.

Mountjoy M, Sundgot-Borgen J, Burke L, Carter S, Constantini N, Lebrun C, Meyer N, Sherman R, ym. The IOC consensus statement: beyond the Female Athlete Triad--Relative Energy Deficiency in Sport (RED-S). Br J Sports Med. 2014 Apr;48(7):491–7.

Oshima S, Miyauchi S, Kawano H, Ishijima T, Asaka M, Taguchi M, Torii S, Higuchi M. Fat-free mass can be utilized to assess resting energy expenditure for male athletes of different body size. J Nutr Sci Vitaminol (Tokyo). 2011;57(6):394–400.

Owen OE, Kavle E, Owen RS, Polansky M, Caprio S, Mozzoli MA, Kendrick ZV, Bushman MC, Boden G, A reappraisal of caloric requirements in healthy women. Am J Clin Nutr. 1986 Jul;44(1):1–19.

- Owen OE, Holup JL, D'Alessio DA, Craig ES, Polansky M, Smalley KJ, Kavle EC, Bushman MC, ym.. A reappraisal of the caloric requirements of men.** Am J Clin Nutr. 1987 Dec;46(6):875–85.
- Paddon-Jones D, Westman E, Mattes RD, Wolfe RR, Astrup A, Westerterp-Plantenga M.** Protein, weight management, and satiety. Am J Clin Nutr. 2008 May;87(5):1558S-1561S.
- Rosenbaum M, Leibel RL.** Adaptive thermogenesis in humans . Int J Obes. 2010;11(Suppl 1):S47–S55.
- Rossov LM, Fukuda DH, Fahs CA, Loenneke JP, Stout JR.** Natural bodybuilding competition preparation and recovery: a 12-month case study. Int J Sports Physiol Perform. 2013;11:582–592.
- Russel RR, Willis KS, Ravussin E, Larson-Meyer ED.** Effects of endurance running and dietary fat on circulating ghrelin and peptide YY. J Sports Sci Med. 2009 Dec 1;8(4):574–583.
- Saarni SE, Rissanen A, Sarna S, Koskenvuo M, Kaprio J.** Weight cycling of athletes and subsequent weight gain in middleage. Int J Obes (Lond). 2006 Nov;30(11):1639–44.
- Sedeaud A, Marc A, Marck A, Dor F, Schipman J, Dorsey M, Haida A, Berthelot G, Toussaint JF.** BMI, a performance parameter for speed improvement. PLoS One. 2014 Feb 25;9(2):e90183.
- Shimizu K, Aizawa K, Suzuki N, Masuchi K, Okada H, Akimoto T, Mesaki N, Kono I ym.** Influences of weight loss on monocytes and T-cell subpopulations in male judo athletes. J Strength Cond Res. 2011 Jul;25(7):1943–50.
- Simmons PS, Miles JM, Gerich JE, Haymond MW.** Increased proteolysis. An effect of increases in plasma cortisol within the physiologic range. J Clin Invest. 1984;11:412–420.
- Singhal V, Lawson EA, Ackerman KE, Fazeli PK, Clarke H, Lee H, Eddy K, Marengi DA, Derrico NP, Bouxsein ML, Misra M.** Irisin levels are lower in young amenorrheic athletes compared with eumenorrheic athletes and non-athletes and are associated with bone density and strength estimates. PLoS One. 2014 Jun 13;9(6):e100218.
- Skårderud F, Fladvad T, Holmlund H, Garthe I, Engebretsen L.** The malnourished athlete--guidelines for interventions. Tidsskr Nor Laegeforen. 2012 Sep 18;132(17):1944.
- Stellingwerff T.** Nutrition for power sports. IOC Consensus In Sports Nutrition Conference Lausanne, Switzerland 25th–27th October 2010
- Stubbs RJ, Hughes DA, Johnstone AM, Whybrow S, Horgan GW, King N, Blundell J.** Rate and extent of compensatory changes in energy intake and expenditure in response to altered exercise and diet composition in humans. Am J Physiol Regul Integr Comp Physiol. 2004 Feb;286(2):R350–8.
- Sundgot-Borgen J, Torstveit MK.** Prevalence of eating disorders in elite athletes is higher than in the general population. Clin J Sport Med. 2004 Jan;14(1):25–32.
- Sumithran P, Prendergast LA, Delbridge E, Purcell K, Shulkes A, Kriketos A, Proietto J.** Long-term persistence of hormonal adaptations to weight loss. N Engl J Med. 2011;11:1597–1604.
- Taguchi M, Ishikawa-Takata K, Tatsuta W, Katsuragi C, Usui C, Sakamoto S, Higuchi M.** Resting energy expenditure can be assessed by fat-free mass in female athletes regardless of body size. J Nutr Sci Vitaminol (Tokyo). 2011;57(1):22–9.
- Tanda G, Knechtle B.** Marathon performance in relation to body fat percentage and training indices in recreational male runners. Open Access J Sports Med. 2013 May 28;4:141–9.
- Thein-Nissenbaum JM, Rauh MJ, Carr KE, Loud KJ, McGuine TA.** Menstrual irregularity and musculoskeletal injury in female high school athletes. J Athl Train. 2012 Jan-Feb;47(1):74–82.
- Thein-Nissenbaum J.** Long term consequences of the female athlete triad. Maturitas. 2013 Jun;75(2):107–12.
- Thompson JL.** Energy balance in young athletes. Int J Sport Nutr. 1998 Jun;8(2):160–74.
- Tomiyama AJ, Mann T, Vinas D, Hunger JM, DeJager J, Taylor SE.** Low calorie dieting increases cortisol. Psychosom Med. 2010;11:357–364.
- Trexler ET, Smith-Ryan AE1, Norton LE.** Metabolic adaptation to weight loss: implications for the athlete. J Int Soc Sports Nutr. 2014 Feb 27;11(1):7.
- Wardle J, Haase AM, Steptoe A.** Body image and weight control in young adults: international comparisons in university students from 22 countries. Int J Obes (Lond). 2006 Apr;30(4):644–51.

Westerterp KR. Physical activity and physical activity induced energy expenditure in humans: measurement, determinants, and effects. *Front Physiol.* 2013 Apr 26;4:90.

Westerterp-Plantenga MS, Lemmens SG, Westerterp KR. Dietary protein – its role in satiety, energetics, weight loss and health. *Br J Nutr.* 2012 Aug;108 Suppl 2:S105–12.

Weyer C, Walford RL, Harper IT, Milner M, MacCallum T, Tataranni PA, Ravussin E. Energy metabolism after 2 y of energy restriction: the biosphere 2 experiment. *Am J Clin Nutr.* 2000;11:946–953

Yao E, Buchholz AC, Edwards AM, Simpson JA. Predicted and measured resting metabolic rate in young, non-obese women. *Can J Diet Pract Res.* 2013 Fall;74(3):124–30.

Lähdeluettelo luku 2

Bath SC, Rayman MP. Iodine deficiency in the U.K.: an overlooked cause of impaired neurodevelopment? *Proc Nutr Soc.* 2013 May;72(2):226–35.

Dangour AD, Lock K, Hayter A, Aikenhead A, Allen E, Uauy R. Nutrition-related health effects of organic foods: a systematic review. *Am J Clin Nutr.* 2010 Jul;92(1):203–10

Hunter D, Foster M, McArthur JO, Ojha R, Petocz P, Samman S. Evaluation of the micronutrient composition of plant foods produced by organic and conventional agricultural methods. *Crit Rev Food Sci Nutr.* 2011 Jul;51(6):571–82.

Napolitano F, Castellini C, Naspetti S, Piasentier E, Girolami A, Braghieri A. Consumer preference for chicken breast may be more affected by information on organic production than by product sensory properties. *Poult Sci.* 2013 Mar;92(3):820–6.

Palupi E, Jayanegara A, Ploeger A, Kahl J. Comparison of nutritional quality between conventional and organic dairy products: a meta-analysis. *J Sci Food Agric.* 2012 Nov;92(14):2774–81

Smith-Spangler C, Brandeau ML, Hunter GE, Bavinger JC, Pearson M, Eschbach PJ, Sundaram V, Liu H, ym. Are organic foods safer or healthier than conventional alternatives?: a systematic review. *Ann Intern Med.* 2012 Sep 4;157(5):348–66.

Załęcka A, Bügel S, Paoletti F, Kahl J, Bonanno A, Dostalova A, Rahmann G. The influence of organic production on food quality – research findings, gaps and future challenges. *J Sci Food Agric.* 2014 Jan 16.

Zhao X, Chambers E 4th, Matta Z, Loughin TM, Carey EE. Consumer sensory analysis of organically and conventionally grown vegetables. *J Food Sci.* 2007 Mar;72(2):S87–91.

Zimmermann MB. Iodine deficiency. *Endocr Rev.* 2009 Jun;30(4):376–408.

Lähdeluettelo luku 3

Abargouei AS, Janghorbani M, Salehi-Marzijarani M, Esmaillzadeh A. Effect of dairy consumption on weight and body composition in adults: a systematic review and meta-analysis of randomized controlled clinical trials. *Int J Obes (Lond).* 2012 Dec;36(12):1485–93.

Ajala O, English P, Pinkney J. Systematic review and meta-analysis of different dietary approaches to the management of type 2 diabetes. *Am J Clin Nutr.* 2013 Mar;97(3):505–16.

Akinkuolie AO, Ngwa JS, Meigs JB, Djousse L (2011) Omega-3 polyunsaturated fatty acid and insulin sensitivity: a meta-analysis of randomized controlled trials. *Clin Nutr* 30: 702–707

Alexander J, Benford D, Cockburn A, Cravedi JP, Dogliotti, E, Domenico AD, Fernández Cruz ML, ym. Opinion of the Scientific Panel on Contaminants in the Food chain on a request from the European Commission to perform a scientific risk assessment on nitrate in vegetables. *The European Food Safety Authority Journal.* 2008;689:1–79.

Allouche Y, Jiménez A, Gaforio JJ, Uceda M, Beltrán G. How heating affects extra virgin olive oil quality indexes and chemical composition. *J Agric Food Chem.* 2007 Nov 14;55(23):9646–54.

- Anderson JW, Waters AR.** Raisin consumption by humans: effects on glycemia and insulinemia and cardiovascular risk factors. *J Food Sci.* 2013 Jun;78 Suppl 1:A11–7.
- Alhazmi A, Stojanovski E, McEvoy M, Garg ML.** The association between dietary patterns and type 2 diabetes: a systematic review and meta-analysis of cohort studies. *J Hum Nutr Diet.* 2014 Jun;27(3):251–60.
- Alonso A, Zozaya C, Vázquez Z, Alfredo Martínez J, Martínez-González MA.** The effect of low-fat versus whole-fat dairy product intake on blood pressure and weight in young normotensive adults. *J Hum Nutr Diet.* 2009 Aug;22(4):336–42.
- Artero EG, Lee DC, Lavie CJ, España-Romero V, Sui X, Church TS, Blair SN.** Effects of muscular strength on cardiovascular risk factors and prognosis. *J Cardiopulm Rehabil Prev.* 2012 Nov-Dec;32(6):351–8.
- Astrup A.** Yogurt and dairy product consumption to prevent cardiometabolic diseases: epidemiologic and experimental studies. *Am J Clin Nutr.* 2014 May;99(5 Suppl):1235S-42S.
- Astrup A, Pedersen SD.** Is a protein calorie better for weight control? *Am J Clin Nutr.* 2012 Mar;95(3):535–6.
- Aune D, Norat T, Romundstad P, Vatten LJ.** Dairy products and the risk of type 2 diabetes: a systematic review and dose-response meta-analysis of cohort studies. *Am J Clin Nutr.* 2013 Oct;98(4):1066–83.
- Baladia E, Basulto J, Manera M, Martínez R, Calbet D.** Effect of green tea or green tea extract consumption on body weight and body composition: systematic review and meta-analysis. *Nutr Hosp.* 2014 Mar 1;29(3):479–90.
- Bao Y, Han J, Hu FB, Giovannucci EL, Stampfer MJ, Willett WC, Fuchs CS.** Association of nut consumption with total and cause-specific mortality. *N Engl J Med.* 2013 Nov 21;369(21):2001–11
- Barr SI, Rideout CA.** Nutritional considerations for vegetarian athletes. *Nutrition.* 2004 Jul-Aug;20(7–8):696–703.
- Barrack MT, Ackerman KE, Gibbs JC.** Update on the female athlete triad. *Curr Rev Musculoskelet Med.* 2013 Jun;6(2):195–204
- Baxheinrich A, Stratmann B, Lee-Barkey YH, Tschoepe D, Wahrburg U.** Effects of a rapeseed oil-enriched hypoenergetic diet with a high content of α-linolenic acid on body weight and cardiovascular risk profile in patients with the metabolic syndrome. *Br J Nutr.* 2012 Aug;108(4):682–91.
- Bazzano LA, Thompson AM, Tees MT, Nguyen CH, Winham DM.** Non-soy legume consumption lowers cholesterol levels: a meta-analysis of randomized controlled trials. *Nutr Metab Cardiovasc Dis.* 2011 Feb;21(2):94–103.
- Beermann C, Hartung J.** Physiological properties of milk ingredients released by fermentation. *Food Funct.* 2013 Feb;4(2):185–99
- Beja-Pereira A, Luikart G, England PR, Bradley DG, Jann OC, Bertorelle G, Chamberlain AT ym.** Gene-culture coevolution between cattle milk protein genes and human lactase genes. *Nat Genet.* 2003 Dec;35(4):311–3.
- Bell PG, McHugh MP, Stevenson E, Howatson G.** The role of cherries in exercise and health. *Scand J Med Sci Sports.* 2014 Jun;24(3):477–90.
- Bender N, Portmann M, Heg Z, Hofmann K, Zwahlen M, Egger M.** Fish or n3-PUFA intake and body composition: a systematic review and meta-analysis. *Obes Rev.* 2014 May 29.
- Bernstein AM, Sun Q, Hu FB, Stampfer MJ, Manson JE, Willett WC.** Major dietary protein sources and risk of coronary heart disease in women. *Circulation.* 2010 Aug 31;122(9):876–83.
- Biesiekierski JR, Peters SL, Newnham ED, Rosella O, Muir JG, Gibson PR.** No effects of gluten in patients with self-reported non-celiac gluten sensitivity after dietary reduction of fermentable, poorly absorbed, short-chain carbohydrates. *Gastroenterology.* 2013 Aug;145(2):320–8.e1–3.
- Black CD, Herring MP, Hurley DJ, O'Connor PJ.** Ginger (*Zingiber officinale*) reduces muscle pain caused by eccentric exercise. *J Pain.* 2010 Sep;11(9):894–903.
- Boirie Y, Dangin M, Gachon P, Vasson MP, Maubois JL, Beaufre B.** Slow and fast dietary proteins differently modulate postprandial protein accretion. *Proc Natl Acad Sci U S A.* 1997 Dec 23;94(26):14930–5.
- Bonjour JP.** Nutritional disturbance in acid-base balance and osteoporosis: a hypothesis that disregards the essential homeostatic role of the kidney. *Br J Nutr.* 2013 Oct;110(7):1168–77

Bonjour JP, Kraenzlin M, Levasseur R, Warren M, Whiting S. Dairy in adulthood: from foods to nutrient interactions on bone and skeletal muscle health. *J Am Coll Nutr.* 2013;32(4):251–63.

Brandhagen M, Forslund HB, Lissner L, Winkvist A, Lindroos AK, Carlsson LM, Sjöström L, Larsson I. Alcohol and macronutrient intake patterns are related to general and central adiposity.. *Eur J Clin Nutr.* 2012 Mar;66(3):305–13.

Brien SE, Ronksley PE, Turner BJ, Mukamal KJ, Ghali WA. Effect of alcohol consumption on biological markers associated with risk of coronary heart disease: systematic review and meta-analysis of interventional studies. *BMJ.* 2011 Feb 22;342:d636

Bryan NS, Alexander DD, Coughlin JR, Milkowski AL, Boffetta P. Ingested nitrate and nitrite and stomach cancer risk: an updated review. *Food Chem Toxicol.* 2012 Oct;50(10):3646–65.

Buckley JD, Howe PR. Long-chain omega-3 polyunsaturated fatty acids may be beneficial for reducing obesity-a review. *Nutrients.* 2010 Dec;2(12):1212–30.

Buijsse B, Feskens EJ, Schulze MB, Forouhi NG, Wareham NJ, Sharp S, Palli D, Tognon G, Halkjaer J ym. Fruit and vegetable intakes and subsequent changes in body weight in European populations: results from the project on Diet, Obesity, and Genes (DiOGenes). *Am J Clin Nutr.* 2009 Jul;90(1):202–9.

Bøhn SK, Blomhoff R, Paur I. Coffee and cancer risk, epidemiological evidence, and molecular mechanisms. *Mol Nutr Food Res.* 2014 May;58(5):915–30.

Cai L, Ma D, Zhang Y, Liu Z, Wang P. The effect of coffee consumption on serum lipids: a meta-analysis of randomized controlled trials. *Eur J Clin Nutr.* 2012 Aug;66(8):872–7.

Calder PC. Omega-3 polyunsaturated fatty acids and inflammatory processes: nutrition or pharmacology? *Br J Clin Pharmacol.* 2013 Mar;75(3):645–62.

Calder PC, Ahluwalia N, Brouns F, Buetler T, Clement K, Cunningham K, Esposito K, Jönsson LS ym. Dietary factors and low-grade inflammation in relation to overweight and obesity. *Br J Nutr.* 2011 Dec;106 Suppl 3:S5–78.

Caldeira D, Martins C, Alves LB, Pereira H, Ferreira JJ, Costa J. Caffeine does not increase the risk of atrial fibrillation: a systematic review and meta-analysis of observational studies. *Heart.* 2013 Oct;99(19):1383–9.

Carman AJ, Dacks PA, Lane RF, Shineman DW, Fillit HM. Current evidence for the use of coffee and caffeine to prevent age-related cognitive decline and Alzheimer's disease. *J Nutr Health Aging.* 2014 Apr;18(4):383–92.

Carter P, Gray LJ, Troughton J, Khunti K, Davies MJ. Fruit and vegetable intake and incidence of type 2 diabetes mellitus: systematic review and meta-analysis. *BMJ.* 2010 Aug 18;341:c4229.

Casal S, Malheiro R, Sendas A, Oliveira BP, Pereira JA. Olive oil stability under deep-frying conditions. *Food Chem Toxicol.* 2010 Oct;48(10):2972–9.

Catassi C, Bai JC, Bonaz B, Bouma G, Calabò A, Carroccio A, Castillejo G, Ciacci C, Cristofori F, ym. Non-Celiac Gluten sensitivity: the new frontier of gluten related disorders. *Nutrients.* 2013 Sep 26;5(10):3839–53

Cerbulis J, Farrell HM Jr. Composition of milks of dairy cattle. I. Protein, lactose, and fat contents and distribution of protein fraction. *J Dairy Sci.* 1975 Jun;58(6):817–27.

Chan Q, Stamler J, Brown IJ, Daviglus ML, Van Horn L, Dyer AR, Oude Griep LM, Miura K, Ueshima H ym. Relation of raw and cooked vegetable consumption to blood pressure: the INTERMAP Study. *J Hum Hypertens.* 2014 Jun;28(6):353–9.

Chatterton DE, Nguyen DN, Bering SB, Sangild PT. Anti-inflammatory mechanisms of bioactive milk proteins in the intestine of newborns. *Int J Biochem Cell Biol.* 2013 Aug;45(8):1730–47

Chen GC, Lv DB, Pang Z, Liu QF. Red and processed meat consumption and risk of stroke: a meta-analysis of prospective cohort studies. *Eur J Clin Nutr.* 2013 Jan;67(1):91–5

Chen B, Shi HF, Wu SC. Tea consumption didn't modify the risk of fracture: a dose-response meta-analysis of observational studies. *Diagn Pathol.* 2014 Mar 3;9(1):44.

Cheng M, Hu Z, Lu X, Huang J, Gu D. Caffeine intake and atrial fibrillation incidence: dose response meta-analysis of prospective cohort studies. *Can J Cardiol.* 2014 Apr;30(4):448–54.

Childs MT, Dorsett CS, King IB, Ostrander JG, Yamanaka WK. Effects of shellfish consumption on lipoproteins in normolipidemic men. *Am J Clin Nutr.* 1990 Jun;51(6):1020–7.

Cho SS, Qi L, Fahey GC Jr, Klurfeld DM. Consumption of cereal fiber, mixtures of whole grains and bran, and whole grains and risk reduction in type 2 diabetes, obesity, and cardiovascular disease. *Am J Clin Nutr.* 2013 Aug;98(2):594–619.

Choi Y, Song S, Song Y, Lee JE. Consumption of red and processed meat and esophageal cancer risk: meta-analysis. *World J Gastroenterol.* 2013 Feb 21;19(7):1020–9.

Chowdhury R, Stevens S, Gorman D, Pan A, Warnakula S, Chowdhury S, Ward H, Johnson L, Crowe F, ym. Association between fish consumption, long chain omega 3 fatty acids, and risk of cerebrovascular disease: systematic review and meta-analysis. *BMJ.* 2012 Oct 30;345:e6698.

Christensen AS, Viggers L, Hasselström K, Gregersen S. Effect of fruit restriction on glycemic control in patients with type 2 diabetes--a randomized trial. *Nutr J.* 2013 Mar 5;12:29.

Cicerale S, Conlan XA, Barnett NW, Sinclair AJ, Keast RS. Influence of heat on biological activity and concentration of oleocanthal--a natural anti-inflammatory agent in virgin olive oil. *J Agric Food Chem.* 2009 Feb 25;57(4):1326–30.

Claesson AL, Holm G, Ernersson A, Lindström T, Nystrom FH. Two weeks of overfeeding with candy, but not peanuts, increases insulin levels and body weight. *Scand J Clin Lab Invest.* 2009;69(5):598–605.

Clarke R, Frost C, Collins R, Appleby P, Peto R. Dietary lipids and blood cholesterol: quantitative meta-analysis of metabolic ward studies. *BMJ.* 1997 Jan 11;314(7074):112–7.

Cordain L, Eaton SB, Miller JB, Mann N, Hill K. The paradoxical nature of hunter-gatherer diets: meat-based, yet non-atherogenic. *Eur J Clin Nutr.* 2002 Mar;56 Suppl 1:S42–52. Review.

Costanzo S, Di Castelnuovo A, Donati MB, Iacoviello L, de Gaetano G. Wine, beer or spirit drinking in relation to fatal and non-fatal cardiovascular events: a meta-analysis. *Eur J Epidemiol.* 2011 Nov;26(11):833–50.

Cross GA, Fung DY. The effect of microwaves on nutrient value of foods. *Crit Rev Food Sci Nutr.* 1982;16(4):355–81. Review.

Davis JM, Murphy EA, Carmichael MD, Zielinski MR, Groschwitz CM, Brown AS, Gangemi JD, Ghaffar A, ym. Curcumin effects on inflammation and performance recovery following eccentric exercise-induced muscle damage. *Am J Physiol Regul Integr Comp Physiol.* 2007 Jun;292(6):R2168–73.

de Oliveira MC, Sichieri R, Venturim Mozzer R. A low-energy-dense diet adding fruit reduces weight and energy intake in women. *Appetite.* 2008 Sep;51(2):291–5.

De Oliveira e Silva ER, Seidman CE, Tian JJ, Hudgins LC, Sacks FM, Breslow JL. Effects of shrimp consumption on plasma lipoproteins. *Am J Clin Nutr.* 1996 Nov;64(5):712–7.

de Oliveira Otto MC, Mozaffarian D, Kromhout D, Bertoni AG, Sibley CT, Jacobs DR Jr, Nettleton JA. Dietary intake of saturated fat by food source and incident cardiovascular disease: the Multi-Ethnic Study of Atherosclerosis. *Am J Clin Nutr.* 2012 Aug;96(2):397–404.

Dias CB, Garg R, Wood LG, Garg ML. Saturated fat consumption may not be the main cause of increased blood lipid levels. *Med Hypotheses.* 2014 Feb;82(2):187–95.

Dideriksen K, Reitelseder S, Holm L. Influence of amino acids, dietary protein, and physical activity on muscle mass development in humans. *Nutrients.* 2013 Mar 13;5(3):852–76.

Ding M, Bhupathiraju SN, Chen M, van Dam RM, Hu FB. Caffeinated and decaffeinated coffee consumption and risk of type 2 diabetes: a systematic review and a dose-response meta-analysis. *Diabetes Care.* 2014 Feb;37(2):569–86.(a)

Ding M, Bhupathiraju SN, Satija A, van Dam RM, Hu FB. Long-term coffee consumption and risk of cardiovascular disease: a systematic review and a dose-response meta-analysis of prospective cohort studies. *Circulation.* 2014 Feb 11;129(6):643–59. (b)

Dinstel RR, Cascio J, Koukel S. The antioxidant level of Alaska's wild berries: high, higher and highest. *Int J Circumpolar Health.* 2013 Aug 5;72.

Dougkas A, Reynolds CK, Givens ID, Elwood PC, Minihane AM. Associations between dairy consumption and body weight: a review of the evidence and underlying mechanisms. *Nutr Res Rev.* 2011 Jun;24(1):72–95.

Dreher ML, Davenport AJ. Hass avocado composition and potential health effects. *Crit Rev Food Sci Nutr.* 2013;53(7):738–50

Du H, van der A DL, Boshuizen HC, Forouhi NG, Wareham NJ, Halkjaer J, Tjønneland A, Overvad K ym. Dietary fiber and subsequent changes in body weight and waist circumference in European men and women. *Am J Clin Nutr.* 2010 Feb;91(2):329–36.

EFSA. Nitrate in vegetables Scientific Opinion of the Panel on Contaminants in the Food chain 1 (Question No EFSA-Q-2006-071). *The EFSA Journal* (2008) 689, 1–79

EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA) (2011). Scientific Opinion on the substantiation of health claims related to caffeine and increased fat oxidation leading to a reduction in body fat mass (ID 735, 1484), increased energy expenditure leading to a reduction in body weight (ID 1487), increased alertness (ID 736, 1101, 1187, 1485, 1491, 2063, 2103) and increased attention (ID 736, 1485, 1491, 2375) pursuant to Article 13(1) of Regulation (EC) No 1924/2006. *EFSA Journal*, 9(4):2054.

EFSA. Scientific Opinion on Dietary Reference Values for energy *EFSA Journal* 2013;11(1):3005 [112 pp.]

EFSA. The 2010 European Union Report on Pesticide Residues in Food. *EFSA Journal* 2013;11(3):3130 [808 pp.]

Eftekhari MH, Mozaffari-Khosravi H, Shidfar F, Zamani A. Relation between Body Iron Status and Cardiovascular Risk Factors in Patients with Cardiovascular Disease. *Int J Prev Med.* 2013 Aug;4(8):911–6.

Ehnholm C, Lukka M, Kuusi T, Nikkilä E, Utermann G. Apolipoprotein E polymorphism in the Finnish population: gene frequencies and relation to lipoprotein concentrations. *J Lipid Res.* 1986 Mar;27(3):227–35.

Elliott P, Stamler J, Dyer AR, Appel L, Dennis B, Kesteloot H, Ueshima H, Okayama A, Chan Q ym. Association between protein intake and blood pressure: the INTERMAP Study. *Arch Intern Med.* 2006 Jan 9;166(1):79–87.

Elwood PC, Pickering JE, Givens DI, Gallacher JE. The consumption of milk and dairy foods and the incidence of vascular disease and diabetes: an overview of the evidence. *Lipids.* 2010 Oct;45(10):925–39.

Engle-Stone R, Yeung A, Welch R, Glahn R. Meat and ascorbic acid can promote Fe availability from Fe-phytate but not from Fe-tannic acid complexes. *J Agric Food Chem.* 2005 Dec 28;53(26):10276–84.

Eslick GD, Howe PR, Smith C, Priest R, Bensoussan A. Benefits of fish oil supplementation in hyperlipidemia: a systematic review and meta-analysis. *Int J Cardiol.* 2009 Jul 24;136(1):4–16.

Esposito K, Kastorini CM, Panagiotakos DB, Giugliano D. Mediterranean diet and weight loss: meta-analysis of randomized controlled trials. *Metab Syndr Relat Disord.* 2011 Feb;9(1):1–12.

Estruch R, Salas-Salvadó J. “Towards an even healthier Mediterranean diet”. *Nutr Metab Cardiovasc Dis.* 2013 Dec;23(12):1163–6.

Estruch R, Ros E, Salas-Salvadó J, Covas MI, Corella D, Arós F, Gómez-Gracia E, Ruiz-Gutiérrez V ym. Primary prevention of cardiovascular disease with a Mediterranean diet. *N Engl J Med.* 2013 Apr 4;368(14):1279–90.

Evira. Ravinnon kasvinsuojeluainejäämät – kumulatiivinen riskinarvointi. *Eviran tutkimuksia* 3/2010.

Evira. Kvantiatiivinen riskinarvointi lasten ja aikuisten altistumisesta nitraatille ja nitriitille. *Eviran tutkimuksia* 2/2013 (a).

Evira. <http://www.evira.fi/portal/fi/elintarvikkeet/tietoa%2Belintarvikkeista/elintarvikkearat/elintarvikkeiden%2Bkayton%2Brajoituksset/kalan%2Bsyontisuositukset/> 2013 (b)

Fenton TR, Eliasziw M, Tough SC, Lyon AW, Brown JP, Hanley DA. Low urine pH and acid excretion do not predict bone fractures or the loss of bone mineral density: a prospective cohort study. *BMC Musculoskelet Disord.* 2010 May 10;11:88.

Fenton TR, Tough SC, Lyon AW, Eliasziw M, Hanley DA. Causal assessment of dietary acid load and bone disease: a systematic review & meta-analysis applying Hill's epidemiologic criteria for causality. *Nutr J.* 2011 Apr 30;10:41.

Fernandez ML. Dietary cholesterol provided by eggs and plasma lipoproteins in healthy populations. *Curr Opin Clin Nutr Metab Care.* 2006 Jan;9(1):8–12. Review.

Fernandez-Cao JC, Arija V1, Aranda N, Bullo M, Basora J, Martínez-González MA, Díez-Espino J, ym. Heme iron intake and risk of new-onset diabetes in a Mediterranean population at high risk of cardiovascular disease: an observational cohort analysis. *BMC Public Health*. 2013 Nov 4;13:1042.

Ferracane R, Pellegrini N, Visconti A, Graziani G, Chiavaro E, Miglio C, Fogliano V. Effects of different cooking methods on antioxidant profile, antioxidant capacity, and physical characteristics of artichoke. *J Agric Food Chem*. 2008 Sep 24;56(18):8601–8.

Feskens EJ, Sluik D, van Woudenberg GJ. Meat consumption, diabetes, and its complications. *Curr Diab Rep*. 2013 Apr;13(2):298–306.

Finravinto 2012 -tutkimus. Helldán A, Raulio S, Kosola M, Tapanainen H, Ovaskainen M-L, Virtanen S. THL Raportti 16/2013. Juvenes Print – Suomen Yliopistopaino Oy. Tampere 2013.

Flores-Mateo G, Rojas-Rueda D, Basora J, Ros E, Salas-Salvadó J. Nut intake and adiposity: meta-analysis of clinical trials. *Am J Clin Nutr*. 2013 Jun;97(6):1346–55.

Fogelholm M, Anderssen S, Gunnarsdottir I, Lahti-Koski M. Dietary macronutrients and food consumption as determinants of long-term weight change in adult populations: a systematic literature review. *Food Nutr Res*. 2012;56.

Frassetto LA, Schloetter M, Mietus-Synder M, Morris RC Jr, Sebastian A. Metabolic and physiologic improvements from consuming a paleolithic, hunter-gatherer type diet. *Eur J Clin Nutr*. 2009 Aug;63(8):947–55.

Fulgoni VL 3rd, Dreher M, Davenport AJ. Avocado consumption is associated with better diet quality and nutrient intake, and lower metabolic syndrome risk in US adults: results from the National Health and Nutrition Examination Survey (NHANES) 2001–2008. *Nutr J*. 2013 Jan 2;12:1.

Fung TT, van Dam RM, Hankinson SE, Stampfer M, Willett WC, Hu FB. Low-carbohydrate diets and all-cause and cause-specific mortality: two cohort studies. *Ann Intern Med*. 2010 Sep 7;153(5):289–98.

Gao D, Ning N, Wang C, Wang Y, Li Q, Meng Z, Liu Y, Li Q. Dairy products consumption and risk of type 2 diabetes: systematic review and dose-response meta-analysis. *PLoS One*. 2013 Sep 27;8(9):e73965.

Gautam S, Platel K, Srinivasan K. Influence of combinations of promoter and inhibitor on the bioaccessibility of iron and zinc from food grains. *Int J Food Sci Nutr*. 2011 Dec;62(8):826–34.

Glatter KA, Myers R, Chiamvimonvat N. Recommendations regarding dietary intake and caffeine and alcohol consumption in patients with cardiac arrhythmias: what do you tell your patients to do or not to do? *Curr Treat Options Cardiovasc Med*. 2012 Oct;14(5):529–35.

Gliszczynska-Swiglo A, Ciska E, Pawlak-Lema ska K, Chmielewski J, Borkowski T, Tyrakowska B. Changes in the content of health-promoting compounds and antioxidant activity of broccoli after domestic processing. *Food Addit Contam*. 2006 Nov;23(11):1088–98.

Gobbetti M, Rizzello CG, Di Cagno R, De Angelis M. How the sourdough may affect the functional features of leavened baked goods. *Food Microbiol*. 2014 Feb;37:30–40.

Gray P, Gabriel B, Thies F, Gray SR. Fish oil supplementation augments post-exercise immune function in young males. *Brain Behav Immun*. 2012 Nov;26(8):1265–72.

Gray P, Chappell A, Jenkinson AM, Thies F, Gray SR. Fish oil supplementation reduces markers of oxidative stress but not muscle soreness after eccentric exercise. *Int J Sport Nutr Exerc Metab*. 2014 Apr;24(2):206–14.

Gregersen NT, Belza A, Jensen MG, Ritz C, Bitz C, Hels O, Frandsen E, Mela DJ, Astrup A. Acute effects of mustard, horseradish, black pepper and ginger on energy expenditure, appetite, ad libitum energy intake and energy balance in human subjects. *Br J Nutr*. 2013 Feb 14;109(3):556–63

Griffin JD, Lichtenstein AH. Dietary Cholesterol and Plasma Lipoprotein Profiles: Randomized-Controlled Trials. *Curr Nutr Rep*. 2013 Dec;2(4):274–282.

Guasch-Ferré M, Bulló M, Martínez-González MÁ, Ros E, Corella D, Estruch R ym; PREDIMED study group. Frequency of nut consumption and mortality risk in the PREDIMED nutrition intervention trial. *BMC Med*. 2013 Jul 16;11:164.

Guasch-Ferré M, Hu FB, Martínez-González MA, Fitó M, Bulló M, Estruch R, Ros E, Corella D, Recondo J ym. Olive oil intake and risk of cardiovascular disease and mortality in the PREDIMED Study. *BMC Med.* 2014 May 13;12(1):78.

Halkjær J, Olsen A, Overvad K, Jakobsen MU, Boeing H, Buijsse B, Palli D, Tognon G, Du H, van der A DL ym. Intake of total, animal and plant protein and subsequent changes in weight or waist circumference in European men and women: the Diogenes project. *Int J Obes (Lond)*. 2011 Aug;35(8):1104–13.

Harman NL, Leeds AR, Griffin BA. Increased dietary cholesterol does not increase plasma low density lipoprotein when accompanied by an energy-restricted diet and weight loss. *Eur J Nutr.* 2008 Sep;47(6):287–93.

Harris RC, Wise JA, Price KA, Kim HJ, Kim CK, Sale C. Determinants of muscle carnosine content. *Amino Acids.* 2012 Jul;43(1):5–12.

Hartley L, Igbinedion E, Holmes J, Flowers N, Thorogood M, Clarke A, Stranges S, Hooper L, Rees K. Increased consumption of fruit and vegetables for the primary prevention of cardiovascular diseases. *Cochrane Database Syst Rev.* 2013 Jun 4;6:CD009874.

Haub MD, Wells AM, Tarnopolsky MA, Campbell WW. Effect of protein source on resistive-training-induced changes in body composition and muscle size in older men. *Am J Clin Nutr.* 2002 Sep;76(3):511–7.

Hauner H, Bechthold A, Boeing H, Brönstrup A, Buyken A, Leschik-Bonnet E ym. ; **German Nutrition Society.** Evidence-based guideline of the German Nutrition Society: carbohydrate intake and prevention of nutrition-related diseases. *Ann Nutr Metab.* 2012;60 Suppl 1:1–58

He K. Fish, long-chain omega-3 polyunsaturated fatty acids and prevention of cardiovascular disease--eat fish or take fish oil supplement? *Prog Cardiovasc Dis.* 2009 Sep-Oct;52(2):95–114.

He FJ, Nowson CA, Lucas M, MacGregor GA. Increased consumption of fruit and vegetables is related to a reduced risk of coronary heart disease: meta-analysis of cohort studies. *J Hum Hypertens.* 2007 Sep;21(9):717–28.

Hill AM, Buckley JD, Murphy KJ, Howe PR. Combining fish-oil supplements with regular aerobic exercise improves body composition and cardiovascular disease risk factors. *Am J Clin Nutr.* 2007 May;85(5):1267–74.

Hodgson AB, Randell RK, Jeukendrup AE. The metabolic and performance effects of caffeine compared to coffee during endurance exercise. *PLoS One.* 2013;8(4):e59561.

Hoffman CJ, Zabik ME. Effects of microwave cooking/reheating on nutrients and food systems: a review of recent studies. *J Am Diet Assoc.* 1985 Aug;85(8):922–6. Review.

Hu FB, Stampfer MJ, Rimm EB, Manson JE, Ascherio A, Colditz GA, Rosner BA, Spiegelman D ym. A prospective study of egg consumption and risk of cardiovascular disease in men and women. *JAMA.* 1999 Apr 21;281(15):1387–94.

Huhtinen H, Rimpelä A. Nuorison energiajuomien käyttö ja kofeiiniin liittyvät oireet. *Suomen Lääkärilehti* 2013;68(39):2451–2455.

Huncharek M, Muscat J, Kupelnick B. Impact of dairy products and dietary calcium on bone-mineral content in children: results of a meta-analysis. *Bone.* 2008 Aug;43(2):312–21

Hursel R, Westerterp-Plantenga MS. Catechin- and caffeine-rich teas for control of body weight in humans. *Am J Clin Nutr.* 2013 Dec;98(6 Suppl):1682S–1693S.

Hurst S, Zainal Z, Caterson B, Hughes CE, Harwood JL. Dietary fatty acids and arthritis. *Prostaglandins Leukot Essent Fatty Acids.* 2010 Apr-Jun;82(4–6):315–8.

Huth PJ, Park KM. Influence of dairy product and milk fat consumption on cardiovascular disease risk: a review of the evidence. *Adv Nutr.* 2012 May 1;3(3):266–85.

Hwang ES, Kim GH. Effects of various heating methods on glucosinolate, carotenoid and tocopherol concentrations in broccoli. *Int J Food Sci Nutr.* 2013 Feb;64(1):103–11.

Jackson CL, Hu FB. Long-term associations of nut consumption with body weight and obesity. *Am J Clin Nutr.* 2014 Jun 4.

Jakulj F, Zernicke K, Bacon SL, van Wielingen LE, Key BL, West SG, Campbell TS. A high-fat meal increases cardiovascular reactivity to psychological stress in healthy young adults. *J Nutr.* 2007 Apr;137(4):935–9.

Janssens PL, Hursel R, Martens EA, Westerterp-Plantenga MS. Acute effects of capsaicin on energy expenditure and fat oxidation in negative energy balance. *PLoS One.* 2013 Jul 2;8(7):e67786.

Janssens PL, Hursel R, Westerterp-Plantenga MS. Capsaicin increases sensation of fullness in energy balance, and decreases desire to eat after dinner in negative energy balance. *Appetite.* 2014 Jun;77:44–9.

Je Y, Giovannucci E. Coffee consumption and total mortality: a meta-analysis of twenty prospective cohort studies. *Br J Nutr.* 2014 Apr 14;111(7):1162–73.

Jiang PY, Jiang ZB, Shen KX, Yue Y. Fish intake and ovarian cancer risk: a meta-analysis of 15 case-control and cohort studies. *PLoS One.* 2014 Apr 14;9(4):e94601. (a)

Jiang X, Zhang D, Jiang W. Coffee and caffeine intake and incidence of type 2 diabetes mellitus: a meta-analysis of prospective studies. *Eur J Nutr.* 2014 Feb;53(1):25–38. (b)

Jiménez-Monreal AM, García-Diz L, Martínez-Tomé M, Mariscal M, Murcia MA. Influence of cooking methods on antioxidant activity of vegetables. *J Food Sci.* 2009 Apr;74(3):H97-H103.

Jin M, Cai S, Guo J, Zhu Y, Li M, Yu Y, Zhang S, Chen K. Alcohol drinking and all cancer mortality: a meta-analysis. *Ann Oncol.* 2013 Mar;24(3):807–16.

Johnson GH, Fritzsche K. Effect of dietary linoleic acid on markers of inflammation in healthy persons: a systematic review of randomized controlled trials. *J Acad Nutr Diet.* 2012 Jul;112(7):1029–41, 1041.e1–15.

Johnson CM, Wei C, Ensor JE, Smolenski DJ, Amos CI, Levin B, Berry DA. Meta-analyses of colorectal cancer risk factors. *Cancer Causes Control.* 2013 Jun;24(6):1207–22.

Jones PJ. Dietary cholesterol and the risk of cardiovascular disease in patients: a review of the Harvard Egg Study and other data. *Int J Clin Pract Suppl.* 2009 Oct;(163):1–8, 28–36.

Jones G. Early life nutrition and bone development in children. *Nestle Nutr Workshop Ser Pediatr Program.* 2011;68:227–33; discussion 233–6.

Jones AM, Bailey SJ, Vanhatalo A, Fulford J, Gilchrist M, Benjamin N, Winyard PG. Reply to Lundberg, Larsen, and Weitzberg. *J Appl Physiol (1985).* 2011 Aug;111(2):619

Jones AM. Dietary nitrate supplementation and exercise performance. *Sports Med.* 2014 May;44 Suppl 1:S35–45.

Josse AR, Phillips SM. Impact of milk consumption and resistance training on body composition of female athletes. *Med Sport Sci.* 2012;59:94–103.

Jouris KB, McDaniel JL, Weiss EP. The Effect of Omega-3 Fatty Acid Supplementation on the Inflammatory Response to eccentric strength exercise. *J Sports Sci Med.* 2011 Sep 1;10(3):432–8. eCollection 2011.

Jurgens TM, Whelan AM, Killian L, Doucette S, Kirk S, Foy E. Green tea for weight loss and weight maintenance in overweight or obese adults. *Cochrane Database Syst Rev.* 2012 Dec 12;12:CD008650.

Jönsson T, Granfeldt Y, Ahrén B, Branell UC, Pålsson G, Hansson A, Söderström M, Lindeberg S. Beneficial effects of a Paleolithic diet on cardiovascular risk factors in type 2 diabetes: a randomized cross-over pilot study. *Cardiovasc Diabetol.* 2009 Jul 16;8:35.

Jönsson T, Granfeldt Y, Erlanson-Albertsson C, Ahrén B, Lindeberg S. A paleolithic diet is more satiating per calorie than a mediterranean-like diet in individuals with ischemic heart disease. *Nutr Metab (Lond).* 2010 Nov 30;7:85.

Jönsson T, Granfeldt Y, Lindeberg S, Hallberg AC. Subjective satiety and other experiences of a Paleolithic diet compared to a diabetes diet in patients with type 2 diabetes. *Nutr J.* 2013 Jul 29;12:105.

Kai SH, Bongard V, Simon C, Ruidavets JB, Arveiler D, Dallongeville J, Wagner A, Amouyel P, Ferrières J. Low-fat and high-fat dairy products are differently related to blood lipids and cardiovascular risk score. *Eur J Prev Cardiol.* 2013 Sep 3.

Kalergis M, Leung Yinko SS, Nedelcu R. Dairy products and prevention of type 2 diabetes: implications for research and practice. *Front Endocrinol (Lausanne).* 2013 Jul 23;4:90.

Kanerva N, Kaartinen NE, Ovaskainen ML, Konttinen H, Kontto J, Männistö S. A diet following Finnish nutrition recommendations does not contribute to the current epidemic of obesity. *Public Health Nutr.* 2013 May;16(5):786–94. (a)

Kanerva N, Kaartinen NE, Schwab U, Lahti-Koski M, Männistö S. Adherence to the Baltic Sea diet consumed in the Nordic countries is associated with lower abdominal obesity. *Br J Nutr.* 2013 Feb 14;109(3):520–8. (b)

Kanerva N, Loo BM, Eriksson JG, Leiviskä J, Kaartinen NE, Jula A, Männistö S. Associations of the Baltic Sea diet with obesity-related markers of inflammation. *Ann Med.* 2014 Mar;46(2):90–6 (a)

Kanerva N, Kaartinen NE, Rissanen H, Knekt P, Eriksson JG, Sääksjärvi K, Sundvall J, Männistö S. Associations of the Baltic Sea diet with cardiometabolic risk factors – a meta-analysis of three Finnish studies. *Br J Nutr.* 2014 May 27:1–11. (b)

Killer SC, Blannin AK, Jeukendrup AE. No evidence of dehydration with moderate daily coffee intake: a counterbalanced cross-over study in a free-living population. *PLoS One.* 2014 Jan 9;9(1):e84154.

Klein BP. Retention of nutrients in microwave-cooked foods. *Bol Asoc Med P R.* 1989 Jul;81(7):277–9.

Knight C. “Most people are simply not designed to eat pasta”: evolutionary explanations for obesity in the low-carbohydrate diet movement. *Public Underst Sci.* 2011 Sep;20(5):706–19.

Kokubo Y, Iso H, Saito I, Yamagishi K, Yatsuya H, Ishihara J, Inoue M, Tsugane S. The impact of green tea and coffee consumption on the reduced risk of stroke incidence in Japanese population: the Japan public health center-based study cohort. *Stroke.* 2013 May;44(5):1369–74.

Kolb H, Mandrup-Poulsen T. The global diabetes epidemic as a consequence of lifestyle-induced low-grade inflammation. *Diabetologia.* 2010 Jan;53(1):10–20.

Koloverou E, Esposito K, Giugliano D, Panagiotakos D. The effect of Mediterranean diet on the development of type 2 diabetes mellitus: A meta-analysis of 10 prospective studies and 136,846 participants. *Metabolism.* 2014 Jul;63(7):903–911.

Kratz M, Baars T, Guyenet S. The relationship between high-fat dairy consumption and obesity, cardiovascular, and metabolic disease. *Eur J Nutr.* 2013 Feb;52(1):1–24.

Kratz M, Marcovina S, Nelson JE, Yeh MM, Kowdley KV, Callahan HS, Song X, Di C, Utzschneider KM. Dairy fat intake is associated with glucose tolerance, hepatic and systemic insulin sensitivity, and liver fat but not -cell function in humans. *Am J Clin Nutr.* 2014 Apr 16;99(6):1385–1396.

Krishnan S, Cooper JA. Effect of dietary fatty acid composition on substrate utilization and body weight maintenance in humans. *Eur J Nutr.* 2014 Apr;53(3):691–710.

Kromhout D, de Goede J. Update on cardiometabolic health effects of -3 fatty acids. *Curr Opin Lipidol.* 2014 Feb;25(1):85–90.

Kuehl KS. Cherry juice targets antioxidant potential and pain relief. *Med Sport Sci.* 2012;59:86–93.

Kunutsor SK, Apekey TA, Walley J, Kain K. Ferritin levels and risk of type 2 diabetes mellitus: an updated systematic review and meta-analysis of prospective evidence. *Diabetes Metab Res Rev.* 2013 May;29(4):308–18.

Lacroix M, Bos C, Léonil J, Airinei G, Luengo C, Daré S, Benamouzig R, Fouillet H, Fauquant J, Tomé D ym. Compared with casein or total milk protein, digestion of milk soluble proteins is too rapid to sustain the anabolic postprandial amino acid requirement. *Am J Clin Nutr.* 2006 Nov;84(5):1070–9.

Lappi J. Effects of wholegrain foods and grain fibre in intestinal tract in relation to glucose metabolism : with an emphasis on wheat and rye bread effects. University of Eastern Finland, Faculty of Health Sciences. Publications of the University of Eastern Finland. Dissertations in Health Sciences. 227. 2014 p. 140

Lee DR, Lee J, Rota M, Lee J, Ahn HS, Park SM, Shin D. Coffee consumption and risk of fractures: a systematic review and dose-response meta-analysis. *Bone.* 2014 Jun;63:20–8.

Leung Yinko SS, Stark KD, Thanassoulis G, Pilote L. Fish Consumption and Acute Coronary Syndrome:A Meta-Analysis. *Am J Med.* 2014 May;3.

Li Y, Zhou C, Zhou X, Li L. Egg consumption and risk of cardiovascular diseases and diabetes: a meta-analysis. *Atherosclerosis.* 2013 Aug;229(2):524–30

Lindeberg S, Jönsson T, Granfeldt Y, Borgstrand E, Soffman J, Sjöström K, Ahrén B. A Palaeolithic diet improves glucose tolerance more than a Mediterranean-like diet in individuals with ischaemic heart disease. *Diabetologia.* 2007 Sep;50(9):1795–807.

- Liu RH.** Health-promoting components of fruits and vegetables in the diet. *Adv Nutr.* 2013 May;4(3):384S-92S.
- Liu H, Yao K, Zhang W, Zhou J, Wu T, He C.** Coffee consumption and risk of fractures: a meta-analysis. *Arch Med Sci.* 2012 Nov;8(5):776–83.
- Lomer MC, Parkes GC, Sanderson JD.** Review article: lactose intolerance in clinical practice--myths and realities. *Aliment Pharmacol Ther.* 2008 Jan;27(2):93–103.
- Lucas L, Russell A, Keast R.** Molecular mechanisms of inflammation. Anti-inflammatory benefits of virgin olive oil and the phenolic compound oleocanthal. *Curr Pharm Des.* 2011;17(8):754–68. Review.
- Lundberg JO, Larsen FJ, Weitzberg E.** Supplementation with nitrate and nitrite salts in exercise: a word of caution. *J Appl Physiol* (1985). 2011 Aug;111(2):616–7.
- Lundin KE, Alaeddini A.** Non-celiac gluten sensitivity. *Gastrointest Endosc Clin N Am.* 2012 Oct;22(4):723–34.
- Lönnadal B.** Calcium and iron absorption--mechanisms and public health relevance. *Int J Vitam Nutr Res.* 2010 Oct;80(4–5):293–9.
- Ma JK, Zhang H, Tsuchiya T, Akiyama Y, Chen JY.** Frying stability of rapeseed Kizakinonatane (*Brassica napus*) oil in comparison with canola oil. *Food Sci Technol Int.* 2014 Jan 28.
- Macaluso F, Barone R, Catanese P, Carini F, Rizzuto L, Farina F, Di Felice V.** Do fat supplements increase physical performance? *Nutrients.* 2013 Feb 7;5(2):509–24.
- Machha A, Schechter AN.** Inorganic nitrate: a major player in the cardiovascular health benefits of vegetables? *Nutr Rev.* 2012 Jun;70(6):367–72.
- Mackey AL.** Does an NSAID a day keep satellite cells at bay? *J Appl Physiol* (1985). 2013 Sep;115(6):900–8.
- Madero M, Arriaga JC, Jalal D, Rivard C, McFann K, Pérez-Méndez O, Vázquez A, Ruiz A, Lanasa MA, ym.** The effect of two energy-restricted diets, a low-fructose diet versus a moderate natural fructose diet, on weight loss and metabolic syndrome parameters: a randomized controlled trial. *Metabolism.* 2011 Nov;60(11):1551–9.
- Magalhães B, Peleteiro B, Lunet N.** Dietary patterns and colorectal cancer: systematic review and meta-analysis. *Eur J Cancer Prev.* 2012 Jan;21(1):15–23.
- Manore MM.** Dietary supplements for improving body composition and reducing body weight: where is the evidence? *Int J Sport Nutr Exerc Metab.* 2012 Apr;22(2):139–54. Review.
- Mannerheim Lastensuojeluliitto & Suomen Sydänliitto. Pusa T, Kara A, Pietiläinen K.** "Ensimmäinen ruokavuosi".
- Mariani J, Doval HC, Nul D, Varini S, Grancelli H, Ferrante D, Tognoni G, Macchia A.** N-3 polyunsaturated fatty acids to prevent atrial fibrillation: updated systematic review and meta-analysis of randomized controlled trials. *J Am Heart Assoc.* 2013 Feb 19;2(1):e005033..
- Marinangeli CP, Jones PJ.** Pulse grain consumption and obesity: effects on energy expenditure, substrate oxidation, body composition, fat deposition and satiety. *Br J Nutr.* 2012 Aug;108 Suppl 1:S46–51.
- Martínez-González MA, Toledo E, Arós F, Fiol M, Corella D, Salas-Salvadó J, Ros E, Covas MI ym.** Extra-Virgin Olive Oil Consumption Reduces Risk of Atrial Fibrillation: The PREDIMED Trial. *Circulation.* 2014 Apr 30
- Martínez-Lapiscina EH, Clavero P, Toledo E, Estruch R, Salas-Salvadó J, San Julián B, Sanchez-Tainta A ym.** Mediterranean diet improves cognition: the PREDIMED-NAVARRA randomised trial. *J Neurol Neurosurg Psychiatry.* 2013 Dec;84(12):1318–25.
- Mashhadi NS, Ghiasvand R, Askari G, Feizi A, Hariri M, Darvishi L, Barani A, Taghiyar M, Shiranian A, ym.** Influence of ginger and cinnamon intake on inflammation and muscle soreness endured by exercise in Iranian female athletes. *Int J Prev Med.* 2013 Apr;4(Suppl 1):S11–5.
- Massey LK.** Dairy food consumption, blood pressure and stroke. *J Nutr.* 2001 Jul;131(7):1875–8. Review.
- Matthews SB, Waud JP, Roberts AG, Campbell AK.** Systemic lactose intolerance: a new perspective on an old problem. *Postgrad Med J.* 2005 Mar;81(953):167–73. Review.
- McClung JP.** Iron status and the female athlete. *J Trace Elem Med Biol.* 2012 Jun;26(2–3):124–6.

McEvoy CT, Temple N, Woodside JV. Vegetarian diets, low-meat diets and health: a review. *Public Health Nutr.* 2012 Dec;15(12):2287–94.

McGartland CP, Robson PJ, Murray LJ, Cran GW, Savage MJ, Watkins DC, Rooney MM, Boreham CA. Fruit and vegetable consumption and bone mineral density: the Northern Ireland Young Hearts Project. *Am J Clin Nutr.* 2004 Oct;80(4):1019–23.

McKeown NM, Troy LM, Jacques PF, Hoffmann U, O'Donnell CJ, Fox CS. Whole- and refined-grain intakes are differentially associated with abdominal visceral and subcutaneous adiposity in healthy adults: the Framingham Heart Study. *Am J Clin Nutr.* 2010 Nov;92(5):1165–71.

Mensinga TT, Speijers GJ, Meulenbelt J. Health implications of exposure to environmental nitrogenous compounds. *Toxicol Rev.* 2003;22(1):41–51.

Micha R, Michas G, Lajous M, Mozaffarian D. Processing of meats and cardiovascular risk: time to focus on preservatives. *BMC Med.* 2013 May 23;11:136.

Mickleborough TD, Head SK, Lindley MR. Exercise-induced asthma: nutritional management. *Curr Sports Med Rep.* 2011 Jul;10(4):197–202.

Miglio C, Chiavaro E, Visconti A, Fogliano V, Pellegrini N. Effects of different cooking methods on nutritional and physicochemical characteristics of selected vegetables. *J Agric Food Chem.* 2008 Jan 9;56(1):139–47

Minihane AM, Khan S, Leigh-Firbank EC, Talmud P, Wright JW, Murphy MC, Griffin BA, Williams CM: ApoE polymorphism and fish oil supplementation in subjects with an atherogenic lipoprotein phenotype. *Arterioscler Thromb Vasc Biol* 2000;20:1990–1997.

Mirvish SS, Grandjean AC, Reimers KJ, Connelly BJ, Chen SC, Morris CR, Wang X, Haorah J, Lyden ER. Effect of ascorbic acid dose taken with a meal on nitrosoproline excretion in subjects ingesting nitrate and proline. *Nutr Cancer.* 1998;31(2):106–10.

Misiak B, Leszek J, Kiejna A. Metabolic syndrome, mild cognitive impairment and Alzheimer's disease--the emerging role of systemic low-grade inflammation and adiposity. *Brain Res Bull.* 2012 Nov 1;89(3–4):144–9.

Monnier L, Mas E, Ginet C, Michel F, Villon L, Cristol JP, Colette C. Activation of oxidative stress by acute glucose fluctuations compared with sustained chronic hyperglycemia in patients with type 2 diabetes. *JAMA.* 2006 Apr 12;295(14):1681–7.

Mozaffarian D. Trans fatty acids – effects on systemic inflammation and endothelial function. *Atheroscler Suppl.* 2006 May;7(2):29–32. Epub 2006 May 18.

Mozaffarian D, Aro A, Willett WC. Health effects of trans-fatty acids: experimental and observational evidence. *Eur J Clin Nutr.* 2009 May;63 Suppl 2:S5–21.

Mozaffarian D, Wu JH. Omega-3 fatty acids and cardiovascular disease: effects on risk factors, molecular pathways, and clinical events. *J Am Coll Cardiol.* 2011 Nov 8;58(20):2047–67.

Mozaffarian D, Hao T, Rimm EB, Willett WC, Hu FB. Changes in diet and lifestyle and long-term weight gain in women and men. *N Engl J Med.* 2011 Jun 23;364(25):2392–404

Murphy RR, Renfroe MH, Brevard PB, Lee RE, Gloeckner JW. Cooking does not decrease hydrophilic antioxidant capacity of wild blueberries. *Int J Food Sci Nutr.* 2009;60 Suppl 2:88–98.

Muñoz A, Costa M. Nutritionally mediated oxidative stress and inflammation. *Oxid Med Cell Longev.* 2013;2013:610950.

Mühlbauer RC, Lozano A, Reinli A. Onion and a mixture of vegetables, salads, and herbs affect bone resorption in the rat by a mechanism independent of their base excess. *J Bone Miner Res.* 2002 Jul;17(7):1230–6

Mäkinen S. Skeptikko 4/2000 s 14–17. Nykypaino.

Nelson M, Poulter J. Impact of tea drinking on iron status in the UK: a review. *J Hum Nutr Diet.* 2004 Feb;17(1):43–54.

Neves Ribeiro D, Gonçalves Alfenas Rde C, Bressan J, Brunoro Costa NM. The effect of oilseed consumption on appetite and on the risk of developing type 2 diabetes mellitus. *Nutr Hosp.* 2013 Mar-Apr;28(2):296–305.

Neville CE, Young IS, Gilchrist SE, McKinley MC, Gibson A, Edgar JD, Woodside JV. Effect of increased fruit and vegetable consumption on bone turnover in older adults: a randomised controlled trial. *Osteoporos Int.* 2014 Jan;25(1):223–33.

Noto H, Goto A, Tsujimoto T, Noda M. Low-carbohydrate diets and all-cause mortality: a systematic review and meta-analysis of observational studies. *PLoS One.* 2013;8(1):e55030.

O'Keefe JH, Gheewala NM, O'Keefe JO. Dietary strategies for improving post-prandial glucose, lipids, inflammation, and cardiovascular health. *J Am Coll Cardiol.* 2008 Jan 22;51(3):249–55.

Onakpoya I, Spencer E, Heneghan C, Thompson M. The effect of green tea on blood pressure and lipid profile: A systematic review and meta-analysis of randomized clinical trials. *Nutr Metab Cardiovasc Dis.* 2014 Jan 31.

O'Neil CE, Keast DR, Nicklas TA, Fulgoni VL 3rd. Nut consumption is associated with decreased health risk factors for cardiovascular disease and metabolic syndrome in U.S. adults: NHANES 1999–2004. *J Am Coll Nutr.* 2011 Dec;30(6):502–10.

Ortega JF, Fernández-Elías VE, Hamouti N, Mora-Rodriguez R. Increased blood cholesterol after a high saturated fat diet is prevented by aerobic exercise training. *Appl Physiol Nutr Metab.* 2013 Jan;38(1):42–8.

Palomäki A, Pohjantähti-Maaroos H, Wallenius M, Kankkunen P, Aro H, Husgafvel S, Pihlava JM, Oksanen K. Effects of dietary cold-pressed turnip rapeseed oil and butter on serum lipids, oxidized LDL and arterial elasticity in men with metabolic syndrome. *Lipids Health Dis.* 2010 Dec 1;9:137.

Pan A, Sun Q, Bernstein AM, Schulze MB, Manson JE, Willett WC, Hu FB. Red meat consumption and risk of type 2 diabetes: 3 cohorts of US adults and an updated meta-analysis. *Am J Clin Nutr.* 2011 Oct;94(4):1088–96.

Parlesak A, Eckoldt J, Winkler K, Bode CJ, Schäfer C. Intercorrelations of lipoprotein subfractions and their covariation with lifestyle factors in healthy men. *J Clin Biochem Nutr.* 2014 May;54(3):174–80.

Pate RR, Miller BJ, Davis JM, Slentz CA, Klingshirn LA. Iron status of female runners. *Int J Sport Nutr.* 1993 Jun;3(2):222–31.

Patel BP, Bellissimo N, Luhovyy B, Bennett LJ, Hurton E, Painter JE, Anderson GH. An after-school snack of raisins lowers cumulative food intake in young children. *J Food Sci.* 2013 Jun;78 Suppl 1:A5-A10. (a)

Patel AM, Adeseun GA, Goldfarb S. Calcium-alkali syndrome in the modern era. *Nutrients.* 2013 Nov 27;5(12):4880–93. (b)

Pearce KL, Clifton PM, Noakes M. Egg consumption as part of an energy-restricted high-protein diet improves blood lipid and blood glucose profiles in individuals with type 2 diabetes. *Br J Nutr.* 2011 Feb;105(4):584–92.

Pedersen A, Baumstark MW, Marckmann P, Gylling H, Sandström B. An olive oil-rich diet results in higher concentrations of LDL cholesterol and a higher number of LDL subfraction particles than rapeseed oil and sunflower oil diets. *J Lipid Res.* 2000 Dec;41(12):1901–11.

Pedersen AN, Kondrup J, Børshøj E. Health effects of protein intake in healthy adults: a systematic literature review. *Food Nutr Res.* 2013 Jul 30;57.

Pérez-Jiménez F, Ruano J, Perez-Martinez P, Lopez-Segura F, Lopez-Miranda J. The influence of olive oil on human health: not a question of fat alone. *Mol Nutr Food Res.* 2007 Oct;51(10):1199–208. Review.

Patel PS, Forouhi NG, Kuijsten A, Schulze MB, van Woudenberg GJ, Ardanaz E, Amiano P, Arriola L ym.; InterAct Consortium. The prospective association between total and type of fish intake and type 2 diabetes in 8 European countries: EPIC-InterAct Study. *Am J Clin Nutr.* 2012 Jun;95(6):1445–53.

Peternelj TT, Coombes JS. Antioxidant supplementation during exercise training: beneficial or detrimental? *Sports Med.* 2011 Dec 1;41(12):1043–69.

Petta S, Marchesini G, Caracausi L, Macaluso FS, Cammà C, Ciminnisi S, Cabibi D, Porcasì R, Craxì A ym. Industrial, not fruit fructose intake is associated with the severity of liver fibrosis in genotype 1 chronic hepatitis C patients. *J Hepatol.* 2013 Dec;59(6):1169–76

Poulsen SK, Crone C, Astrup A, Larsen TM. Long-term adherence to the New Nordic Diet and the effects on body weight, anthropometry and blood pressure: a 12-month follow-up study. *Eur J Nutr.* 2014 Mar 25. (a)

Poulsen SK, Due A, Jordy AB, Kiens B, Stark KD, Stender S, Holst C, Astrup A, Larsen TM. Health effect of the New Nordic Diet in adults with increased waist circumference: a 6-mo randomized controlled trial. *Am J Clin Nutr.* 2014 Jan;99(1):35–45. (b)

Power O, Jakeman P, FitzGerald RJ. Antioxidative peptides: enzymatic production, in vitro and in vivo antioxidant activity and potential applications of milk-derived antioxidative peptides. *Amino Acids.* 2013 Mar;44(3):797–820.

Prospective Studies Collaboration, Lewington S, Whitlock G, Clarke R, Sherliker P, Emberson J, Halsey J, ym. Blood cholesterol and vascular mortality by age, sex, and blood pressure: a meta-analysis of individual data from 61 prospective studies with 55,000 vascular deaths. *Lancet.* 2007 Dec 1;370(9602):1829–39. Review.

Qi H, Li S. Dose-response meta-analysis on coffee, tea and caffeine consumption with risk of Parkinson's disease. *Geriatr Gerontol Int.* 2014 Apr;14(2):430–9.

Ratliff J, Leite JO, de Ogburn R, Puglisi MJ, VanHeest J, Fernandez ML. Consuming eggs for breakfast influences plasma glucose and ghrelin, while reducing energy intake during the next 24 hours in adult men. *Nutr Res.* 2010 Feb;30(2):96–103.

Rasmussen LB, Andersen LF, Borodulin K, Enghardt Barbieri H, Fagt S, Matthiessen J, Sveinsson T, Thorgeirsdottir H and Trolle E. Nordic monitoring of diet, physical activity and overweight. First collection of data in all Nordic Countries 2011. *TemaNord* 2012:552

Rebello CJ, Greenway FL, Finley JW. A review of the nutritional value of legumes and their effects on obesity and its related co-morbidities. *Obes Rev.* 2014 May;15(5):392–407.

Reiss R, Johnston J, Tucker K, DeSesso JM, Keen CL. Estimation of cancer risks and benefits associated with a potential increased consumption of fruits and vegetables. *Food Chem Toxicol.* 2012 Dec;50(12):4421–7.

Rodriguez-Leyva D, Dupasquier CM, McCullough R, Pierce GN. The cardiovascular effects of flaxseed and its omega-3 fatty acid, alpha-linolenic acid. *Can J Cardiol.* 2010 Nov;26(9):489–96. Review.

Roehrs T, Roth T. Caffeine: sleep and daytime sleepiness. *Sleep Med Rev.* 2008 Apr;12(2):153–62.

Rohrmann S, Overvad K, Bueno-de-Mesquita HB, Jakobsen MU, Egeberg R, Tjønneland A, Nailler L ym. Meat consumption and mortality--results from the European Prospective Investigation into Cancer and Nutrition. *BMC Med.* 2013 Mar 7;11:63.

Rolls BJ, Ello-Martin JA, Tohill BC. What can intervention studies tell us about the relationship between fruit and vegetable consumption and weight management? *Nutr Rev.* 2004 Jan;62(1):1–17

Rong Y, Chen L, Zhu T, Song Y, Yu M, Shan Z, Sands A, Hu FB, Liu L. Egg consumption and risk of coronary heart disease and stroke: dose-response meta-analysis of prospective cohort studies. *BMJ.* 2013 Jan 7;346:e8539.

Ronksley PE, Brien SE, Turner BJ, Mukamal KJ, Ghali WA. Association of alcohol consumption with selected cardiovascular disease outcomes: a systematic review and meta-analysis. *BMJ.* 2011;342:d671.

Rylander C, Sandanger TM, Engeset D, Lund E. Consumption of lean fish reduces the risk of type 2 diabetes mellitus: a prospective population based cohort study of Norwegian women. *PLoS One.* 2014 Feb 24;9(2):e89845

Sacks FM, Katan M. Randomized clinical trials on the effects of dietary fat and carbohydrate on plasma lipoproteins and cardiovascular disease. *Am J Med.* 2002 Dec 30;113 Suppl 9B:13S-24S.

Salas-Salvadó J, Bulló M, Babio N, Martínez-González MÁ, Ros E, ym. ; PREDIMED Study Investigators. Reduction in the incidence of type 2 diabetes with the Mediterranean diet: results of the PREDIMED-Reus nutrition intervention randomized trial. *Diabetes Care.* 2011 Jan;34(1):14–9.

Santos VC, Levada-Pires AC, Alves SR, Pithon-Curi TC, Curi R, Cury-Boaventura MF. Effects of DHA-rich fish oil supplementation on lymphocyte function before and after a marathon race. *Int J Sport Nutr Exerc Metab.* 2013 Apr;23(2):161–9.

Sarkkinen E, Korhonen M, Erkkilä A, Ebeling T, Uusitupa M. Effect of apolipoprotein E polymorphism on serum lipid response to the separate modification of dietary fat and dietary cholesterol. *Am J Clin Nutr.* 1998 Dec;68(6):1215–22.

Savage JS, Marini M, Birch LL. Dietary energy density predicts women's weight change over 6 y. *Am J Clin Nutr.* 2008 Sep;88(3):677–84.

Schlemmer U, Frølich W, Prieto RM, Grases F. Phytate in foods and significance for humans: food sources, intake, processing, bioavailability, protective role and analysis. *Mol Nutr Food Res.* 2009 Sep;53 Suppl 2:S330–75.

Schwingshackl L, Hoffmann G. Monounsaturated fatty acids and risk of cardiovascular disease: synopsis of the evidence available from systematic reviews and meta-analyses. *Nutrients.* 2012 Dec 11;4(12):1989–2007.

Seagle HM, Strain GW, Makris A, Reeves RS; American Dietetic Association. Position of the American Dietetic Association: weight management. *J Am Diet Assoc* 2009; 109: 330–46.

Shimamoto T, Yamamichi N, Kodashima S, Takahashi Y, Fujishiro M, Oka M, Mitsushima T, Koike K. No association of coffee consumption with gastric ulcer, duodenal ulcer, reflux esophagitis, and non-erosive reflux disease: a cross-sectional study of 8,013 healthy subjects in Japan. *PLoS One.* 2013 Jun 12;8(6):e65996.

Shin JY, Xun P, Nakamura Y, He K. Egg consumption in relation to risk of cardiovascular disease and diabetes: a systematic review and meta-analysis. *Am J Clin Nutr.* 2013 Jul;98(1):146–59.

Si R, Qu K, Jiang Z, Yang X, Gao P. Egg consumption and breast cancer risk: a meta-analysis. *Breast Cancer.* 2014 May;21(3):251–61.

Sievenpiper JL, de Souza RJ, Mirrahimi A, Yu ME, Carleton AJ, Beyene J, Chiavaroli L, Di Buono M **ym** Effect of fructose on body weight in controlled feeding trials: a systematic review and meta-analysis. *Ann Intern Med.* 2012 Feb 21;156(4):291–304. (a)

Silva AM, Júdice PB, Matias CN, Santos DA, Magalhães JP, St-Onge MP, Gonçalves EM **ym**. Total body water and its compartments are not affected by ingesting a moderate dose of caffeine in healthy young adult males. *Appl Physiol Nutr Metab.* 2013 Jun;38(6):626–32.

Silver HJ, Kang H, Keil CD, Muldowney JA 3rd, Kocalis H, Fazio S, Vaughan DE, Niswender KD. Consuming a balanced high fat diet for 16 weeks improves body composition, inflammation and vascular function parameters in obese premenopausal women. *Metabolism.* 2014 Apr;63(4):562–73.

Simopoulos AP. Omega-3 fatty acids and athletics. *Curr Sports Med Rep.* 2007 Jul;6(4):230–6.

Siri-Tarino PW. Effects of diet on high-density lipoprotein cholesterol. *Curr Atheroscler Rep.* 2011 Dec;13(6):453–60

Skulas-Ray AC, Kris-Etherton PM, Harris WS, Vanden Heuvel JP, Wagner PR, West SG: Dose-response effects of omega-3 fatty acids on triglycerides, inflammation, and endothelial function in healthy persons with moderate hypertriglyceridemia. *Am J Clin Nutr.* 2011;93:243–252.

Sluijs I, Forouhi NG, Beulens JW, van der Schouw YT, Agnoli C, Arriola L, Balkau B, Barricarte A, Boeing H **ym**; **InterAct Consortium.** The amount and type of dairy product intake and incident type 2 diabetes: results from the EPIC-InterAct Study. *Am J Clin Nutr.* 2012 Aug;96(2):382–90.

Soczynska JK, Kennedy SH, Woldeyohannes HO, Liauw SS, Alsuwaidan M, Yim CY, McIntyre RS. Mood disorders and obesity: understanding inflammation as a pathophysiological nexus. *Neuromolecular Med.* 2011 Jun;13(2):93–116.

Soedamah-Muthu SS, Ding EL, Al-Delaimy WK, Hu FB, Engberink MF, Willett WC, Geleijnse JM. Milk and dairy consumption and incidence of cardiovascular diseases and all-cause mortality: dose-response meta-analysis of prospective cohort studies. *Am J Clin Nutr.* 2011 Jan;93(1):158–71.

Soerjomataram I, Oomen D, Lemmens V, Oenema A, Benetou V, Trichopoulou A, Coebergh JW **ym**. Increased consumption of fruit and vegetables and future cancer incidence in selected European countries. *Eur J Cancer.* 2010 Sep;46(14):2563–80.

Song Y, Stampfer MJ, Liu S: Meta-analysis: apolipoprotein E genotypes and risk for coronary heart disease. *Ann Intern Med.* 2004;141:137–147.

Song J, Su H, Wang BL, Zhou YY, Guo LL. Fish consumption and lung cancer risk: systematic review and meta-analysis. *Nutr Cancer.* 2014;66(4):539–49.

Soop M, Nehra V, Henderson GC, Boirie Y, Ford GC, Nair KS. Coingestion of whey protein and casein in a mixed meal: demonstration of a more sustained anabolic effect of casein. *Am J Physiol Endocrinol Metab.* 2012 Jul 1;303(1):E152–62.

Sousa M, Teixeira VH, Soares J. Dietary strategies to recover from exercise-induced muscle damage. *Int J Food Sci Nutr.* 2014 Mar;65(2):151–63.

Sontag-Strohm T. Kvinoa jää kauralle kakkoseksi terveellisyydessä. HS 7.11.2013. <http://www.hs.fi/terveys/a1383718884269>

Spector S, Tan R. Exercise-induced bronchoconstriction update: therapeutic management. *Allergy Asthma Proc.* 2012 Jan-Feb;33(1):7–12.

Spence JD, Jenkins DJ, Davignon J. Dietary cholesterol and egg yolks: not for patients at risk of vascular disease. *Can J Cardiol.* 2010 Nov;26(9):e336–9.

Stacewicz-Sapuntzakis M. Dried plums and their products: composition and health effects--an updated review. *Crit Rev Food Sci Nutr.* 2013;53(12):1277–302

Stein O, Thiery J, Stein Y. Is there a genetic basis for resistance to atherosclerosis? *Atherosclerosis.* 2002 Jan;160(1):1–10. Review.

Stratton SL, Henrich CL, Matthews NI, Bogusiewicz A, Dawson AM, Horvath TD, Owen SN, Boysen G, ym. Marginal biotin deficiency can be induced experimentally in humans using a cost-effective outpatient design. *J Nutr.* 2012 Jan;142(1):22–6.

Schwingshackl L, Strasser B, Hoffmann G. Effects of monounsaturated fatty acids on cardiovascular risk factors: a systematic review and meta-analysis. *Ann Nutr Metab.* 2011;59(2–4):176–86.

USDA. <http://ndb.nal.usda.gov/>

Takahashi M, Suzuki K, Kim HK, Otsuka Y, Imaizumi A, Miyashita M, Sakamoto S. Effects of curcumin supplementation on exercise-induced oxidative stress in humans. *Int J Sports Med.* 2014 Jun;35(6):469–75

Tang Y, Jiang H, Bryan NS. Nitrite and nitrate: cardiovascular risk-benefit and metabolic effect. *Curr Opin Lipidol.* 2011 Feb;22(1):11–5.

Temme EH, Van Hoydonck PG. Tea consumption and iron status. *Eur J Clin Nutr.* 2002 May;56(5):379–86.

Teucher B, Olivares M, Cori H. Enhancers of iron absorption: ascorbic acid and other organic acids. *Int J Vitam Nutr Res.* 2004 Nov;74(6):403–19.

Toledo E, Delgado-Rodríguez M, Estruch R, Salas-Salvadó J, Corella D, Gomez-Gracia E, Fiol M, ym. Low-fat dairy products and blood pressure: follow-up of 2290 older persons at high cardiovascular risk participating in the PREDIMED study. *Br J Nutr.* 2009 Jan;101(1):59–67.

Trappe TA, Liu SZ. Effects of prostaglandins and COX-inhibiting drugs on skeletal muscle adaptations to exercise. *J Appl Physiol (1985).* 2013 Sep;115(6):909–19.

Tresserra-Rimbau A, Rimm E, Medina-Remón A, Martínez-González MA, Carmen Lopez-Sabater M, ym. Polyphenol intake and mortality risk: a re-analysis of the PREDIMED trial. *BMC Med.* 2014; 12:77. (a)

Tresserra-Rimbau A, Rimm E, Medina-Remón A, Martínez-González MA, de la Torre R, Corella D, ym. Inverse association between habitual polyphenol intake and incidence of cardiovascular events in the PREDIMED study. *Nutr Metab Cardiovasc Dis.* 2014 (b)

Trichopoulou A, Bamia C, Trichopoulos D. Anatomy of health effects of Mediterranean diet: Greek EPIC prospective cohort study. *BMJ.* 2009 Jun 23;338:b2337.

Tse G, Eslick GD. Egg consumption and risk of GI neoplasms: dose-response meta-analysis and systematic review. *Eur J Nutr.* 2014 Feb 6

Urpi-Sarda M, Casas R, Chiva-Blanch G, Romero-Mamani ES, Valderas-Martínez P, Arranz S, ym. Virgin olive oil and nuts as key foods of the Mediterranean diet effects on inflammatory biomarkers related to atherosclerosis. *Pharmacol Res.* 2012 Jun;65(6):577–83.

Uusitupa M, Sarkkinen E. Kananmunat ja kolesteroli – onko pääkirjoituksen sanoma tulkittu oikein? *Lääketieteellinen Aikakauskirja Duodecim* 2000;116(6):628–630

Vadivel V, Kunyanga CN, Biesalski HK. Health benefits of nut consumption with special reference to body weight control. *Nutrition.* 2012 Nov-Dec;28(11–12):1089–97

- Vainio H, Weiderpass E.** Fruit and vegetables in cancer prevention. *Nutr Cancer*. 2006;54(1):111–42.
- Vander Wal JS, Gupta A, Khosla P, Dhurandhar NV.** Egg breakfast enhances weight loss. *Int J Obes (Lond)*. 2008 Oct;32(10):1545–51.
- van het Hof KH, de Boer BC, Tijburg LB, Lucius BR, Zijp I, West CE, Hautvast JG, Weststrate JA.** Carotenoid bioavailability in humans from tomatoes processed in different ways determined from the carotenoid response in the triglyceride-rich lipoprotein fraction of plasma after a single consumption and in plasma after four days of consumption. *J Nutr*. 2000 May;130(5):1189–96.
- Van Wijck K, Lenaerts K, Van Bijnen AA, Boonen B, Van Loon LJ, Dejong CH, Buurman WA.** Aggravation of exercise-induced intestinal injury by Ibuprofen in athletes. *Med Sci Sports Exerc*. 2012 Dec;44(12):2257–62
- Vartiainen E ym.** Terveyden ja hyvinvoinnin laitos, syöpätautien asiantuntijaryhmä. Syövän ehkäisyn, varhaisen toteamisen ja kuntoutumisen tuen kehittäminen vuosina 2014–2025. Kansallisen syöpäsuounnitelman II osa. Juvenes print, Suomen Yliopistopaino Oy. Tampere 2014.
- Vázquez C, Botella-Carretero JI, Corella D, Fiol M, Lage M, ym. WISH-CARE Study Investigators.** White fish reduces cardiovascular risk factors in patients with metabolic syndrome: the WISH-CARE study, a multicenter randomized clinical trial. *Nutr Metab Cardiovasc Dis*. 2014 Mar;24(3):328–35.
- Venkatraman JT, Leddy J, Pendergast D.** Dietary fats and immune status in athletes: clinical implications. *Med Sci Sports Exerc*. 2000 Jul;32(7 Suppl):S389–95. Review.
- Venkatraman JT, Pendergast DR.** Effect of dietary intake on immune function in athletes. *Sports Med*. 2002;32(5):323–37. Review.
- Vergnaud A-C, Norat T, Romaguera D, Mouw T, May AM, Travier N, et al.** Meat consumption and prospective weight change in participants of the EPIC-PANACEA study. *Am J Clin Nutr*. 2010;92:398–407.
- Vislocky LM, Pikosky MA, Rubin KH, Vega-López S, Gaine PC, Martin WF, Zern TL, Lofgren IE ym.** Habitual consumption of eggs does not alter the beneficial effects of endurance training on plasma lipids and lipoprotein metabolism in untrained men and women. *J Nutr Biochem*. 2009 Jan;20(1):26–34.
- Volek JS, Kraemer WJ, Bush JA, Incledon T, Boetes M.** Testosterone and cortisol in relationship to dietary nutrients and resistance exercise. *J Appl Physiol* (1985). 1997 Jan;82(1):49–54.
- VRN.** Terveytä ruuasta – Suomalaiset ravitsemussuoitus 2014. Juvenes Oy, Helsinki 2014.
- Vucenik I, Shamsuddin AM.** Protection against cancer by dietary IP6 and inositol. *Nutrition and Cancer* 2006;55:109–125.
- Wallin A, Di Giuseppe D, Orsini N, Patel PS, Forouhi NG, Wolk A.** Fish consumption, dietary long-chain n-3 fatty acids, and risk of type 2 diabetes: systematic review and meta-analysis of prospective studies. *Diabetes Care*. 2012 Apr;35(4):918–29.
- Waterman JJ, Kapur R.** Upper gastrointestinal issues in athletes. *Curr Sports Med Rep*. 2012 Mar-Apr;11(2):99–104.
- Wei MY, Jacobson TA.** Effects of eicosapentaenoic acid versus docosahexaenoic acid on serum lipids: a systematic review and meta-analysis. *Curr Atheroscler Rep*. 2011 Dec;13(6):474–83.
- Weitzberg E, Lundberg JO.** Novel aspects of dietary nitrate and human health. *Annu Rev Nutr*. 2013;33:129–59
- Welch AA, Mulligan A, Bingham SA, Khaw KT.** Urine pH is an indicator of dietary acid-base load, fruit and vegetables and meat intakes: results from the European Prospective Investigation into Cancer and Nutrition (EPIC)-Norfolk population study. *Br J Nutr*. 2008 Jun;99(6):1335–43.
- Wien M, Haddad E, Oda K, Sabaté J.** A randomized 3x3 crossover study to evaluate the effect of Hass avocado intake on post-ingestive satiety, glucose and insulin levels, and subsequent energy intake in overweight adults. *Nutr J*. 2013 Nov 27;12:155. doi: 10.1186/1475–2891–12–155.
- Woodside JV, Young IS, McKinley MC.** Fruit and vegetable intake and risk of cardiovascular disease. *Proc Nutr Soc*. 2013 Nov;72(4):399–406
- World Health Organisation.** Geneva: Evaluation of Certain Food Additives; 2002. Fifty ninth Report of the Joint FAO/WHO Expert Committee on Food Additives (JECFA)

World Cancer Research Fund / American Institute for Cancer Research. Washington DC: AICR; 2007. Food, Nutrition, Physical Activity, and the Prevention of Cancer: A Global Perspective.

Wu JH, Micha R, Imamura F, Pan A, Biggs ML, Ajaz O, Djousse L, Hu FB, Mozaffarian D. Omega-3 fatty acids and incident type 2 diabetes: a systematic review and meta-analysis. *Br J Nutr.* 2012 Jun;107 Suppl 2:S214–27.

Wu L, Ma D, Walton-Moss B, He Z. Effects of low-fat diet on serum lipids in premenopausal and postmenopausal women: a meta-analysis of randomized controlled trials. *Menopause.* 2014 Jan;21(1):89–99.

Xu F, Zheng Y, Yang Z, Cao S, Shao X, Wang H. Domestic cooking methods affect the nutritional quality of red cabbage. *Food Chem.* 2014 Oct 15;161:162–7.

Xun P, He K. Fish Consumption and Incidence of Diabetes: meta-analysis of data from 438,000 individuals in 12 independent prospective cohorts with an average 11-year follow-up. *Diabetes Care.* 2012 Apr;35(4):930–8.

Yang WS, Wang WY, Fan WY, Deng Q, Wang X. Tea consumption and risk of type 2 diabetes: a dose-response meta-analysis of cohort studies. *Br J Nutr.* 2014 Apr 28;111(8):1329–39.

Ye EQ, Chacko SA, Chou EL, Kugizaki M, Liu S. Greater whole-grain intake is associated with lower risk of type 2 diabetes, cardiovascular disease, and weight gain. *J Nutr.* 2012 Jul;142(7):1304–13.

Yi M, Fu J, Zhou L, Gao H, Fan C, Shao J, Xu B, Wang Q, Li J, Huang G, Lapsley K, Blumberg JB, Chen CY. The effect of almond consumption on elements of endurance exercise performance in trained athletes. *J Int Soc Sports Nutr.* 2014 May 11;11:18.

Young JF, Therkildsen M, Ekstrand B, Che BN, Larsen MK, Oksbjerg N, Stagsted J. Novel aspects of health promoting compounds in meat. *Meat Sci.* 2013 Dec;95(4):904–11.

Yu X, Bao Z, Zou J, Dong J. Coffee consumption and risk of cancers: a meta-analysis of cohort studies. *BMC Cancer.* 2011 Mar 15;11:96.

Zahradka P, Wright B, Weighell W, Blewett H, Baldwin A, O K, Guzman RP, Taylor CG. Daily non-soy legume consumption reverses vascular impairment due to peripheral artery disease. *Atherosclerosis.* 2013 Oct;230(2):310–4.

Załęcka A, Bügel S, Paoletti F, Kahl J, Bonanno A, Dostalova A, Rahmann G. The influence of organic production on food quality – research findings, gaps and future challenges. *J Sci Food Agric.* 2014 Jan 16.

Zemel MB. Proposed role of calcium and dairy food components in weight management and metabolic health. *Phys Sportsmed.* 2009;37:29–39.

Zhao Z, Li S, Liu G, Yan F, Ma X, Huang Z, Tian H. Body iron stores and heme-iron intake in relation to risk of type 2 diabetes: a systematic review and meta-analysis. *PLoS One.* 2012;7(7):e41641.

Zheng W, Lee SA. Well-done meat intake, heterocyclic amine exposure, and cancer risk. *Nutr Cancer.* 2009;61(4):437–46.

Zheng XX, Xu YL, Li SH, Liu XX, Hui R, Huang XH. Green tea intake lowers fasting serum total and LDL cholesterol in adults: a meta-analysis of 14 randomized controlled trials. *Am J Clin Nutr.* 2011 Aug;94(2):601–10.

Zheng JS, Huang T, Yang J, Fu YQ, Li D. Marine N-3 polyunsaturated fatty acids are inversely associated with risk of type 2 diabetes in Asians: a systematic review and meta-analysis. *PLoS One.* 2012;7(9):e44525.

Zhou Y, Tian C, Jia C. Association of fish and n-3 fatty acid intake with the risk of type 2 diabetes: a meta-analysis of prospective studies. *Br J Nutr.* 2012 Aug;108(3):408–17.

Zhou D, Yu H, He F, Reilly KH, Zhang J, Li S, Zhang T, Wang B, Ding Y, Xi B. Nut consumption in relation to cardiovascular risk and type 2 diabetes: a systematic review and meta-analysis of prospective studies. *Am J Clin Nutr.* 2014 May 7

Zijp IM, Korver O, Tijburg LB. Effect of tea and other dietary factors on iron absorption. *Crit Rev Food Sci Nutr.* 2000 Sep;40(5):371–98. Review.

Ziltener JL, Leal S, Fournier PE. Non-steroidal anti-inflammatory drugs for athletes: an update. *Ann Phys Rehabil Med.* 2010 May;53(4):278–82, 282–8.

Österdahl M, Kocturk T, Koochek A, Wändell PE. Effects of a short-term intervention with a paleolithic diet in healthy volunteers. *Eur J Clin Nutr.* 2008 May;62(5):682–5.

Lähdeluettelo luku 4

- Arciero PJ, Ormsbee MJ, Gentile CL, Nindl BC, Brestoff JR, Ruby M.** Increased protein intake and meal frequency reduces abdominal fat during energy balance and energy deficit. *Obesity (Silver Spring)*. 2013 Jul;21(7):1357–66
- Barnosky AR, Hoddy KK, Unterman TG, Varady KA.** Intermittent fasting vs daily caloric restriction for type 2 diabetes prevention: a review of human findings. *Transl Res*. 2014 Jun 12. pii: S1931-5244(14)00200-X.
- Baron KG, Reid KJ, Horn LV, Zee PC.** Contribution of evening macronutrient intake to total caloric intake and body mass index. *Appetite*. 2013 Jan;60(1):246–51.
- Bellisle F, McDevitt R, Prentice AM.** Meal frequency and energy balance. *Br J Nutr*. 1997;77(Suppl 1):S57–70.
- Bhutani S, Klempel MC, Kroeger CM, Trepanowski JF, Varady KA.** Alternate day fasting and endurance exercise combine to reduce body weight and favorably alter plasma lipids in obese humans. *Obesity (Silver Spring)*. 2013 Jul;21(7):1370–9.
- Brown D, Wyon M.** The effect of moderate glycemic energy bar consumption on blood glucose and mood in dancers. *Med Probl Perform Art*. 2014 Mar;29(1):27–31.
- Burke LM, Kiens B, Ivy JL.** Carbohydrates and fat for training and recovery. *J Sports Sci*. 2004 Jan;22(1):15–30.
- Carlson O, Martin B, Stote KS, Golden E, Maudsley S, Najjar SS, Ferrucci L, Ingram DK, Longo DL, Rumpler WV, Baer DJ, Egan Jym.** Impact of reduced meal frequency without caloric restriction on glucose regulation in healthy, normal-weight middle-aged men and women. *Metabolism*. 2007 Dec;56(12):1729–34.
- de Castro JM.** When, how much and what foods are eaten are related to total daily food intake. *Br J Nutr*. 2009 Oct;102(8):1228–37.
- Deutz RC, Benardot D, Martin DE, Cody MM.** Relationship between energy deficits and body composition in elite female gymnasts and runners. *Med Sci Sports Exerc*. 2000 Mar;32(3):659–68.
- Edelstein SL, Barrett-Connor EL, Wingard DL, Cohn BA.** Increased meal frequency associated with decreased cholesterol concentrations; Rancho Bernardo, CA, 1984–1987. *Am J Clin Nutr*. 1992;55(3):664–9
- Erdman KA, Tunnicliffe J, Lun VM, Reimer RA.** Eating patterns and composition of meals and snacks in elite Canadian athletes. *Int J Sport Nutr Exerc Metab*. 2013 Jun;23(3):210–9.
- Fricker J, Giroux S, Fumeron F, Apfelbaum M.** Circadian rhythm of energy intake and corpulence status in adults. *Int J Obes*. 1990;14(5):387–93
- Giovannini M, Agostoni C, Shamir R.** Symposium overview: Do we all eat breakfast and is it important? *Crit Rev Food Sci Nutr*. 2010 Feb;50(2):97–9.
- Griffin BA.** Nonpharmacological approaches for reducing serum low-density lipoprotein cholesterol. *Curr Opin Cardiol*. 2014 Jul;29(4):360–5.
- Halson SL.** Sleep in elite athletes and nutritional interventions to enhance sleep. *Sports Med*. 2014 May;44 Suppl 1:S13–23
- Hatami Zargaran Z, Salehi M, Heydari ST, Babajafari S.** The Effects of 6 Isocaloric Meals on Body Weight, Lipid Profiles, Leptin, and Adiponectin in Overweight Subjects (BMI > 25). *Int Cardiovasc Res J*. 2014 Apr;8(2):52–6.
- Holmback I, Ericson U, Gullberg B, Wirkkula E.** A high eating frequency is associated with an overall healthy lifestyle in middle-aged men and women and reduced likelihood of general and central obesity in men. *Br J Nutr*. 2010;104(7):1065–73.
- Heilbronn LK, Smith SR, Martin CK, Anton SD, Ravussin E.** Alternate-day fasting in nonobese subjects: effects on body weight, body composition, and energy metabolism. *Am J Clin Nutr*. 2005;12:69–73.
- Horikawa C, Kodama S, Yachi Y, Heianza Y, Hirasawa R, Ibe Y, Saito K, Shimano H, Yamada N, Sone H.** Skipping breakfast and prevalence of overweight and obesity in Asian and Pacific regions: a meta-analysis. *Prev Med*. 2011 Oct;53(4–5):260–7.
- Hoyland A, Dye L, Lawton CL.** A systematic review of the effect of breakfast on the cognitive performance of children and adolescents. *Nutr Res Rev*. 2009 Dec;22(2):220–43.

Iwao S, Mori K, Sato Y. Effects of meal frequency on body composition during weight control in boxers. *Scand J Med Sci Sports*. 1996 Oct;6(5):265–72.

Klempel MC, Kroeger CM, Varady KA. Alternate day fasting (ADF) with a high-fat diet produces similar weight loss and cardio-protection as ADF with a low-fat diet. *Metabolism*. 2013;12:137–143.

Kobayashi F, Ogata H, Omi N, Nagasaka S, Yamaguchi S, Hibi M, Tokuyama K. Effect of breakfast skipping on diurnal variation of energy metabolism and blood glucose. *Obes Res Clin Pract*. 2014 May-Jun;8(3):e201–98.

Koletzko B, Toschke AM. Meal patterns and frequencies: do they affect body weight in children and adolescents? *Crit Rev Food Sci Nutr*. 2010 Feb;50(2):100–5.

Kutsuma A, Nakajima K, Suwa K. Potential Association between Breakfast Skipping and Concomitant Late-Night-Dinner Eating with Metabolic Syndrome and Proteinuria in the Japanese Population. *Scientifica (Cairo)*. 2014;2014:253581.

La Bounty PM, Campbell BI, Wilson J, Galvan E, Berardi J, Kleiner SM, et al. International Society of Sports Nutrition position stand: meal frequency. *J Int Soc Sports Nutr*. 2011;8:4.

LeCheminant JD, Christenson E, Bailey BW, Tucker LA. Restricting night-time eating reduces daily energy intake in healthy young men: a short-term cross-over study. *Br J Nutr*. 2013 Dec 14;110(11):2108–13.

Leidy HJ, Campbell WW. The effect of eating frequency on appetite control and food intake: brief synopsis of controlled feeding studies. *J Nutr*. 2011 Jan;141(1):154–7.

Leidy HJ, Ortinau LC, Douglas SM, Hoertel HA. Beneficial effects of a higher-protein breakfast on the appetitive, hormonal, and neural signals controlling energy intake regulation in overweight/obese, “breakfast-skipping,” late-adolescent girls. *Am J Clin Nutr*. 2013 Apr;97(4):677–88.

Levitsky DA, Pacanowski CR. Effect of skipping breakfast on subsequent energy intake. *Physiol Behav*. 2013 Jul 2;119:9–16.

Liu AG, Most MM, Brashears MM, Johnson WD, Cefalu WT, Greenway FL. Reducing the glycemic index or carbohydrate content of mixed meals reduces postprandial glycemia and insulinemia over the entire day but does not affect satiety. *Diabetes Care*. 2012 Aug;35(8):1633–7.

Lombardo M, Bellia A, Padua E, Annino G, Guglielmi V, D'Adamo M, Iellamo F, Sbraccia P. Morning meal more efficient for fat loss in a 3-month lifestyle intervention. *J Am Coll Nutr*. 2014;33(3):198–205.

Louis-Sylvestre J, Lluch A, Neant F, Blundell JE. Highlighting the positive impact of increasing feeding frequency on metabolism and weight management. *Forum Nutr*. 2003;56:126–8.

Ma Y, Bertone ER, Stanek E, Reed GW, Hebert JR, Cohen NL, et al. Association between eating patterns and obesity in a free-living US adult population. *Am J Epidemiol*. 2003;158(1):85–92.

McCrory MA. Meal skipping and variables related to energy balance in adults: A brief review, with emphasis on the breakfast meal. *Physiol Behav*. 2014 May 11. pii: S0031-9384(14)00260-1.

Morgan LM, Shi JW, Hampton SM, Frost G. Effect of meal timing and glycaemic index on glucose control and insulin secretion in healthy volunteers. *Br J Nutr*. 2012 Oct;108(7):1286–91.

Munsters MJ, Saris WH. Body weight regulation and obesity: dietary strategies to improve the metabolic profile. *Annu Rev Food Sci Technol*. 2014;5:39–51.

Odegaard AO, Jacobs DR Jr, Steffen LM, Van Horn L, Ludwig DS, Pereira MA. Breakfast frequency and development of metabolic risk. *Diabetes Care*. 2013 Oct;36(10):3100–6.

Robinson E, Aveyard P, Daley A, Jolly K, Lewis A, Lycett D, Higgs S. Eating attentively: a systematic review and meta-analysis of the effect of food intake memory and awareness on eating. *Am J Clin Nutr*. 2013 Apr;97(4):728–42.

Robinson E, Almiron-Roig E, Rutters F, de Graaf C, Forde CG, Tudur Smith C, Nolan SJ, Jebb SA. A systematic review and meta-analysis examining the effect of eating rate on energy intake and hunger. *Am J Clin Nutr*. 2014 May 21;100(1):123–151

Ruidavets JB, Bongard V, Bataille V, Gourdy P, Ferrieres J. Eating frequency and body fatness in middle-aged men. *Int J Obes Relat Metab Disord*. 2002;26(11):1476–83.

Seppälä L. Yhdessä syöminen kannattaa. HS 26.6.2014. s. C11.

Smeets AJ, Westerterp-Plantenga MS. Acute effects on metabolism and appetite profile of one meal difference in the lower range of meal frequency. *Br J Nutr.* 2008;99(6):1316–21.

Stote KS, Baer DJ, Spears K, Paul DR, Harris GK, Rumpler WV, Strycula P, Najjar SS, Ferrucci L, Ingram DK, Longo DL, Mattson MP. A controlled trial of reduced meal frequency without caloric restriction in healthy, normal-weight, middle-aged adults. *Am J Clin Nutr.* 2007 Apr;85(4):981–8.

Szajewska H, Ruszczynski M. Systematic review demonstrating that breakfast consumption influences body weight outcomes in children and adolescents in Europe. *Crit Rev Food Sci Nutr.* 2010 Feb;50(2):113–9.

Tanaka M, Mizuno K, Fukuda S, Shigihara Y, Watanabe Y. Relationships between dietary habits and the prevalence of fatigue in medical students. *Nutrition.* 2008 Oct;24(10):985–9.

Rothschild J, Hoddy KK, Jambazian P, Varady KA. Time-restricted feeding and risk of metabolic disease: a review of human and animal studies. *Nutr Rev.* 2014 May;72(5):308–18.

Varady KA, Bhutani S, Church EC, Klempel MC. Short-term modified alternate-day fasting: a novel dietary strategy for weight loss and cardioprotection in obese adults. *Am J Clin Nutr.* 2009;12:1138–1143

Varady KA, Bhutani S, Klempel MC, Kroeger CM, Trepanowski JF, Haus JM, Hoddy KK, Calvo Y. Alternate day fasting for weight loss in normal weight and overweight subjects: a randomized controlled trial. *Nutr J.* 2013 Nov 12;12(1):146.

Westerterp-Plantenga MS, Wijckmans-Duysens NA, ten Hoor F. Food intake in the daily environment after energy-reduced lunch, related to habitual meal frequency. *Appetite.* 1994;22(2):173–82.

Zilberman T, Zilberman EY. Breakfast: to skip or not to skip? *Front Public Health.* 2014 Jun 3;2:59.

Lähdeluettelo luku 5

Achten J, Jentjens RL, Brouns F, Jeukendrup AE. Exogenous oxidation of isomaltulose is lower than that of sucrose during exercise in men. *J Nutr.* 2007 May;137(5):1143–8.

Achten J, Halson SL, Moseley L, Rayson MP, Casey A, Jeukendrup AE. Higher dietary carbohydrate content during intensified running training results in better maintenance of performance and mood state. *J Appl Physiol (1985).* 2004 Apr;96(4):1331–40.

Aguilar F, Riccardo Crebelli, Birgit Dusemund, Pierre Galtier, David Gott, Ursula Gundert-Remy ym. Scientific Opinion on the re-evaluation of aspartame (E 951) as a food additive. *EFSA Journal* 2013;11(12):3496 [263 pp.].

Alhazmi A, Stojanovski E, McEvoy M, Garg ML. The association between dietary patterns and type 2 diabetes: a systematic review and meta-analysis of cohort studies. *J Hum Nutr Diet.* 2014 Jun;27(3):251–60.

Alvarez-Suarez JM, Giampieri F, Battino M. Honey as a source of dietary antioxidants: structures, bioavailability and evidence of protective effects against human chronic diseases. *Curr Med Chem.* 2013;20(5):621–38.

Anderson JW, Baird P, Davis RH Jr, Ferreri S, Knudtson M, Koraym A, Waters V, Williams CL. Health benefits of dietary fiber. *Nutr Rev.* 2009 Apr;67(4):188–205.

Anderson GH, Foreyt J, Sigman-Grant M, Allison DB. The use of low-calorie sweeteners by adults: impact on weight management. *J Nutr.* 2012 Jun;142(6):1163S–9S.

Appel LJ, Sacks FM, Carey VJ, Obarzanek E, Swain JF, Miller ER 3rd, Conlin PR, Erlinger TP, Rosner BA ym. Effects of protein, monounsaturated fat, and carbohydrate intake on blood pressure and serum lipids: results of the OmniHeart randomized trial. *JAMA.* 2005 Nov 16;294(19):2455–64.

Aragon AA, Schoenfeld BJ. Nutrient timing revisited: is there a post-exercise anabolic window? *J Int Soc Sports Nutr.* 2013 Jan 29;10(1):5.

Areta JL, Burke LM, Ross ML, Camera DM, West DW, Broad EM, Jeacocke NA ym. Timing and distribution of protein ingestion during prolonged recovery from resistance exercise alters myofibrillar protein synthesis. *J Physiol.* 2013 May 1;591(Pt 9):2319–31.

Arnall DA, Nelson AG, Quigley J, Lex S, Dehart T, Fortune P. Supercompensated glycogen loads persist 5 days in resting trained cyclists. *Eur J Appl Physiol*. 2007 Feb;99(3):251–6. Epub 2006 Nov 22.

Baar K. Nutrition and the adaptation to endurance training. *Sports Med*. 2014 May;44 Suppl 1:S5–12.

Backhouse SH, Bishop NC, Biddle SJ, Williams C. Effect of carbohydrate and prolonged exercise on affect and perceived exertion. *Med Sci Sports Exerc*. 2005 Oct;37(10):1768–73.

Bao J, Atkinson F, Petocz P, Willett WC, Brand-Miller JC. Prediction of postprandial glycemia and insulinemia in lean, young, healthy adults: glycemic load compared with carbohydrate content alone. *Am J Clin Nutr*. 2011;93(5):984–96.

Barclay AW, Petocz P, McMillan-Price J, Flood VM, Prvan T, Mitchell P, Brand-Miller JC. Glycemic index, glycemic load, and chronic disease risk--a meta-analysis of observational studies. *Am J Clin Nutr*. 2008 Mar;87(3):627–37. Review.

Barenys M, Recasens MA, Martí-Henneberg C, Salas-Salvadó J. Effect of exercise and protein intake on energy expenditure in adolescents. *Rev Esp Fisiol*. 1993 Dec;49(4):209–17.

Bartlett JD, Louhelainen J, Iqbal Z, Cochran AJ, Gibala MJ, Gregson W, Close GL, Drust B, Morton JP. Reduced carbohydrate availability enhances exercise-induced p53 signaling in human skeletal muscle: implications for mitochondrial biogenesis. *Am J Physiol Regul Integr Comp Physiol*. 2013 Mar 15;304(6):R450–8.

Bazzano LA. Effects of soluble dietary fiber on low-density lipoprotein cholesterol and coronary heart disease risk. *Curr Atheroscler Rep*. 2008 Dec;10(6):473–7. Review.

Beaven CM, Maulder P, Pooley A, Kilduff L, Cook C. Effects of caffeine and carbohydrate mouth rinses on repeated sprint performance. *Appl Physiol Nutr Metab*. 2013 Jun;38(6):633–7.

Beelen M, Burke LM, Gibala MJ, van Loon L JC. Nutritional strategies to promote postexercise recovery. *Int J Sport Nutr Exerc Metab*. 2010 Dec;20(6):515–32.

Beelen M, Zorenc A, Pennings B, Senden JM, Kuipers H, van Loon LJ. Impact of protein coingestion on muscle protein synthesis during continuous endurance type exercise. *Am J Physiol Endocrinol Metab*. 2011 Jun;300(6):E945–54.

Beis LY, Willkomm L, Ross R, Bekele Z, Wolde B, Fudge B, Pitsiladis YP. Food and macronutrient intake of elite Ethiopian distance runners. *J Int Soc Sports Nutr*. 2011 May 19;8:7

Bell EA, Castellanos VH, Pelkman CL, Thorwart ML, Rolls BJ. Energy density of foods affects energy intake in normal-weight women. *Am J Clin Nutr*. 1998 Mar;67(3):412–20.

Bennett CB, Chilibeck PD, Barss T, Vatanparast H, Vandenberg A, Zello GA. Metabolism and performance during extended high-intensity intermittent exercise after consumption of low- and high-glycaemic index pre-exercise meals. *Br J Nutr*. 2012 Aug;108 Suppl 1:S81–90

Bentley DJ, Cox GR, Green D, Laursen PB. Maximising performance in triathlon: applied physiological and nutritional aspects of elite and non-elite competitions. *J Sci Med Sport*. 2008 Jul;11(4):407–16.

Berardi JM, Noreen EE, Lemon PW. Recovery from a cycling time trial is enhanced with carbohydrate-protein supplementation vs. isoenergetic carbohydrate supplementation. *J Int Soc Sports Nutr*. 2008 Dec 24;5:24.

Betts JA, Williams C. Short-term recovery from prolonged exercise: exploring the potential for protein ingestion to accentuate the benefits of carbohydrate supplements. *Sports Med*. 2010 Nov 1;40(11):941–59.

Bidwell AJ, Fairchild TJ, Redmond J, Wang L, Keslacy S, Kanaley JA. Physical Activity Offsets the Negative Effects of a High Fructose Diet. *Med Sci Sports Exerc*. 2014 May 20.

Bijkerk CJ, Muris JW, Knottnerus JA, Hoes AW, de Wit NJ. Systematic review: the role of different types of fibre in the treatment of irritable bowel syndrome. *Aliment Pharmacol Ther*. 2004 Feb 1;19(3):245–51.

Bird SP, Tarpenning KM, Marino FE. Independent and combined effects of liquid carbohydrate/essential amino acid ingestion on hormonal and muscular adaptations following resistance training in untrained men. *Eur J Appl Physiol*. 2006 May;97(2):225–38.

Blomstrand E, Saltin B. Effect of muscle glycogen on glucose, lactate and amino acid metabolism during exercise and recovery in human subjects. *J Physiol*. 1999 Jan 1;514 (Pt 1):293–302.

- Blundell JE, Hill AJ.** Paradoxical effects of an intense sweetener (aspartame) on appetite. *Lancet* 1986;1:1092–3
- Bonnema AL, Kolberg LW, Thomas W, Slavin JL.** Gastrointestinal tolerance of chicory inulin products. *J Am Diet Assoc.* 2010 Jun;110(6):865–8
- Bornet FR.** Undigestible sugars in food products. *Am J Clin Nutr.* 1994 Mar;59(3 Suppl):763S-769S.
- Bottoms L, Sinclair J, Taylor K, Polman R, Fewtrell D.** The effects of carbohydrate ingestion on the badminton serve after fatiguing exercise. *J Sports Sci.* 2012;30(3):285–93
- Brand-Miller J, Hayne S, Petocz P, et al. (2003)** Low-glycemic index diets in the management of diabetes: a meta-analysis of randomized controlled trials. *Diabetes Care* 26, 2261–2267.
- Brand-Miller J, Buyken AE.** The glycemic index issue. *Curr Opin Lipidol.* 2012 Feb;23(1):62–7.
- Bray GA.** Energy and fructose from beverages sweetened with sugar or high-fructose corn syrup pose a health risk for some people. *Adv Nutr.* 2013 Mar 1;4(2):220–5.
- Breen L, Philp A, Witard OC, Jackman SR, Selby A, Smith K, Baar K, Tipton KD.** The influence of carbohydrate-protein co-ingestion following endurance exercise on myofibrillar and mitochondrial protein synthesis. *J Physiol.* 2011 Aug 15;589(Pt 16):4011–25.
- Brown LJ, Midgley AW, Vince RV, Madden LA, McNaughton LR.** High versus low glycemic index 3-h recovery diets following glycogen-depleting exercise has no effect on subsequent 5-km cycling time trial performance. *J Sci Med Sport.* 2013 Sep;16(5):450–4
- Brun JF, Fedou C, Mercier J.** Postprandial reactive hypoglycemia. *Diabetes Metab.* 2000 Nov;26(5):337–51. Review.
- Burke LM, Millet G, Tarnopolsky MA; International Association of Athletics Federations.** Nutrition for distance events. *J Sports Sci.* 2007;25 Suppl 1:S29–38.
- Bussau VA, Fairchild TJ, Rao A, Steele P, Fournier PA.** Carbohydrate loading in human muscle: an improved 1 day protocol. *Eur J Appl Physiol.* 2002 Jul;87(3):290–5.
- Carlson LA, Headley S, DeBruin J, Tuckow AT, Koch AJ, Kenefick RW.** Carbohydrate supplementation and immune responses after acute exhaustive resistance exercise. *Int J Sport Nutr Exerc Metab.* 2008 Jun;18(3):247–59.
- Carter J, Jeukendrup AE, Mundel T, Jones DA.** Carbohydrate supplementation improves moderate and high-intensity exercise in the heat. *Pflugers Arch.* 2003 May;446(2):211–9.
- Carter JM, Jeukendrup AE, Mann CH, Jones DA.** The effect of glucose infusion on glucose kinetics during a 1-h time trial. *Med Sci Sports Exerc.* 2004 Sep;36(9):1543–50. (a)
- Carter JM, Jeukendrup AE, Jones DA.** The effect of carbohydrate mouth rinse on 1-h cycle time trial performance. *Med Sci Sports Exerc.* 2004 Dec;36(12):2107–11. (b)
- Cermak NM, van Loon LJ.** The use of carbohydrates during exercise as an ergogenic aid. *Sports Med.* 2013 Nov;43(11):1139–55.
- Chambers ES, Bridge MW, Jones DA.** Carbohydrate sensing in the human mouth: effects on exercise performance and brain activity. *J Physiol.* 2009 Apr 15;587(Pt 8):1779–94.
- Chan TF, Lin WT, Chen YL, Huang HL, Yang WZ, Lee CY, Chen MH, Wang TN, Huang MC, Chiu YW, et al.** Elevated serum triglyceride and retinol-binding protein 4 levels associated with fructose-sweetened beverages in adolescents. *PLoS One.* 2014 Jan 27;9(1):e82004.
- Charalampopoulos D, Rastall RA.** Prebiotics in foods. *Curr Opin Biotechnol.* 2012 Apr;23(2):187–91.
- Chen YJ, Wong SH, Wong CK, Lam CW, Huang YJ, Siu PM.** The effect of a pre-exercise carbohydrate meal on immune responses to an endurance performance run. *Br J Nutr.* 2008 Dec;100(6):1260–8. (a)
- Chen Y, Wong SH, Xu X, Hao X, Wong CK, Lam CW.** Effect of CHO loading patterns on running performance. *Int J Sports Med.* 2008 Jul;29(7):598–606. (b)

Chiu S, Sievenpiper JL, de Souza RJ, Cozma AI, Mirrahimi A, Carleton AJ, Ha V, Di Buono M, Jenkins AL ym.
Effect of fructose on markers of non-alcoholic fatty liver disease (NAFLD): a systematic review and meta-analysis of controlled feeding trials. Eur J Clin Nutr. 2014 Apr;68(4):416–23.

Choi Y, Giovannucci E, Lee JE. Glycaemic index and glycaemic load in relation to risk of diabetes-related cancers: a meta-analysis. Br J Nutr. 2012 Dec 14;108(11):1934–47

Chong E, Guelfi KJ, Fournier PA. Effect of a carbohydrate mouth rinse on maximal sprint performance in competitive male cyclists. J Sci Med Sport. 2011 Mar;14(2):162–7.

Chou SW, Lai CH, Hsu TH, Cho YM, Ho HY, Lai YC, Chen SM, Ho CF, Kuo CH. Characteristics of glycemic control in elite power and endurance athletes. Prev Med. 2005 May;40(5):564–9.

Churchward-Venne TA, Burd NA, Phillips SM. Nutritional regulation of muscle protein synthesis with resistance exercise: strategies to enhance anabolism. Nutr Metab (Lond). 2012; 9: 40.

Coggan AR, Swanson SC. Nutritional manipulations before and during endurance exercise: effects on performance. Med Sci Sports Exerc. 1992 Sep;24(9 Suppl):S331–5.

Colombani PC, Mannhart C, Mettler S. Carbohydrates and exercise performance in non-fasted athletes: a systematic review of studies mimicking real-life. Nutr J. 2013 Jan 28;12:16.

Cordain L, Eaton SB, Miller JB, Mann N, Hill K. The paradoxical nature of hunter-gatherer diets: meat-based, yet non-atherogenic. Eur J Clin Nutr. 2002 Mar;56 Suppl 1:S42–52. Review.

Coyle EF. Physiological determinants of endurance exercise performance. J Sci Med Sport. 1999;2(3):181–189.

Coyle EF, Coggan AR, Hemmert MK, Ivy JL. Muscle glycogen utilization during prolonged strenuous exercise when fed carbohydrate. J Appl Physiol (1985). 1986 Jul;61(1):165–72.

Coyle EF, Jeukendrup AE, Wagenmakers AJ, Saris WH. Fatty acid oxidation is directly regulated by carbohydrate metabolism during exercise. Am J Physiol. 1997 Aug;273(2 Pt 1):E268–75.

Cozma AI, Sievenpiper JL, de Souza RJ, Chiavaroli L, Ha V, Wang DD, Mirrahimi A, Yu ME, Carleton AJ ym.
Effect of fructose on glycemic control in diabetes: a systematic review and meta-analysis of controlled feeding trials. Diabetes Care. 2012 Jul;35(7):1611–20.

Currell K, Jeukendrup AE. Superior endurance performance with ingestion of multiple transportable carbohydrates. Med Sci Sports Exerc. 2008 Feb;40(2):275–81.

David Wang D, Sievenpiper JL, de Souza RJ, Cozma AI, Chiavaroli L, Ha V, Mirrahimi A, Carleton AJ ym. Effect of fructose on postprandial triglycerides: a systematic review and meta-analysis of controlled feeding trials. Atherosclerosis. 2014 Jan;232(1):125–33.

de Ataide e Silva T, Di Cavalcanti Alves de Souza ME, de Amorim JF, Stathis CG, Leandro CG, Lima-Silva AE.
Can carbohydrate mouth rinse improve performance during exercise? A systematic review. Nutrients. 2013 Dec 19;6(1):1–10.

de Koning L, Malik VS, Rimm EB, Willett WC, Hu FB. Sugar-sweetened and artificially sweetened beverage consumption and risk of type 2 diabetes in men. Am J Clin Nutr. 2011 Jun;93(6):1321–7.

de Ruyter JC, Olthof MR, Seidell JC, Katan MB. A trial of sugar-free or sugar-sweetened beverages and body weight in children. N Engl J Med. 2012 Oct 11;367(15):1397–406. (a)

de Ruyter JC, Katan MB, Kuijper LD, Liem DG, Olthof MR. The effect of sugar-free versus sugar-sweetened beverages on satiety, liking and wanting: an 18 month randomized double-blind trial in children. PLoS One. 2013 Oct 22;8(10):e78039 (b)

Doherty M, Smith PM. Effects of caffeine ingestion on rating of perceived exertion during and after exercise: a meta-analysis. Scand J Med Sci Sports. 2005;15(2):69–78.

Dolan LC, Potter SM, Burdock GA. Evidence-based review on the effect of normal dietary consumption of fructose on development of hyperlipidemia and obesity in healthy, normal weight individuals. Crit Rev Food Sci Nutr. 2010 Jan;50(1):53–84. (a)

- Dolan LC, Potter SM, Burdock GA.** Evidence-based review on the effect of normal dietary consumption of fructose on blood lipids and body weight of overweight and obese individuals. *Crit Rev Food Sci Nutr.* 2010 Nov;50(10):889–918. (b)
- Donaldson CM, Perry TL, Rose MC.** Glycemic index and endurance performance. *Int J Sport Nutr Exerc Metab.* 2010 Apr;20(2):154–65.
- Dong JY, Zhang L, Zhang YH, Qin LQ.** Dietary glycaemic index and glycaemic load in relation to the risk of type 2 diabetes: a meta-analysis of prospective cohort studies. *Br J Nutr.* 2011 Dec;106(11):1649–54.
- Dong JY, Zhang YH, Wang P, Qin LQ.** Meta-analysis of dietary glycemic load and glycemic index in relation to risk of coronary heart disease. *Am J Cardiol.* 2012 Jun 1;109(11):1608–13.
- Dougherty KA, Baker LB, Chow M, Kenney WL.** Two percent dehydration impairs and six percent carbohydrate drink improves boys basketball skills. *Med Sci Sports Exerc.* 2006 Sep;38(9):1650–8.
- Drenowatz C, Eisenmann JC, Carlson JJ, Pfeiffer KA, Pivarnik JM.** Energy expenditure and dietary intake during high-volume and low-volume training periods among male endurance athletes. *Appl Physiol Nutr Metab.* 2012 Apr;37(2):199–205.
- Dubé JJ, Allison KF, Rousson V, Goodpaster BH, Amati F.** Exercise dose and insulin sensitivity: relevance for diabetes prevention. *Med Sci Sports Exerc.* 2012 May;44(5):793–9.
- Egli L, LeCoultrre V, Theytaz F, Campos V, Hodson L, Schneiter P, Mittendorfer B, ym..** Exercise prevents fructose-induced hypertriglyceridemia in healthy young subjects. *Diabetes.* 2013 Jul;62(7):2259–65.
- Elfhag K, Tynelius P, Rasmussen F.** Sugar-sweetened and artificially sweetened soft drinks in association to restrained, external and emotional eating. *Physiol Behav.* 2007 Jun 8;91(2–3):191–5. **Ello-Martin JA, Ledikwe JH, Rolls BJ.** The influence of food portion size and energy density on energy intake: implications for weight management. *Am J Clin Nutr.* 2005 Jul;82(1 Suppl):236S-241S. Review.
- el-Sayed MS, Balmer J, Rattu AJ.** Carbohydrate ingestion improves endurance performance during a 1 h simulated cycling time trial. *J Sports Sci.* 1997 Apr;15(2):223–30.
- Erith S, Williams C, Stevenson E, Chamberlain S, Crews P, Rushbury I.** The effect of high carbohydrate meals with different glycemic indices on recovery of performance during prolonged intermittent high-intensity shuttle running. *Int J Sport Nutr Exerc Metab.* 2006 Aug;16(4):393–404.
- Esfahani A, Wong JM, Mirrahimi A, Villa CR, Kendall CW.** The application of the glycemic index and glycemic load in weight loss: A review of the clinical evidence. *IUBMB Life.* 2011 Jan;63(1):7–13.
- Eswaran S1, Muir J, Chey WD.** Fiber and functional gastrointestinal disorders. *Am J Gastroenterol.* 2013 May;108(5):718–27.
- Fares EJ, Kayser B.** Carbohydrate mouth rinse effects on exercise capacity in pre- and postprandial States. *J Nutr Metab.* 2011;2011:385962.
- Fenzl N, Bartsch K, Koenigstorfer J.** Labeling exercise fat-burning increases post-exercise food consumption in self-imposed exercisers. *Appetite.* 2014 May 28;81C:1–7.
- Ferguson-Stegall L, McCleave E, Ding Z, Doerner Iii PG, Liu Y, Wang B, Healy M, ym.** Aerobic exercise training adaptations are increased by postexercise carbohydrate-protein supplementation. *J Nutr Metab.* 2011;2011:623182.
- Figueiredo VC, Cameron-Smith D.** Is carbohydrate needed to further stimulate muscle protein synthesis/hypertrophy following resistance exercise? *J Int Soc Sports Nutr.* 2013 Sep 25;10(1):42.
- Finravinto 2012 -tutkimus. Heldan A, Raulio S, Kosola M, Tapanainen H, Ovaskainen M-L, Virtanen S.** THL Raportti 16/2013. Juvenes Print – Suomen Yliopistopaino Oy. Tampere 2013.
- Fogelholm M, Anderssen S, Gunnarsdottir I, Lahti-Koski M.** Dietary macronutrients and food consumption as determinants of long-term weight change in adult populations: a systematic literature review. *Food Nutr Res.* 2012;56.
- Fordtran JS.** Stimulation of active and passive sodium absorption by sugars in the human jejunum. *J Clin Invest.* 1975 Apr;55(4):728–37

Foskett A, Williams C, Boobis L, Tsintzas K. Carbohydrate availability and muscle energy metabolism during intermittent running. *Med Sci Sports Exerc.* 2008 Jan;40(1):96–103.

Gam S., Guelfi K.J., Fournier P.A. (2013) Opposition of carbohydrate in a mouth-rinse solution to the detrimental effect of mouth rinsing during cycling time trials. *International Journal of Sport Nutrition and Exercise Metabolism* 23, 48–56

Gardner C, Wylie-Rosett J, Gidding SS, Steffen LM, Johnson RK, Reader D, Lichtenstein AH; American Heart Association Nutrition Committee of the Council on Nutrition, Physical Activity and Metabolism, Council on Arteriosclerosis, Thrombosis and Vascular Biology, Council on Cardiovascular Disease in the Young; American Diabetes Association. Nonnutritive sweeteners: current use and health perspectives: a scientific statement from the American Heart Association and the American Diabetes Association. *Diabetes Care.* 2012 Aug;35(8):1798–808. Epub 2012 Jul 9.

Gibson GR, Probert HM, Loo JV, Rastall RA, Roberfroid MB. Dietary modulation of the human colonic microbiota: updating the concept of prebiotics. *Nutr Res Rev.* 2004 Dec;17(2):259–75.

Gleeson M, Williams C. Intense exercise training and immune function. *Nestle Nutr Inst Workshop Ser.* 2013;76:39–50.

Goedecke JH, White NJ, Chicktay W, Mahomed H, Durandt J, Lambert MI. The effect of carbohydrate ingestion on performance during a simulated soccer match. *Nutrients.* 2013 Dec 16;5(12):5193–204.

Goforth HW Jr, Laurent D, Prusaczyk WK, Schneider KE, Petersen KF, Shulman GI. Effects of depletion exercise and light training on muscle glycogen supercompensation in men. *Am J Physiol Endocrinol Metab.* 2003 Dec;285(6):E1304–11.

Goh Q, Boop CA, Luden ND, Smith AG, Womack CJ, Saunders MJ. Recovery from cycling exercise: effects of carbohydrate and protein beverages. *Nutrients.* 2012 Jul;4(7):568–84.

Grabitske HA, Slavin JL. Gastrointestinal effects of low-digestible carbohydrates. *Crit Rev Food Sci Nutr.* 2009 Apr;49(4):327–60.

Haff GG, Lehmkuhl MJ, McCoy LB, Stone MH. Carbohydrate supplementation and resistance training. *J Strength Cond Res.* 2003 Feb;17(1):187–96.

Hargreaves M, Hawley JA, Jeukendrup A. Pre-exercise carbohydrate and fat ingestion: effects on metabolism and performance. *J Sports Sci.* 2004 Jan;22(1):31–8. Review.

Hauner H, Bechthold A, Boeing H, Brönstrup A, Buyken A, Leschik-Bonnet E ym. ; German Nutrition Society. Evidence-based guideline of the German Nutrition Society: carbohydrate intake and prevention of nutrition-related diseases. *Ann Nutr Metab.* 2012;60 Suppl 1:1–58

Hawley JA. Nutritional strategies to modulate the adaptive response to endurance training. *Nestle Nutr Inst Workshop Ser.* 2013;75:1–14.

Hawley JA, Schabert EJ, Noakes TD, Dennis SC. Carbohydrate-loading and exercise performance. An update. *Sports Med.* 1997 Aug;24(2):73–81. Review.

Heinonen M. Ruuan lisääineet ja terveellisyys. *Duodecim.* 2014;130(7):683–8.

Highton J, Twist C, Lamb K, Nicholas C. Carbohydrate-protein coingestion improves multiple-sprint running performance. *J Sports Sci.* 2013;31(4):361–9.

Holway F. Team Sports. IOC Consensus In Sports Nutrition Conference Lausanne, Switzerland 25th–27th October 2010

Heesch MW, Mieras ME, Slivka DR. The performance effect of early versus late carbohydrate feedings during prolonged exercise. *Appl Physiol Nutr Metab.* 2014 Jan;39(1):58–63.

Hottenrott K, Hass E, Kraus M, Neumann G, Steiner M, Knechtle B. A scientific nutrition strategy improves time trial performance by $\approx 6\%$ when compared with a self-chosen nutrition strategy in trained cyclists: a randomized cross-over study. *Appl Physiol Nutr Metab.* 2012 Aug;37(4):637–45.

Howarth KR, Moreau NA, Phillips SM, Gibala MJ. Coingestion of protein with carbohydrate during recovery from endurance exercise stimulates skeletal muscle protein synthesis in humans. *J Appl Physiol (1985).* 2009 Apr;106(4):1394–402

Howarth KR, Phillips SM, MacDonald MJ, Richards D, Moreau NA, Gibala MJ. Effect of glycogen availability on human skeletal muscle protein turnover during exercise and recovery. *J Appl Physiol* (1985). 2010 Aug;109(2):431–8.

Hu FB. Are refined carbohydrates worse than saturated fat? *Am J Clin Nutr.* 2010 Jun;91(6):1541–2.

Hu FB. Resolved: there is sufficient scientific evidence that decreasing sugar-sweetened beverage consumption will reduce the prevalence of obesity and obesity-related diseases. *Obes Rev.* 2013 Aug;14(8):606–19.

Hulmi JJ, Volek JS, Selänne H, Mero AA. Protein ingestion prior to strength exercise affects blood hormones and metabolism. *Med Sci Sports Exerc.* 2005 Nov;37(11):1990–7.

Hulston CJ, Venables MC, Mann CH, Martin C, Philp A, Baar K, Jeukendrup AE. Training with low muscle glycogen enhances fat metabolism in well-trained cyclists. *Med Sci Sports Exerc.* 2010 Nov;42(11):2046–55.

Hulton AT, Gregson W, MacLaren D, Doran DA. Effects of GI meals on intermittent exercise. *Int J Sports Med.* 2012 Sep;33(9):756–62.

ICQC (International Carbohydrate Quality Consortium) Jenkins DJ, Willett WC, Astrup A ym. Glycaemic index: did Health Canada get it wrong? Position from the International Carbohydrate Quality Consortium (ICQC). *Br J Nutr.* 2014 Jan 28;111(2):380–2.

IOC. Nutrition Working Group of the International Olympic Committee. Nutrition for Athletes. A practical guide to eating for health and performance. Revised and updated 2012.

Jakobsen MU, Dethlefsen C, Joensen AM, Stegger J, Tjønneland A, Schmidt EB, Overvad K. Intake of carbohydrates compared with intake of saturated fatty acids and risk of myocardial infarction: importance of the glycemic index. *Am J Clin Nutr.* 2010 Jun;91(6):1764–8.

Jenkins DJ, Kendall CW, Augustin LS, Franceschi S, Hamidi M, Marchie A, Jenkins AL, Axelsen M. Glycemic index: overview of implications in health and disease. *Am J Clin Nutr.* 2002 Jul;76(1):266S–73S. Review.

Jenkins DJ, Srihaikul K, Kendall CW, Sievenpiper JL, Abdulnour S, Mirrahimi A, Meneses C, Nishi S ym. The relation of low glycaemic index fruit consumption to glycaemic control and risk factors for coronary heart disease in type 2 diabetes. *Diabetologia.* 2011 Feb;54(2):271–9.

Jenkins DJ, Kendall CW, Augustin LS, et al. (2012) Effect of legumes as part of a low glycemic index diet on glycemic control and cardiovascular risk factors in type 2 diabetes mellitus: a randomized controlled trial. *Arch Intern Med* 172, 1653–1660.

Jentjens R, Jeukendrup A. Determinants of post-exercise glycogen synthesis during short-term recovery. *Sports Med.* 2003;33(2):117–44. Review

Jentjens RL, Jeukendrup AE. High rates of exogenous carbohydrate oxidation from a mixture of glucose and fructose ingested during prolonged cycling exercise. *Br J Nutr.* 2005 Apr;93(4):485–92.

Jeukendrup AE. Nutrition for endurance sports: marathon, triathlon, road cycling. IOC Consensus In Sports Nutrition Conference Lausanne, Switzerland 25th–27th October 2010 (a)

Jeukendrup AE. Carbohydrate and exercise performance: the role of multiple transportable carbohydrates. *Curr Opin Clin Nutr Metab Care.* 2010 Jul;13(4):452–7. (b)

Jeukendrup AE. <http://www.nutritionworks.com.au/wp-content/uploads/2011/11/Carbohydrate.blends.pdf>

Jeukendrup AE. Oral carbohydrate rinse: placebo or beneficial? *Curr Sports Med Rep.* 2013 Jul-Aug;12(4):222–7.

Jeukendrup A, Brouns F, Wagenmakers AJ, Saris WH. Carbohydrate-electrolyte feedings improve 1 h time trial cycling performance. *Int J Sports Med.* 1997 Feb;18(2):125–9.

Jeukendrup AE, Jentjens R. Oxidation of carbohydrate feedings during prolonged exercise: current thoughts, guidelines and directions for future research. *Sports Med.* 2000 Jun;29(6):407–24.

Jeukendrup AE, Hopkins S, Aragón-Vargas LF, Hulston C. No effect of carbohydrate feeding on 16 km cycling time trial performance. *Eur J Appl Physiol.* 2008 Nov;104(5):831–7.

Jeukendrup AE, Killer SC. The myths surrounding pre-exercise carbohydrate feeding. *Ann Nutr Metab.* 2010;57 Suppl 2:18–25

Jeukendrup AE, Moseley L. Multiple transportable carbohydrates enhance gastric emptying and fluid delivery. *Scand J Med Sci Sports*. 2010 Feb;20(1):112–21.

Jeukendrup AE, McLaughlin J. Carbohydrate ingestion during exercise: effects on performance, training adaptations and trainability of the gut. *Nestle Nutr Inst Workshop Ser*. 2011;69:1–12; discussion 13–7.

Johnson RJ, Murray R. Fructose, exercise, and health. *Curr Sports Med Rep*. 2010 Jul-Aug;9(4):253–8.

Kang J, Raines E, Rosenberg J, Ratamess N, Naclerio F, Faigenbaum A. Metabolic responses during postprandial exercise. *Res Sports Med*. 2013;21(3):240–52.

Karamanolis IA, Laparidis KS, Volaklis KA, Douda HT, Tokmakidis SP. The effects of pre-exercise glycemic index food on running capacity. *Int J Sports Med*. 2011 Sep;32(9):666–71.

Kerksick C, Harvey T, Stout J, Campbell B, Wilborn C, Kreider R, Kalman D, et al. International Society of Sports Nutrition position stand: nutrient timing. *J Int Soc Sports Nutr*. 2008; 5: 17.

Koh-Banerjee P, Franz M, Sampson L, Liu S, Jacobs DR, Jr, Spiegelman D, et al. Changes in whole-grain, bran, and cereal fiber consumption in relation to 8-y weight gain among men. *Am J Clin Nutr*. 2004;80(5):1237–45.

Kulik JR, Touchberry CD, Kawamori N, Blumert PA, Crum AJ, Haff GG. Supplemental carbohydrate ingestion does not improve performance of high-intensity resistance exercise. *J Strength Cond Res*. 2008 Jul;22(4):1101–7.

Lane SC, Bird SR, Burke LM, Hawley JA. Effect of a carbohydrate mouth rinse on simulated cycling time-trial performance commenced in a fed or fasted state. *Appl Physiol Nutr Metab*. 2013 Feb;38(2):134–9.

Leiper J.B., Broad N.P., Maughan R.J. (2001) Effect of intermittent high-intensity exercise on gastric emptying in man. *Medicine and Science in Sports and Exercise* 33, 1270–1278

Lemon PW, Nagle FJ. Effects of exercise on protein and amino acid metabolism. *Med Sci Sports Exerc*. 1981;13(3):141–9.

Lieberman HR, Falco CM, Slade SS. Carbohydrate administration during a day of sustained aerobic activity improves vigilance, as assessed by a novel ambulatory monitoring device, and mood. *Am J Clin Nutr*. 2002 Jul;76(1):120–7.

Lima-Silva AE, Pires FO, Bertuzzi R, Silva-Cavalcante MD, Oliveira RS, Kiss MA, Bishop D. Effects of a low- or a high-carbohydrate diet on performance, energy system contribution, and metabolic responses during supramaximal exercise. *Appl Physiol Nutr Metab*. 2013 Sep;38(9):928–34.

Little JP, Chilibeck PD, Ciona D, Forbes S, Rees H, Vandenberg A, Zello GA. Effect of low- and high-glycemic-index meals on metabolism and performance during high-intensity, intermittent exercise. *Int J Sport Nutr Exerc Metab*. 2010 Dec;20(6):447–56.

Livesey G, Taylor R, Livesey H, Liu S. Is there a dose-response relation of dietary glycemic load to risk of type 2 diabetes? Meta-analysis of prospective cohort studies. *Am J Clin Nutr*. 2013 Mar;97(3):584–96.

Lomax AR, Calder PC. Prebiotics, immune function, infection and inflammation: a review of the evidence. *Br J Nutr*. 2009 Mar;101(5):633–58.

Lunn WR, Pasiakos SM, Colletto MR, Karfonta KE, Carbone JW, Anderson JM, Rodriguez NR. Chocolate milk and endurance exercise recovery: protein balance, glycogen, and performance. *Med Sci Sports Exerc*. 2012 Apr;44(4):682–91.

Malik VS, Popkin BM, Bray GA, Després JP, Willett WC, Hu FB. Sugar-sweetened beverages and risk of metabolic syndrome and type 2 diabetes: a meta-analysis. *Diabetes Care*. 2010 Nov;33(11):2477–83.

Malik VS, Pan A, Willett WC, Hu FB. Sugar-sweetened beverages and weight gain in children and adults: a systematic review and meta-analysis. *Am J Clin Nutr*. 2013 Oct;98(4):1084–102.

Mattes RD, Popkin BM. Nonnutritive sweetener consumption in humans: effects on appetite and food intake and their putative mechanisms. *Am J Clin Nutr*. 2009 Jan;89(1):1–14.

Matsui T, Soya S, Okamoto M, Ichitani Y, Kawanaka K, Soya H. Brain glycogen decreases during prolonged exercise. *J Physiol*. 2011 Jul 1;589(Pt 13):3383–93.

Maughan RJ. Fundamentals of Sports Nutrition: Application to Sports Drinks. Kirjassa: Sports Drinks – Basic Science and Practical Aspects. Maughan RJ, Murray R (toim.). CRC Press. Boca Raton, USA, 2001.

Maughan RJ, Meyer NL. Hydration during intense exercise training. *Nestle Nutr Inst Workshop Ser.* 2013;76:25–37.

McConell G, Kloot K, Hargreaves M. Effect of timing of carbohydrate ingestion on endurance exercise performance. *Med Sci Sports Exerc.* 1996 Oct;28(10):1300–4.

McGawley K, Shannon O, Betts J. Ingesting a high-dose carbohydrate solution during the cycle section of a simulated Olympic-distance triathlon improves subsequent run performance. *Appl Physiol Nutr Metab.* 2012 Aug;37(4):664–71.

McGlory C, Morton JP. The effects of postexercise consumption of high-molecular-weight versus low-molecular-weight carbohydrate solutions on subsequent high-intensity interval-running capacity. *Int J Sport Nutr Exerc Metab.* 2010 Oct;20(5):361–9.

McLellan TM, Pasiakos SM, Lieberman HR. Effects of protein in combination with carbohydrate supplements on acute or repeat endurance exercise performance: a systematic review. *Sports Med.* 2014 Apr;44(4):535–50.

McRae KA, Galloway SD. Carbohydrate-electrolyte drink ingestion and skill performance during and after 2 hr of indoor tennis match play. *Int J Sport Nutr Exerc Metab.* 2012 Feb;22(1):38–46.

Mensink RP, Zock PL, Kester AD, Katan MB. Effects of dietary fatty acids and carbohydrates on the ratio of serum total to HDL cholesterol and on serum lipids and apolipoproteins: a meta-analysis of 60 controlled trials. *Am J Clin Nutr.* 2003 May;77(5):1146–55.

Marinovich M, Galli CL, Bosetti C, Gallus S, La Vecchia C. Aspartame, low-calorie sweeteners and disease: regulatory safety and epidemiological issues. *Food Chem Toxicol.* 2013 Oct;60:109–15

Miller PE, Perez V. Low-calorie sweeteners and body weight and composition: a meta-analysis of randomized controlled trials and prospective cohort studies. *Am J Clin Nutr.* 2014 Jun 18.

Mitchell JB. Ingestion of carbohydrate during recovery in exercising people. *Curr Opin Clin Nutr Metab Care.* 2013 Jul;16(4):461–5

Mitchell JB, DiLauro PC, Pizza FX, Cavender DL. The effect of preexercise carbohydrate status on resistance exercise performance. *Int J Sport Nutr.* 1997 Sep;7(3):185–96.

Moore LJ, Midgley AW, Thurlow S, Thomas G, McNaughton LR. Effect of the glycaemic index of a pre-exercise meal on metabolism and cycling time trial performance. *J Sci Med Sport.* 2010 Jan;13(1):182–8

Moore LJ, Midgley A, Vince R, McNaughton LR. The effects of low and high glycemic index 24-h recovery diets on cycling time trial performance. *J Sports Med Phys Fitness.* 2011 Jun;51(2):233–40.

Moore DR, Areta J, Coffey VG, Stellingwerff T, Phillips SM, Burke LM, Cléroux M, Godin JP, Hawley JA. Daytime pattern of post-exercise protein intake affects whole-body protein turnover in resistance-trained males. *Nutr Metab (Lond).* 2012 Oct 16;9(1):91.

Moore L, Szpalek HM, McNaughton LR. Preexercise high and low glycemic index meals and cycling performance in untrained females: randomized, cross-over trial of efficacy. *Res Sports Med.* 2013;21(1):24–36.

Morton JP, Croft L, Bartlett JD, Maclare DP, Reilly T, Evans L, McArdle A, Drust B. Reduced carbohydrate availability does not modulate training-induced heat shock protein adaptations but does upregulate oxidative enzyme activity in human skeletal muscle. *J Appl Physiol (1985).* 2009 May;106(5):1513–21.

Mozaffarian D, Hao T, Rimm EB, Willett WC, Hu FB. Changes in diet and lifestyle and long-term weight gain in women and men. *N Engl J Med.* 2011 Jun 23;364(25):2392–404

Nicholas CW, Nuttall FE, Williams C. The Loughborough Intermittent Shuttle Test: a field test that simulates the activity pattern of soccer. *J Sports Sci.* 2000 Feb;18(2):97–104.

Noakes TD. Fatigue is a Brain-Derived Emotion that Regulates the Exercise Behavior to Ensure the Protection of Whole Body Homeostasis. *Front Physiol.* 2012 Apr 11;3:82.

Nybo L. CNS fatigue and prolonged exercise: effect of glucose supplementation. *Med Sci Sports Exerc.* 2003 Apr;35(4):589–94.

Nybo L, Møller K, Pedersen BK, Nielsen B, Secher NH. Association between fatigue and failure to preserve cerebral energy turnover during prolonged exercise. *Acta Physiol Scand.* 2003 Sep;179(1):67–74.

O'Brien WJ, Rowlands DS. Fructose-maltodextrin ratio in a carbohydrate-electrolyte solution differentially affects exogenous carbohydrate oxidation rate, gut comfort, and performance. *Am J Physiol Gastrointest Liver Physiol.* 2011 Jan;300(1):G181–9.

O'Brien WJ, Stannard SR, Clarke JA, Rowlands DS. Fructose-maltodextrin ratio governs exogenous and other CHO oxidation and performance. *Med Sci Sports Exerc.* 2013 Sep;45(9):1814–24

Ojuka EO, Jones TE, Han DH, et al. Intermittent increases in cytosolic Ca²⁺ stimulate mitochondrial biogenesis in muscle cells. *Am J Physiol.* 2002;283(5):E1040–E1045.

Onywera VO, Kiplamai FK, Boit MK, Pitsiladis YP. Food and macronutrient intake of elite kenyan distance runners. *Int J Sport Nutr Exerc Metab.* 2004 Dec;14(6):709–19.

O'Reilly J1, Wong SH, Chen Y. Glycaemic index, glycaemic load and exercise performance. *Sports Med.* 2010 Jan 1;40(1):27–39.

Pal S, Radavelli-Bagatini S. Effects of psyllium on metabolic syndrome risk factors. *Obes Rev.* 2012 Nov;13(11):1034–47.

Pal S, Khossousi A, Binns C, et al. The effect of a fibre supplement compared to a healthy diet on body composition, lipids, glucose, insulin and other metabolic syndrome risk factors in overweight and obese individuals. *Br J Nutr.* 2011;105:90–100.

Pan A, Hu FB. Effects of carbohydrates on satiety: differences between liquid and solid food. *Curr Opin Clin Nutr Metab Care.* 2011 Jul;14(4):385–90.

Pan A, Malik VS, Hao T, Willett WC, Mozaffarian D, Hu FB. Changes in water and beverage intake and long-term weight changes: results from three prospective cohort studies. *Int J Obes (Lond).* 2013 Oct;37(10):1378–85.

Paoli A, Marcolin G, Zonin F, Neri M, Sivieri A, Pacelli QF. Exercising fasting or fed to enhance fat loss? Influence of food intake on respiratory ratio and excess postexercise oxygen consumption after a bout of endurance training. *Int J Sport Nutr Exerc Metab.* 2011 Feb;21(1):48–54.

Parnell JA, Reimer RA. Prebiotic fiber modulation of the gut microbiota improves risk factors for obesity and the metabolic syndrome. *Gut Microbes.* 2012 Jan-Feb;3(1):29–34.

Patterson SD, Gray SC. Carbohydrate-gel supplementation and endurance performance during intermittent high-intensity shuttle running. *Int J Sport Nutr Exerc Metab.* 2007 Oct;17(5):445–55.

Peternelj TT, Coombes JS. Antioxidant supplementation during exercise training: beneficial or detrimental? *Sports Med.* 2011;41(12):1043–1069

Peters HP, van Schelven FW, Verstappen PA, de Boer RW, Bol E, Erich WB, van der Togt CR, de Vries WR. Gastrointestinal problems as a function of carbohydrate supplements and mode of exercise. *Med Sci Sports Exerc.* 1993 Nov;2

Pfeiffer B, Stellingwerff T, Zaltas E, Jeukendrup AE. CHO oxidation from a CHO gel compared with a drink during exercise. *Med Sci Sports Exerc.* 2010 Nov;42(11):2038–45.5(11):1211–24. (a)

Pfeiffer B, Stellingwerff T, Zaltas E, Jeukendrup AE. Oxidation of solid versus liquid CHO sources during exercise. *Med Sci Sports Exerc.* 2010 Nov;42(11):2030–7.

Pfeiffer B, Stellingwerff T, Hodgson AB, Jeukendrup AE. Carbohydrate oxidation from a drink during running compared with cycling exercise. *Med Sci Sports Exerc.* 2011 Feb;43(2):327–34.

Pfeiffer B, Stellingwerff T, Hodgson AB, Randell R, Pöttgen K, Res P, Jeukendrup AE. Nutritional intake and gastrointestinal problems during competitive endurance events. *Med Sci Sports Exerc.* 2012 Feb;44(2):344–51.

Phillips SM, Sproule J, Turner AP. Carbohydrate ingestion during team games exercise: current knowledge and areas for future investigation. *Sports Med.* 2011 Jul 1;41(7):559–85.

Phillips SM, Turner AP, Sanderson MF, Sproule J. Carbohydrate gel ingestion significantly improves the intermittent endurance capacity, but not sprint performance, of adolescent team games players during a simulated team games protocol. *Eur J Appl Physiol.* 2012 Mar;112(3):1133–41.

Phillips SM, Turner AP, Gray S, Sanderson MF, Sproule J. Ingesting a 6% carbohydrate-electrolyte solution improves endurance capacity, but not sprint performance, during intermittent, high-intensity shuttle running in adolescent team games players aged 12–14 years. *Eur J Appl Physiol*. 2010 Jul;109(5):811–21

Phillips SM, Findlay S, Kavaliauskas M, Grant MC. The Influence of Serial Carbohydrate Mouth Rinsing on Power Output during a Cycle Sprint. *J Sports Sci Med*. 2014 May 1;13(2):252–8. eCollection 2014.

Philp A, Hargreaves M, Baar K. More than a store: regulatory roles for glycogen in skeletal muscle adaptation to exercise. *Am J Physiol Endocrinol Metab*. 2012 Jun 1;302(11):E1343–51.

Piehl Aulin K, Söderlund K, Hultman E. Muscle glycogen resynthesis rate in humans after supplementation of drinks containing carbohydrates with low and high molecular masses. *Eur J Appl Physiol*. 2000 Mar;81(4):346–51.

Piernas C, Tate DF, Wang X, Popkin BM. Does diet-beverage intake affect dietary consumption patterns? Results from the Choose Healthy Options Consciously Everyday (CHOICE) randomized clinical trial. *Am J Clin Nutr*. 2013 Mar;97(3):604–11

Pritchett K, Pritchett R. Chocolate milk: a post-exercise recovery beverage for endurance sports. *Med Sport Sci*. 2012;59:127–34

Psilander N, Frank P, Flockhart M, Sahlin K. Exercise with low glycogen increases PGC-1 gene expression in human skeletal muscle. *Eur J Appl Physiol*. 2013 Apr;113(4):951–63.

Raben A, Vasilaras TH, Møller AC, Astrup AA. Sucrose compared with artificial sweeteners: different effects on ad libitum food intake and body weight after 10 wk of supplementation in overweight subjects. *Am J Clin Nutr*. 2002;76:721–9.

Raman A, Macdermid PW, Mündel T, Mann M, Stannard SR. The Effects of Carbohydrate Loading 48 Hours Prior to a Simulated Squash Match. *Int J Sport Nutr Exerc Metab*. 2013 Oct 2.

Rauch HG, St Clair Gibson A, Lambert EV, Noakes TD. A signalling role for muscle glycogen in the regulation of pace during prolonged exercise. *Br J Sports Med*. 2005 Jan;39(1):34–8.

Rollo I, Williams C, Nevill M. Influence of ingesting versus mouth rinsing a carbohydrate solution during a 1-h run. *Med Sci Sports Exerc*. 2011 Mar;43(3):468–75.

Rolls BJ. The relationship between dietary energy density and energy intake. *Physiol Behav*. 2009;97:609–15.

Romijn JA, Coyle EF, Sidossis LS, Zhang XJ, Wolfe RR. Relationship between fatty acid delivery and fatty acid oxidation during strenuous exercise. *J Appl Physiol* (1985). 1995 Dec;79(6):1939–45.

Romijn JA, Coyle EF, Sidossis LS, Rosenblatt J, Wolfe RR. Substrate metabolism during different exercise intensities in endurance-trained women. *J Appl Physiol* (1985). 2000 May;88(5):1707–14.

Rowlands DS, Wallis GA, Shaw C, Jentjens RL, Jeukendrup AE. Glucose polymer molecular weight does not affect exogenous carbohydrate oxidation. *Med Sci Sports Exerc*. 2005 Sep;37(9):1510–6.

Rowlands DS, Swift M, Ros M, Green JG. Composite versus single transportable carbohydrate solution enhances race and laboratory cycling performance. *Appl Physiol Nutr Metab*. 2012 Jun;37(3):425–36.

Russell M, Kingsley M. The Efficacy of Acute Nutritional Interventions on Soccer Skill Performance. *Sports Med*. 2014 Apr 12.

Satija A, Hu FB. Cardiovascular benefits of dietary fiber. *Curr Atheroscler Rep*. 2012 Dec;14(6):505–14.

Saunders MJ. Coingestion of carbohydrate-protein during endurance exercise: influence on performance and recovery. *Int J Sport Nutr Exerc Metab*. 2007 Aug;17 Suppl:S87–103.

Saunders MJ. Carbohydrate-protein intake and recovery from endurance exercise: is chocolate milk the answer? *Curr Sports Med Rep*. 2011 Jul;10(4):203–10.

Sedlock DA. The latest on carbohydrate loading: a practical approach. *Curr Sports Med Rep*. 2008 Jul-Aug;7(4):209–13.

Shiraev T, Barclay G. Evidence based exercise – clinical benefits of high intensity interval training. *Aust Fam Physician*. 2012 Dec;41(12):960–2.

Sievenpiper JL, de Souza RJ, Mirrahimi A, Yu ME, Carleton AJ, Beyene J, Chiavaroli L, Di Buono M et al. Effect of fructose on body weight in controlled feeding trials: a systematic review and meta-analysis. *Ann Intern Med.* 2012 Feb 21;156(4):291–304. (a)

Sievenpiper JL, Chiavaroli L, de Souza RJ, Mirrahimi A, Cozma AI, Ha V, Wang DD, Yu ME, et al. ‘Catalytic’ doses of fructose may benefit glycaemic control without harming cardiometabolic risk factors: a small meta-analysis of randomised controlled feeding trials. *Br J Nutr.* 2012 Aug;108(3):418–23. (b)

Sinclair J, Bottoms L, Flynn C, Bradley E, Alexander G, McCullagh S, Finn T, Hurst HT. The effect of different durations of carbohydrate mouth rinse on cycling performance. *Eur J Sport Sci.* 2014 Apr;14(3):259–64.

Siri-Tarino PW, Sun Q, Hu FB, Krauss RM. Saturated fat, carbohydrate, and cardiovascular disease. *Am J Clin Nutr.* 2010 Mar;91(3):502–9.

Skein M, Duffield R, Kelly BT, Marino FE. The effects of carbohydrate intake and muscle glycogen content on self-paced intermittent-sprint exercise despite no knowledge of carbohydrate manipulation. *Eur J Appl Physiol.* 2012 Aug;112(8):2859–70.

Slater G. Nutrition for strength sports. IOC Consensus In Sports Nutrition Conference Lausanne, Switzerland 25th–27th October 2010

Slater G, Phillips SM. Nutrition guidelines for strength sports: sprinting, weightlifting, throwing events, and bodybuilding. *J Sports Sci.* 2011;29 Suppl 1:S67–77.

Sluijs I, Beulens JW, van der Schouw YT, van der A DL, Buckland G, Kuijsten A et al.; InterAct consortium. Dietary glycemic index, glycemic load, and digestible carbohydrate intake are not associated with risk of type 2 diabetes in eight European countries. *J Nutr.* 2013 Jan;143(1):93–9.

Smith JW, Pascoe DD, Passe DH, Ruby BC, Stewart LK, Baker LB, Zachwieja JJ. Curvilinear dose-response relationship of carbohydrate (0–120 g·h⁻¹) and performance. *Med Sci Sports Exerc.* 2013 Feb;45(2):336–41.

Soop M, Nehra V, Henderson GC, Boirie Y, Ford GC, Nair KS. Coingestion of whey protein and casein in a mixed meal: demonstration of a more sustained anabolic effect of casein. *Am J Physiol Endocrinol Metab.* 2012 Jul 1;303(1):E152–62.

Souglis AG, Chryssanthopoulos CI, Travlos AK, Zorzou AE, Gassis IT, Papadopoulos CN, Sotiropoulos AA. The effect of high vs. low carbohydrate diets on distances covered in soccer. *J Strength Cond Res.* 2013 Aug;27(8):2235–47.

Spaccarella KJ, Andzel WD. The effects of low fat chocolate milk on postexercise recovery in collegiate athletes. *J Strength Cond Res.* 2011 Dec;25(12):3456–60.

Stearns RL, Emmanuel H, Volek JS, Casa DJ. Effects of ingesting protein in combination with carbohydrate during exercise on endurance performance: a systematic review with meta-analysis. *J Strength Cond Res.* 2010 Aug;24(8):2192–202.

Stellingwerff T. Contemporary nutrition approaches to optimize elite marathon performance. *Int J Sports Physiol Perform.* 2013 Sep;8(5):573–8. Epub 2013 Apr 9. Review.

Stellingwerff T, Spriet LL, Watt MJ, Kimber NE, Hargreaves M, Hawley JA, Burke LM. Decreased PDH activation and glycogenolysis during exercise following fat adaptation with carbohydrate restoration. *Am J Physiol Endocrinol Metab.* 2006 Feb;290(2):E380–8.

Stellman SD, Garfinkel L. Artificial sweetener use and one-year weight change among women. *Prev Med* 1986;15:195–202

Stephens FB, Roig M, Armstrong G, Greenhaff PL. Post-exercise ingestion of a unique, high molecular weight glucose polymer solution improves performance during a subsequent bout of cycling exercise. *J Sports Sci.* 2008 Jan 15;26(2):149–54

Stevenson L, Phillips F, O’Sullivan K, Walton J. Wheat bran: its composition and benefits to health, a European perspective. *Int J Food Sci Nutr.* 2012 Dec;63(8):1001–13.

Strobel NA, Peake JM, Matsumoto A, et al. Antioxidant supplementation reduces skeletal muscle mitochondrial biogenesis. *Med Sci Sports Exerc.* 2011;43(6):1017–1024.

Sullo A, Monda M, Brizzi G, Meninno V, Papa A, Lombardi P, Fabbri B. The effect of a carbohydrate loading on running performance during a 25-km treadmill time trial by level of aerobic capacity in athletes. *Eur Rev Med Pharmacol Sci.* 1998 Sep-Dec;2(5–6):195–202.

Sørensen LB, Vasilaras TH, Astrup A, Raben A. Sucrose compared with artificial sweeteners: a clinical intervention study of effects on energy intake, appetite, and energy expenditure after 10 wk of supplementation in overweight subjects. *Am J Clin Nutr.* 2014 Apr 30.

Tate DF, Turner-McGrievy G, Lyons E, Stevens J, Erickson K, Polzien K, Diamond M, Wang X, Popkin B. Replacing caloric beverages with water or diet beverages for weight loss in adults: main results of the Choose Healthy Options Consciously Everyday (CHOICE) randomized clinical trial. *Am J Clin Nutr.* 2012 Mar;95(3):555–63.

Taylor C, Bartlett JD, van de Graaf CS, Louhelainen J, Coyne V, Iqbal Z, Maclarens DP, ym. Protein ingestion does not impair exercise-induced AMPK signalling when in a glycogen-depleted state: implications for train-low compete-high. *Eur J Appl Physiol.* 2013 Jun;113(6):1457–68.

Te Morenga L, Mallard S, Mann J. Dietary sugars and body weight: systematic review and meta-analyses of randomised controlled trials and cohort studies. *BMJ.* 2012 Jan 15;346:e7492.

Te Morenga LA, Howatson AJ, Jones RM, Mann J. Dietary sugars and cardiometabolic risk: systematic review and meta-analyses of randomized controlled trials of the effects on blood pressure and lipids. *Am J Clin Nutr.* 2014 May 7;100(1):65–79.

Theuwissen E, Mensink RP. Water-soluble dietary fibers and cardiovascular disease. *Physiol Behav.* 2008 May 23;94(2):285–92.

Threapleton DE, Greenwood DC, Evans CE, Cleghorn CL, Nykjaer C, Woodhead C, Cade JE, ym. Dietary fibre intake and risk of cardiovascular disease: systematic review and meta-analysis. *BMJ.* 2013 Dec 19;347:f6879.

Triplett D, Doyle JA, Rupp JC, Benardot D. An isocaloric glucose-fructose beverage's effect on simulated 100-km cycling performance compared with a glucose-only beverage. *Int J Sport Nutr Exerc Metab.* 2010 Apr;20(2):122–31.

Tucker LA, Thomas KS. Increasing total fiber intake reduces risk of weight and fat gains in women. *J Nutr.* 2009;139(3):576–81.

U.S. Department of Agriculture, U.S. Department of Health. Dietary Guidelines for Americans 2010. www.dietaryguidelines.gov

Valentine RJ, Saunders MJ, Todd MK, St Laurent TG. Influence of carbohydrate-protein beverage on cycling endurance and indices of muscle disruption. *Int J Sport Nutr Exerc Metab.* 2008 Aug;18(4):363–78.

Vandenbogaerde TJ, Hopkins WG. Effects of acute carbohydrate supplementation on endurance performance: a meta-analysis. *Sports Med.* 2011 Sep 1;41(9):773–92.

Van Proeyen K, Szlufcik K, Nielens H, Ramaekers M, Hespel P. Beneficial metabolic adaptations due to endurance exercise training in the fasted state. *J Appl Physiol (1985).* 2011 Jan;110(1):236–45.

Vega-López S, Mayol-Kreiser SN. Use of the glycemic index for weight loss and glycemic control: a review of recent evidence. *Curr Diab Rep.* 2009 Oct;9(5):379–88. Review.

Vergauwen L1, Brouns F, Hespel P. Carbohydrate supplementation improves stroke performance in tennis. *Med Sci Sports Exerc.* 1998 Aug;30(8):1289–95.

Vos MB, Lavine JE. Dietary fructose in nonalcoholic fatty liver disease. *Hepatology.* 2013 Jun;57(6):2525–31.

VRN. Terveytä ruuasta – Suomalaiset ravitsemussuositukset 2014. Juvenes Oy, Helsinki 2014.

Walsh NP, Gleeson M, Pyne DB, Nieman DC, Dhabhar FS, Shephard RJ, Oliver SJ, Bermon S, Kajeniene A. Position statement. Part two: Maintaining immune health. *Exerc Immunol Rev.* 2011;17:64–103. Review.

Wanders AJ, van den Borne JJ, de Graaf C, Hulshof T, Jonathan MC, Kristensen M, Mars M, ym. Effects of dietary fibre on subjective appetite, energy intake and body weight: a systematic review of randomized controlled trials. *Obes Rev.* 2011 Sep;12(9):724–39.

Weir JP, Beck TW, Cramer JT, Housh TJ. Is fatigue all in your head? A critical review of the central governor model. *Br J Sports Med.* 2006 Jul;40(7):573–86; discussion 586. Review.

Welsh RS, Davis JM, Burke JR, Williams HG. Carbohydrates and physical/mental performance during intermittent exercise to fatigue. *Med Sci Sports Exerc.* 2002 Apr;34(4):723–31.

Whitney EN, Rolfes SR. The carbohydrates: Sugars, starch and fibers. Kirjassa: Whitney EN, Rolfes SR (toim.). Understanding Nutrition 7th ed. West Publishing Company, USA, 1996

WHO <http://www.who.int/mediacentre/news/notes/2014/consultation-sugar-guideline/en/>

Winnick JJ, Davis JM, Welsh RS, Carmichael MD, Murphy EA, Blackmon JA. Carbohydrate feedings during team sport exercise preserve physical and CNS function. *Med Sci Sports Exerc.* 2005 Feb;37(2):306–15.

Wolever TM. Relationship between dietary fiber content and composition in foods and the glycemic index. *Am J Clin Nutr.* 1990;51(1):72–5.

Wolever TM, Jenkins DJ, Jenkins AL, Josse RG. The glycemic index: methodology and clinical implications. *Am J Clin Nutr.* 1991 Nov;54(5):846–54.

Wolever TM, Gibbs AL, Mehling C, et al. (2008) The Canadian Trial of Carbohydrates in Diabetes (CCD), a 1-y controlled trial of low-glycemic-index dietary carbohydrate in type 2 diabetes: no effect on glycated hemoglobin but reduction in C-reactive protein. *Am J Clin Nutr* 87, 114–125

Wolever TM. Is glycaemic index (GI) a valid measure of carbohydrate quality? *Eur J Clin Nutr.* 2013 May;67(5):522–31

Yaghoobi N, Al-Waili N, Ghayour-Mobarhan M, Parizadeh SM, Abasalti Z, Yaghoobi Z, Yaghoobi F ym. Natural honey and cardiovascular risk factors; effects on blood glucose, cholesterol, triacylglycerole, CRP, and body weight compared with sucrose. *ScientificWorldJournal.* 2008 Apr 20;8:463–9.

Yao B, Fang H, Xu W, Yan Y, Xu H, Liu Y, Mo M, Zhang H, Zhao Y. Dietary fiber intake and risk of type 2 diabetes: a dose-response analysis of prospective studies. *Eur J Epidemiol.* 2014 Feb;29(2):79–88.

Ye EQ, Chacko SA, Chou EL, Kugizaki M, Liu S. Greater whole-grain intake is associated with lower risk of type 2 diabetes, cardiovascular disease, and weight gain. *J Nutr.* 2012 Jul;142(7):1304–13.

Yeo WK, Paton CD, Garnham AP, Burke LM, Carey AL, Hawley JA. Skeletal muscle adaptation and performance responses to once a day versus twice every second day endurance training regimens. *J Appl Physiol (1985).* 2008 Nov;105(5):1462–70.

Zajac A, Poprzecki S, Maszczyk A, Czuba M, Michalczuk M, Zydek G. The effects of a ketogenic diet on exercise metabolism and physical performance in off-road cyclists. *Nutrients.* 2014 Jun 27;6(7):2493–508.

Åkermark C, Jacobs I, Rasmusson M, Karlsson J. Diet and muscle glycogen concentration in relation to physical performance in Swedish elite ice hockey players. *Int J Sport Nutr.* 1996 Sep;6(3):272–84.

Lähdeluettelo luku 6

Acheson KJ, Blondel-Lubrano A, Oguey-Araymon S, Beaumont M, Emady-Azar S, Ammon-Zufferey C, ym. Protein choices targeting thermogenesis and metabolism. *Am J Clin Nutr.* 2011 Mar;93(3):525–34.

Andersen LL, Tufekovic G, Zebis MK, Crameri RM, Verlaan G, Kjaer M, Suetta C ym. The effect of resistance training combined with timed ingestion of protein on muscle fiber size and muscle strength. *Metabolism.* 2005 Feb;54(2):151–6.

Antonio J, Peacock CA, Ellerbroek A, Fromhoff B, Silver T. The effects of consuming a high protein diet (4.4 g/kg/d) on body composition in resistance-trained individuals. *J Int Soc Sports Nutr.* 2014 May 12;11:19.

Aragon AA, Schoenfeld BJ. Nutrient timing revisited: is there a post-exercise anabolic window? *J Int Soc Sports Nutr.* 2013 Jan 29;10(1):5.

Arciero PJ, Ormsbee MJ, Gentile CL, Nindl BC, Brestoff JR, Ruby M. Increased protein intake and meal frequency reduces abdominal fat during energy balance and energy deficit. *Obesity (Silver Spring).* 2013 Jul;21(7):1357–66

- Areta JL, Burke LM, Ross ML, Camera DM, West DW, Broad EM, Jeacocke NA, Moore DR, Stellingwerff T, ym.** Timing and distribution of protein ingestion during prolonged recovery from resistance exercise alters myofibrillar protein synthesis. *J Physiol.* 2013 May 1;591(Pt 9):2319–31.
- Atkins RC.** Dr. Atkins' diet revolution: the high calorie way to stay thin forever. New York: David McKay, 1972.
- Beelen M, Koopman R, Gijsen AP, Vandereyt H, Kies AK, Kuipers H, Saris WH, van Loon LJ.** Protein coingestion stimulates muscle protein synthesis during resistance-type exercise. *Am J Physiol Endocrinol Metab.* 2008 Jul;295(1):E70–7.
- Betts JA, Williams C.** Short-term recovery from prolonged exercise: exploring the potential for protein ingestion to accentuate the benefits of carbohydrate supplements. *Sports Med.* 2010 Nov 1;40(11):941–59
- Betts JA, Beelen M, Stokes KA, Saris W HM, van Loon L JC.** Endocrine responses during overnight recovery from exercise: impact of nutrition and relationships with muscle protein synthesis. *Int J Sport Nutr Exerc Metab.* 2011 Oct;21(5):398–409. Epub 2011 Aug 10.
- Bilsborough S, Mann N.** A review of issues of dietary protein intake in humans. *Int J Sport Nutr Exerc Metab.* 2006 Apr;16(2):129–52.
- Biolo G, Maggi SP, Williams BD, Tipton KD, Wolfe RR.** Increased rates of muscle protein turnover and amino acid transport after resistance exercise in humans. *Am J Physiol.* 1995 Mar;268(3 Pt 1):E514–20.
- Biolo G, Tipton KD, Klein S, Wolfe RR.** An abundant supply of amino acids enhances the metabolic effect of exercise on muscle protein. *Am J Physiol.* 1997 Jul;273(1 Pt 1):E122–9.
- Blatt AD, Roe LS, Rolls BJ.** Increasing the protein content of meals and its effect on daily energy intake. *J Am Diet Assoc* 2011; 111: 290–4.
- Boirie Y, Dangin M, Gachon P, Vasson MP, Maubois JL, Beaufre B.** Slow and fast dietary proteins differently modulate postprandial protein accretion. *Proc Natl Acad Sci U S A.* 1997 Dec 23;94(26):14930–5.
- Bosse JD, Dixon BM.** Dietary protein to maximize resistance training: a review and examination of protein spread and change theories. *J Int Soc Sports Nutr.* 2012 Sep 8;9(1):42.
- Bray GA, Smith SR, de Jonge L, Xie H, Rood J, Martin CK, Most M, Brock C, Mancuso S, Redman LM.** Effect of dietary protein content on weight gain, energy expenditure, and body composition during overeating: a randomized controlled trial. *JAMA.* 2012 Jan 4;307(1):47–55.
- Brechue WF, Abe T.** The role of FFM accumulation and skeletal muscle architecture in powerlifting performance. *Eur J Appl Physiol.* 2002 Feb;86(4):327–36.
- Breen L, Philp A, Witard OC, Jackman SR, Selby A, Smith K, Baar K, Tipton KD.** The influence of carbohydrate-protein co-ingestion following endurance exercise on myofibrillar and mitochondrial protein synthesis. *J Physiol.* 2011 Aug 15;589(Pt 16):4011–25.
- Buckley JD, Thomson RL, Coates AM, Howe PR, DeNichilo MO, Rowney MK.** Supplementation with a whey protein hydrolysate enhances recovery of muscle force-generating capacity following eccentric exercise. *J Sci Med Sport.* 2010; 13: 178–81.
- Bueno NB, de Melo IS, de Oliveira SL, da Rocha Ataide T.** Very-low-carbohydrate ketogenic diet v. low-fat diet for long-term weight loss: a meta-analysis of randomised controlled trials. *Br J Nutr.* 2013 Oct;110(7):1178–87.
- Burd NA, Yang Y, Moore DR, Tang JE, Tarnopolsky MA, Phillips SM.** Greater stimulation of myofibrillar protein synthesis with ingestion of whey protein isolate v. micellar casein at rest and after resistance exercise in elderly men. *Br J Nutr.* 2012; 108: 958–62.
- Burke LM, Slater G, Broad EM, Haukka J, Modulon S, Hopkins WG.** Eating patterns and meal frequency of elite Australian athletes. *Int J Sport Nutr Exerc Metab.* 2003 Dec;13(4):521–38.
- Calloway DH, Spector H.** Nitrogen balance as related to caloric and protein intake in active young men. *Am J Clin Nutr.* 1954 Nov-Dec;2(6):405–12
- Carbone JW, McClung JP, Pasiakos SM.** Skeletal muscle responses to negative energy balance: effects of dietary protein. *Adv Nutr.* 2012 Mar 1;3(2):119–26.

Cassady BA, Considine RV, Mattes RD. Beverage consumption, appetite, and energy intake: what did you expect? Am J Clin Nutr. 2012 Mar;95(3):587–93

Cerbulis J, Farrell HM Jr. Composition of milks of dairy cattle. I. Protein, lactose, and fat contents and distribution of protein fraction. J Dairy Sci. 1975 Jun;58(6):817–27.

Cermak NM, Res PT, de Groot LC, Saris WH, van Loon LJ. Protein supplementation augments the adaptive response of skeletal muscle to resistance-type exercise training: a meta-analysis. Am J Clin Nutr. 2012 Dec;96(6):1454–64.

Churchward-Venne TA, Burd NA, Phillips SM. Nutritional regulation of muscle protein synthesis with resistance exercise: strategies to enhance anabolism. Nutr Metab (Lond). 2012; 9: 40.

Churchward-Venne TA, Murphy CH, Longland TM, Phillips SM. Role of protein and amino acids in promoting lean mass accretion with resistance exercise and attenuating lean mass loss during energy deficit in humans. Amino Acids. 2013 Aug;45(2):231–40.

Clarys P, Deliens T, Huybrechts I, Deriemaeker P, Vanaelst B, De Keyzer W, Hebbelinck M, Mullie P. Comparison of nutritional quality of the vegan, vegetarian, semi-vegetarian, pesco-vegetarian and omnivorous diet. Nutrients. 2014 Mar 24;6(3):1318–32.

Clifton PM, Condo D, Keogh JB. Long term weight maintenance after advice to consume low carbohydrate, higher protein diets – A systematic review and meta analysis. Nutr Metab Cardiovasc Dis. 2014 Mar;24(3):224–35

Cockburn E, Stevenson E, Hayes PR, Robson-Ansley P, Howatson G. Effect of milk-based carbohydrate-protein supplement timing on the attenuation of exercise-induced muscle damage. Appl Physiol Nutr Metab. 2010; 35: 270–7.

Cockburn E, Robson-Ansley P, Hayes PR, Stevenson E. Effect of volume of milk consumed on the attenuation of exercise-induced muscle damage. Eur J Appl Physiol. 2012 Sep;112(9):3187–94

Coffey VG, Moore DR, Burd NA, Rerecich T, Stellingwerff T, Garnham AP, Phillips SM, Hawley JA. Nutrient provision increases signalling and protein synthesis in human skeletal muscle after repeated sprints. Eur J Appl Physiol. 2011 Jul;111(7):1473–83

Dattilo M, Antunes HK, Medeiros A, Mônico Neto M, Souza HS, Tufik S, de Mello MT. Sleep and muscle recovery: endocrinological and molecular basis for a new and promising hypothesis. Med Hypotheses. 2011 Aug;77(2):220–2.

de Souza RJ, Bray GA, Carey VJ, Hall KD, LeBoff MS, Loria CM, Laranjo NM, Sacks FM, Smith SR. Effects of 4 weight-loss diets differing in fat, protein, and carbohydrate on fat mass, lean mass, visceral adipose tissue, and hepatic fat: results from the POUNDS LOST trial. Am J Clin Nutr. 2012 Mar;95(3):614–25.

Dideriksen KJ, Reitelseder S, Petersen SG, Hjort M, Helmark IC, Kjaer M, Holm L. Stimulation of muscle protein synthesis by whey and caseinate ingestion after resistance exercise in elderly individuals. Scand J Med Sci Sports. 2011; 21: 372–83.

Dideriksen K, Reitelseder S, Holm L. Influence of amino acids, dietary protein, and physical activity on muscle mass development in humans. Nutrients. 2013 Mar 13;5(3):852–76.

Dreyer HC, Fujita S, Cadenas JG, Chinkes DL, Volpi E, Rasmussen BB. Resistance exercise increases AMPK activity and reduces 4E-BP1 phosphorylation and protein synthesis in human skeletal muscle. J Physiol. 2006 Oct 15;576(Pt 2):613–24. Epub 2006 Jul 27.

European Food Safety Authority. Scientific Opinion on the substantiation of health claims related to protein and increase in satiety leading to a reduction in energy intake (ID 414, 616, 730), contribution to the maintenance or achievement of a normal body weight (ID 414, 616, 730), maintenance of normal bone (ID 416) and growth or maintenance of muscle mass (ID 415, 417, 593, 594, 595, 715) pursuant to Article 13(1) of Regulation (EC) No 1924/2006. EFSA Journal 2010; 8: 1811.

Egli L, Lecoultrre V, Theytaz F, Campos V, Hodson L, Schneiter P, Mittendorfer B, Patterson BW, Fielding BA, Gerber PA, Giusti V, Berneis K, Tappy L. Exercise prevents fructose-induced hypertriglyceridemia in healthy young subjects. Diabetes. 2013 Jul;62(7):2259–65

Elliot TA, Cree MG, Sanford AP, Wolfe RR, Tipton KD. Milk ingestion stimulates net muscle protein synthesis following resistance exercise. Med Sci Sports Exerc. 2006; 38: 667–74.

Ericson U, Sonestedt E, Gullberg B, Hellstrand S, Hindy G, Wirkkula E, Orho-Melander M. High intakes of protein and processed meat associate with increased incidence of type 2 diabetes. *Br J Nutr.* 2013 Mar 28;109(6):1143–53.

Faber M, Benade AJ, van Eck M. Dietary intake, anthropometric measurements, and blood lipid values in weight training athletes (body builders). *Int J Sports Med.* 1986 Dec;7(6):342–6.

Feinman RD, Fine EJ. Nonequilibrium thermodynamics and energy efficiency in weight loss diets. *Theor Biol Med Model.* 2007 Jul 30;4:27.

Ferguson-Stegall L, McCleave E, Ding Z, Doerner Iii PG, Liu Y, Wang B, Healy M, Kleinert M, Dessard B, ym. Aerobic exercise training adaptations are increased by postexercise carbohydrate-protein supplementation. *J Nutr Metab.* 2011;2011:623182.

Figueiredo VC, Cameron-Smith D. Is carbohydrate needed to further stimulate muscle protein synthesis/hypertrophy following resistance exercise? *J Int Soc Sports Nutr.* 2013 Sep 25;10(1):42.

Fogelholm M. Suomalaiset ravitsemussuositukset 2014. Mitä, miksi ja miten? Ruokapohjaiset suositukset. Ravitsemussuositusten julkaisutilaisuus Helsinki 23.1.2014.

Fogelholm M, Anderssen S, Gunnarsdottir I, Lahti-Koski M. Dietary macronutrients and food consumption as determinants of long-term weight change in adult populations: a systematic literature review. *Food Nutr Res.* 2012;56.

Forbes GB, Brown MR, Welle SL, Lipinski BA. Deliberate overfeeding in women and men: energy cost and composition of the weight gain. *Br J Nutr.* 1986; 56: 1–9.

Fouillet H, Gaudichon C, Mariotti F, Bos C, Huneau JF, Tomé D. Energy nutrients modulate the splanchnic sequestration of dietary nitrogen in humans: a compartmental analysis. *Am J Physiol Endocrinol Metab.* 2001; 281: 248–60.

Fung TT, van Dam RM, Hankinson SE, Stampfer M, Willett WC, Hu FB. Low-carbohydrate diets and all-cause and cause-specific mortality: two cohort studies. *Ann Intern Med.* 2010 Sep 7;153(5):289–98.

Garthe I, Raastad T, Refsnes PE, Koivisto A, Sundgot-Borgen J. Effect of two different weight-loss rates on body composition and strength and power-related performance in elite athletes. *Int J Sport Nutr Exerc Metab.* 2011 Apr;21(2):97–104.

Garthe I, Raastad T, Refsnes PE, Sundgot-Borgen J. Effect of nutritional intervention on body composition and performance in elite athletes. *Eur J Sport Sci.* 2013;13(3):295–303.

Gibala M. Molecular responses to high-intensity interval exercise. *Appl Physiol Nutr Metab.* 2009 Jun;34(3):428–32. doi: 10.1139/H09-046. Review.

Gilani GS1, Cockell KA, Sepehr E. Effects of antinutritional factors on protein digestibility and amino acid availability in foods. *J AOAC Int.* 2005 May-Jun;88(3):967–87.

Gilani GS, Xiao C, Lee N. Need for accurate and standardized determination of amino acids and bioactive peptides for evaluating protein quality and potential health effects of foods and dietary supplements. *J AOAC Int.* 2008 Jul-Aug;91(4):894–900. Review.

Gow ML, Ho M, Burrows TL, Baur LA, Stewart L, Hutchesson MJ, Cowell CT, Collins CE, Garnett SP. Impact of dietary macronutrient distribution on BMI and cardiometabolic outcomes in overweight and obese children and adolescents: a systematic review. *Nutr Rev.* 2014 Jun 11.

Haakonssen EC, Martin DT, Burke LM, Jenkins DG. Increased lean mass with reduced fat mass in an elite female cyclist returning to competition: case study. *Int J Sports Physiol Perform.* 2013 Nov;8(6):699–701.

Haakonssen EC, Ross ML, Cato LE, Nana A, Knight EJ, Jenkins DG, Martin DT, Burke LM. Dairy Based Pre-Exercise Meal Does Not Affect Gut Comfort or Time-Trial Performance in Female Cyclists. *Int J Sport Nutr Exerc Metab.* 2014 Jul 14.

Halson SL. Sleep in elite athletes and nutritional interventions to enhance sleep. *Sports Med.* 2014 May;44 Suppl 1:13–23.

Halton TL, Hu FB. The effects of high protein diets on thermogenesis, satiety and weight loss: a critical review. *J Am Coll Nutr.* 2004 Oct;23(5):373–85.

Hartman JW, Tang JE, Wilkinson SB, Tarnopolsky MA, Lawrence RL, Fullerton AV, Phillips SM. Consumption of fat-free fluid milk after resistance exercise promotes greater lean mass accretion than does consumption of soy or carbohydrate in young, novice, male weightlifters. *Am J Clin Nutr.* 2007 Aug;86(2):373–81.

Heaney RP, Layman DK. Amount and type of protein influences bone health. *Am J Clin Nutr.* 2008 May;87(5):1567S–1570S.

Helms ER, Zinn C, Rowlands DS, Brown SR. A systematic review of dietary protein during caloric restriction in resistance trained lean athletes: a case for higher intakes. *Int J Sport Nutr Exerc Metab.* 2014 Apr;24(2):127–38.

Hoffman JR, Falvo MJ. Protein – Which is Best? *J Sports Sci Med.* 2004 Sep 1;3(3):118–130.

House JD, Neufeld J, Leson G. Evaluating the quality of protein from hemp seed (*Cannabis sativa L.*) products through the use of the protein digestibility-corrected amino acid score method. *J Agric Food Chem.* 2010 Nov 24;58(22):11801–7.

Howatson G, Hoad M, Goodall S, Tallent J, Bell PG, French DN. Exercise-induced muscle damage is reduced in resistance-trained males by branched chain amino acids: a randomized, double-blind, placebo controlled study. *J Int Soc Sports Nutr.* 2012 May 8;9(1):20.

Howarth KR, Moreau NA, Phillips SM, Gibala MJ. Coingestion of protein with carbohydrate during recovery from endurance exercise stimulates skeletal muscle protein synthesis in humans. *J Appl Physiol (1985).* 2009 Apr;106(4):1394–402

Hughes GJ, Ryan DJ, Mukherjea R, Schasteen CS. Protein digestibility-corrected amino acid scores (PDCAAS) for soy protein isolates and concentrate: criteria for evaluation. *J Agric Food Chem.* 2011 Dec 14;59(23):12707–12.

Hulmi JJ, Volek JS, Selänne H, Mero AA. Protein ingestion prior to strength exercise affects blood hormones and metabolism. *Med Sci Sports Exerc.* 2005; 37: 1990–7.

Hulmi JJ, Lockwood CM, Stout JR. Effect of protein/essential amino acids and resistance training on skeletal muscle hypertrophy: A case for whey protein. *Nutr Metab (Lond).* 2010; 7: 51.

IOC consensus statement on sports nutrition 2010. *J Sports Sci.* 2011;29 Suppl 1:S3–4

Institute of Medicine (2005) Dietary Reference Intakes for energy, carbohydrate, fiber, fat, fatty acids, cholesterol, protein, and amino acids. Washington, DC: National Academies Press.

Jabekk PT, Moe IA, Meen HD, Tomten SE, Høstmark AT

Jia Y, Hwang SY, House JD, Ogborn MR, Weiler HA, et al. (2010) Long-term high intake of whole proteins results in renal damage in pigs. *J Nutr* 140: 1646–1652

Josse AR, Phillips SM. Impact of milk consumption and resistance training on body composition of female athletes. *Med Sport Sci.* 2012;59:94–103.

Katan MB. Weight-loss diets for the prevention and treatment of obesity. *N Engl J Med.* 2009 Feb 26;360(9):923–5.

Kerksick C, Harvey T, Stout J, Campbell B, Wilborn C, Kreider R, Kalman D, Ziegenfuss T, Lopez H, Landis J, Ivy JL, Antonio J. International Society of Sports Nutrition position stand: nutrient timing. *J Int Soc Sports Nutr.* 2008; 5: 17.

Kim H, Lee S, Choue R. Metabolic responses to high protein diet in Korean elite bodybuilders with high-intensity resistance exercise. *J Int Soc Sports Nutr.* 2011 Jul 4;8:10.)

Kleiner SM, Bazzarre TL, Litchford MD. Metabolic profiles, diet, and health practices of championship male and female bodybuilders. *J Am Diet Assoc.* 1990 Jul;90(7):962–7.

Kraemer WJ, Solomon-Hill G, Volk BM, Kupchak BR, Looney DP, Dunn-Lewis C, Comstock BA ym. The effects of soy and whey protein supplementation on acute hormonal responses to resistance exercise in men. *J Am Coll Nutr.* 2013;32(1):66–74.

Kreider RB, Klesges R, Harmon K, Grindstaff P, Ramsey L, Bullen D, Wood L, Li Y, Almada A. Effects of ingesting supplements designed to promote lean tissue accretion on body composition during resistance training. *Int J Sport Nutr.* 1996; 6: 234–46.

Kreider RB, Wilborn CD, Taylor L, Campbell B, Almada AL, Collins R, Cooke M, Earnest CP ym. ISSN exercise & sport nutrition review: research & recommendations. *J Int Soc Sports Nutr.* 2010; 7: 7.

Kulik JR, Touchberry CD, Kawamori N, Blumert PA, Crum AJ, Haff GG. Supplemental carbohydrate ingestion does not improve performance of high-intensity resistance exercise. *J Strength Cond Res.* 2008 Jul;22(4):1101–7.

Kumar V, Atherton P, Smith K, Rennie MJ. Human muscle protein synthesis and breakdown during and after exercise. *J Appl Physiol.* 2009;106:2026–2039.

Lacroix M, Bos C, Léonil J, Airinei G, Luengo C, Daré S, Benamouzig R, Fouillet H, Fauquant J, Tomé D ym. Compared with casein or total milk protein, digestion of milk soluble proteins is too rapid to sustain the anabolic postprandial amino acid requirement. *Am J Clin Nutr.* 2006 Nov;84(5):1070–9.

Lambert CP, Frank LL, Evans WJ. Macronutrient considerations for the sport of bodybuilding. *Sports Med.* 2004;34(5):317–27.

Lemon PW, Berardi JM, Noreen EE. The role of protein and amino acid supplements in the athlete's diet: does type or timing of ingestion matter? *Curr Sports Med Rep.* 2002 Aug;1(4):214–21.

Lepe M, Bacardí Gascón M, Jiménez Cruz A. Long-term efficacy of high-protein diets: a systematic review. *Nutr Hosp.* 2011 Nov-Dec;26(6):1256–9.

Lis D, Stellingwerff T, Shing CM, Ahuja K DK, Fell J. Exploring the Popularity, Experiences and Beliefs Surrounding Gluten-Free Diets in Non-Coeliac Athletes. *Int J Sport Nutr Exerc Metab.* 2014 Jun 5.

Macdermid PW, Stannard SR. A whey-supplemented, high-protein diet versus a high-carbohydrate diet: effects on endurance cycling performance. *Int J Sport Nutr Exerc Metab.* 2006 Feb;16(1):65–77.

Mamerow MM, Mettler JA, English KL, Casperson SL, Arentson-Lantz E, Sheffield-Moore M, Layman DK ym. Dietary protein distribution positively influences 24-h muscle protein synthesis in healthy adults. *J Nutr.* 2014 Jun;144(6):876–80.

Manore MM. Weight management in the performance athlete. *Nestle Nutr Inst Workshop Ser.* 2013;75:123–33.

McLellan TM, Pasiakos SM, Lieberman HR. Effects of protein in combination with carbohydrate supplements on acute or repeat endurance exercise performance: a systematic review. *Sports Med.* 2014 Apr;44(4):535–50

McPherson RA, Hardy G. Clinical and nutritional benefits of cysteine-enriched protein supplements. *Curr Opin Clin Nutr Metab Care.* 2011 Nov;14(6):562–8.

Metzler S, Mitchell N, Tipton KD. Increased protein intake reduces lean body mass loss during weight loss in athletes. *Med Sci Sports Exerc.* 2010 Feb;42(2):326–37

Miller PE, Alexander DD, Perez V. Effects of whey protein and resistance exercise on body composition: a meta-analysis of randomized controlled trials. *J Am Coll Nutr.* 2014 Apr;33(2):163–75.

Milsom J, Barreria P, Burgess DJ, Iqbal Z, Morton JP. Case-Study: Muscle Atrophy and Hypertrophy in a Premier League Soccer Player During Rehabilitation From ACL Injury. *Int J Sport Nutr Exerc Metab.* 2014 Jan 17

Moore DR, Robinson MJ, Fry JL, Tang JE, Glover EI, Wilkinson SB, Prior T, Tarnopolsky MA, Phillips SM. Ingested protein dose response of muscle and albumin protein synthesis after resistance exercise in young men. *Am J Clin Nutr.* 2009; 89: 161–8.

Moore DR, Areta J, Coffey VG, Stellingwerff T, Phillips SM, Burke LM, Cléroux M, Godin JP, Hawley JA. Daytime pattern of post-exercise protein intake affects whole-body protein turnover in resistance-trained males. *Nutr Metab (Lond).* 2012 Oct 16;9(1):91.

Mullins NM, Sinning WE. Effects of resistance training and protein supplementation on bone turnover in young adult women. *Nutr Metab (Lond).* 2005 Aug 17;2:19.

Navas-Carretero S, Abete I, Zuleta MA, Martínez JA. Chronologically scheduled snacking with high-protein products within the habitual diet in type-2 diabetes patients leads to a fat mass loss: a longitudinal study. *Nutr J.* 2011 Jul 14;10:74

Noto H, Goto A, Tsujimoto T, Noda M. Low-carbohydrate diets and all-cause mortality: a systematic review and meta-analysis of observational studies. *PLoS One.* 2013;8(1):e55030.

Pan A, Hu FB. Effects of carbohydrates on satiety: differences between liquid and solid food. *Curr Opin Clin Nutr Metab Care.* 2011 Jul;14(4):385–90.

Paoli A. Ketogenic diet for obesity: friend or foe? *Int J Environ Res Public Health.* 2014 Feb 19;11(2):2092–107.

Paoli A, Grimaldi K, D'Agostino D, Cenci L, Moro T, Bianco A, Palma A. Ketogenic diet does not affect strength performance in elite artistic gymnasts. *J Int Soc Sports Nutr.* 2012 Jul;9(1):34.

Paoli A, Rubini A, Volek JS, Grimaldi KA. Beyond weight loss: a review of the therapeutic uses of very-low-carbohydrate (ketogenic) diets. *Eur J Clin Nutr.* 2013 Aug;67(8):789–96.

Paddon-Jones D, Westman E, Mattes RD, Wolfe RR, Astrup A, Westerterp-Plantenga M. Protein, weight management, and satiety. *Am J Clin Nutr.* 2008 May;87(5):1558S-1561S.

Pasiakos SM, Lieberman HR, McLellan TM. Effects of Protein Supplements on Muscle Damage, Soreness and Recovery of Muscle Function and Physical Performance: A Systematic Review. *Sports Med.* 2014 Jan 17

Pedersen AN, Kondrup J, Børshøj E. Health effects of protein intake in healthy adults: a systematic literature review. *Food Nutr Res.* 2013 Jul 30;57.

Pennings B, Groen B, de Lange A, Gijsen AP, Zorenc AH, Senden JM, van Loon LJ. Amino acid absorption and subsequent muscle protein accretion following graded intakes of whey protein in elderly men. *Am J Physiol Endocrinol Metab.* 2012; 302: 992–9.

Phillips SM. Muscle growth, adaption and repair. IOC Consensus In Sports Nutrition Conference Lausanne, Switzerland 25th–27th October 2010

Phillips SM, Tipton KD, Aarsland A, Wolf SE, Wolfe RR. Mixed muscle protein synthesis and breakdown after resistance exercise in humans. *Am J Physiol.* 1997;273:E99–E107

Phillips SM, Moore DR, Tang JE. A critical examination of dietary protein requirements, benefits, and excesses in athletes. *Int J Sport Nutr Exerc Metab.* 2007 Aug;17 Suppl:S58–76.

Phillips SM, Van Loon LJ. Dietary protein for athletes: from requirements to optimum adaptation. *J Sports Sci.* 2011;29 Suppl 1:S29–38.

Poortmans JR, Dellalieux O. Do regular high protein diets have potential health risks on kidney function in athletes? *Int J Sport Nutr Exerc Metab.* 2000 Mar;10(1):28–38.

Poortmans JR, Carpentier A, Pereira -Lancha LO, Lancha A Jr. Protein turnover, amino acid requirements and recommendations for athletes and active populations. *Braz J Med Biol Res.* 2012 Oct;45(10):875–90.

Power O, Hallihan A, Jakeman P. Human insulinotropic response to oral ingestion of native and hydrolysed whey protein. *Amino Acids.* 2009; 37: 333–9.

Pritchett K, Pritchett R. Chocolate milk: a post-exercise recovery beverage for endurance sports. *Med Sport Sci.* 2012;59:127–34

Raben A, Agerholm-Larsen L, Flint A, Holst JJ, Astrup A. Meals with similar energy densities but rich in protein, fat, carbohydrate, or alcohol have different effects on energy expenditure and substrate metabolism but not on appetite and energy intake. *Am J Clin Nutr.* 2003; 77: 91–100.

Reidy PT, Walker DK, Dickinson JM, Gundermann DM, Drummond MJ, Timmerman KL, Fry CS ym. Protein blend ingestion following resistance exercise promotes human muscle protein synthesis. *J Nutr.* 2013 Apr;143(4):410–6.

Reitelseder S, Agergaard J, Doessing S, Helmark IC, Lund P, Kristensen NB, Frystyk J, Flyvbjerg A ym. Whey and casein labeled with L-[1–13C]leucine and muscle protein synthesis: effect of resistance exercise and protein ingestion. *Am J Physiol Endocrinol Metab.* 2011; 300: 231–42.

Res PT, Groen B, Pennings B, Beelen M, Wallis GA, Gijsen AP, Senden JM, VAN Loon LJ. Protein ingestion before sleep improves postexercise overnight recovery. *Med Sci Sports Exerc.* 2012 Aug;44(8):1560–9.

Robinson MJ1, Burd NA, Breen L, Rerecich T, Yang Y, Hector AJ, Baker SK, Phillips SM. Dose-dependent responses of myofibrillar protein synthesis with beef ingestion are enhanced with resistance exercise in middle-aged men. *Appl Physiol Nutr Metab.* 2013 Feb;38(2):120–5.

Sacks FM, Bray GA, Carey VJ, Smith SR, Ryan DH, Anton SD, McManus K, Champagne CM, Bishop LM ym. Comparison of weight-loss diets with different compositions of fat, protein, and carbohydrates. *N Engl J Med.* 2009 Feb 26;360(9):859–73.

Sallinen J, Pakarinen A, Ahtiainen J, Kraemer WJ, Volek JS, Hakkinen K. Relationship between diet and serum anabolic hormone responses to heavy-resistance exercise in men. *Int J Sports Med.* 2004 Nov;25(8):627–33.

Santesso N, Akl EA, Bianchi M, Mente A, Mustafa R, Heels-Ansdell D, Schünemann HJ. Effects of higher-versus lower-protein diets on health outcomes: a systematic review and meta-analysis. *Eur J Clin Nutr.* 2012 Jul;66(7):780–8

Sarwar G, McDonough FE. Evaluation of protein digestibility-corrected amino acid score method for assessing protein quality of foods. *J Assoc Off Anal Chem.* 1990 May-Jun;73(3):347–56.

Sawyer JC, Wood RJ, Davidson PW, Collins SM, Matthews TD, Gregory SM, Paolone VJ. Effects of a short-term carbohydrate-restricted diet on strength and power performance. *J Strength Cond Res.* 2013 Aug;27(8):2255–62.

Schaafsma G. The protein digestibility-corrected amino acid score. *J Nutr.* 2000 Jul;130(7):1865S–7S.

Schaafsma G. Advantages and limitations of the protein digestibility-corrected amino acid score (PDCAAS) as a method for evaluating protein quality in human diets. *Br J Nutr.* 2012 Aug;108 Suppl 2:S333–6.

Schoenfeld BJ, Aragon AA, Krieger JW. The effect of protein timing on muscle strength and hypertrophy: a meta-analysis. *J Int Soc Sports Nutr.* 2013 Dec 3;10(1):53.

Schwingshackl L, Hoffmann G. Long-term effects of low-fat diets either low or high in protein on cardiovascular and metabolic risk factors: a systematic review and meta-analysis. *Nutr J.* 2013 Apr 15;12:48. doi: 10.1186/1475–2891–12–48. Review.

Schwingshackl L, Hoffmann G. Comparison of high vs. normal/low protein diets on renal function in subjects without chronic kidney disease: a systematic review and meta-analysis. *PLoS One.* 2014 May 22;9(5):e97656.

Seene T, Kaasik P, Alev K. Muscle protein turnover in endurance training: a review. *Int J Sports Med.* 2011 Dec;32(12):905–11.

Shimomura Y, Inaguma A, Watanabe S, Yamamoto Y, Muramatsu Y, Bajotto G, Sato J, Shimomura N ym. Branched-chain amino acid supplementation before squat exercise and delayed-onset muscle soreness. *Int J Sport Nutr Exerc Metab.* 2010; 20: 236–44.

Slater G. Nutrition for strength sports. IOC Consensus In Sports Nutrition Conference Lausanne, Switzerland 25th–27th October 2010

Slater G, Phillips SM. Nutrition guidelines for strength sports: sprinting, weightlifting, throwing events, and bodybuilding. *J Sports Sci.* 2011;29 Suppl 1:S67–77.

Soenen S, Bonomi AG, Lemmens SG, Scholte J, Thijssen MA, van Berkum F, Westerterp-Plantenga MS. Relatively high-protein or ‘low-carb’ energy-restricted diets for body weight loss and body weight maintenance? *Physiol Behav.* 2012 Oct 10;107(3):374–80.

Soop M, Nehra V, Henderson GC, Boirie Y, Ford GC, Nair KS. Coingestion of whey protein and casein in a mixed meal: demonstration of a more sustained anabolic effect of casein. *Am J Physiol Endocrinol Metab.* 2012 Jul 1;303(1):E152–62.

Stark M, Lukaszuk J, Prawitz A, Salacinski A. Protein timing and its effects on muscular hypertrophy and strength in individuals engaged in weight-training. *J Int Soc Sports Nutr.* 2012 Dec 14;9(1):54

Stearns RL, Emmanuel H, Volek JS, Casa DJ. Effects of ingesting protein in combination with carbohydrate during exercise on endurance performance: a systematic review with meta-analysis. *J Strength Cond Res.* 2010 Aug;24(8):2192–202.

Stellingwerff T. Nutrition for power sports. IOC Consensus In Sports Nutrition Conference Lausanne, Switzerland 25th–27th October 2010

Stull AJ, Apolzan JW, Thalacker-Mercer AE, Iglay HB, Campbell WW. Liquid and solid meal replacement products differentially affect postprandial appetite and food intake in older adults. *J Am Diet Assoc.* 2008 Jul;108(7):1226–30

Suárez López MM, Kizlansky A, López LB. [Assessment of protein quality in foods by calculating the amino acids score corrected by digestibility]. *Nutr Hosp.* 2006 Jan-Feb;21(1):47–51.

Schwingshackl L, Hoffmann G. Long-term effects of low-fat diets either low or high in protein on cardiovascular and metabolic risk factors: a systematic review and meta-analysis. *Nutr J.* 2013 Apr 15;12:48.

Tang JE, Moore DR, Kujbida GW, Tarnopolsky MA, Phillips SM. Ingestion of whey hydrolysate, casein, or soy protein isolate: effects on mixed muscle protein synthesis at rest and following resistance exercise in young men. *J Appl Physiol.* 2009; 107: 987–92.

Tang M, O'Connor LE, Campbell WW. Diet-induced weight loss: the effect of dietary protein on bone. *J Acad Nutr Diet.* 2014 Jan;114(1):72–85.

Tarnopolsky MA, Atkinson SA, MacDougall JD, Senor BB, Lemon PW, Schwarcz H. Whole body leucine metabolism during and after resistance exercise in fed humans. *Med Sci Sports Exerc.* 1991 Mar;23(3):326–33.

Tarnopolsky MA, Atkinson SA, MacDougall JD, Chesley A, Phillips S, Schwarcz HP. Evaluation of protein requirements for trained strength athletes. *J Appl Physiol.* 1992 Nov;73(5):1986–95.

Tipton KD, Rasmussen BB, Miller SL, Wolf SE, Owens-Stovall SK, Petrucci BE, Wolfe RR. Timing of amino acid-carbohydrate ingestion alters anabolic response of muscle to resistance exercise. *Am J Physiol Endocrinol Metab.* 2001 Aug;281(2):E197–206.

Tipton KD, Elliott TA, Cree MG, Aarsland AA, Sanford AP, Wolfe RR. Stimulation of net muscle protein synthesis by whey protein ingestion before and after exercise. *Am J Physiol Endocrinol Metab.* 2007; 292: 71–6.

Tipton KD. Efficacy and consequences of very-high-protein diets for athletes and exercisers. *Proc Nutr Soc.* 2011 May;70(2):205–14.

Tipton KD. Dietary strategies to attenuate muscle loss during recovery from injury. *Nestle Nutr Inst Workshop Ser.* 2013;75:51–61

Vandenbogaerde TJ, Hopkins WG. Effects of acute carbohydrate supplementation on endurance performance: a meta-analysis. *Sports Med.* 2011 Sep 1;41(9):773–92.

van Loon LJ, Kruijshoop M, Verhagen H, Saris WH, Wagenmakers AJ. Ingestion of protein hydrolysate and amino acid-carbohydrate mixtures increases postexercise plasma insulin responses in men. *J Nutr.* 2000 Oct;130(10):2508–13.

van Loon LJ, Gibala MJ. Dietary protein to support muscle hypertrophy. *Nestle Nutr Inst Workshop Ser.* 2011;69:79–89; discussion 89–95

van Wijck K1, Pennings B, van Bijnen AA, Senden JM, Buurman WA, Dejong CH, van Loon LJ, Lenaerts K. Dietary protein digestion and absorption are impaired during acute postexercise recovery in young men. *Am J Physiol Regul Integr Comp Physiol.* 2013 Mar 1;304(5):R356–61

Veldhorst M, Smeets A, Soenen S, Hochstenbach-Waelen A, Hursel R, Diepvens K, Lejeune M, Luscombe-Marsh N ym. Protein-induced satiety: effects and mechanisms of different proteins. *Physiol Behav.* 2008 May 23;94(2):300–7.

Volek JS, Kraemer WJ, Bush JA, Incledon T, Boetes M. Testosterone and cortisol in relationship to dietary nutrients and resistance exercise. *J Appl Physiol.* 1997 Jan;82(1):49–54

Volek JS, Volk BM, Gómez AL, Kunce LJ, Kupchak BR, Freidenreich DJ, Aristizabal JC, Saenz C, Dunn-Lewis C, Ballard KD ym. Whey protein supplementation during resistance training augments lean body mass. *J Am Coll Nutr.* 2013;32(2):122–35.

VRN. Terveyttä ruoasta – Suomalaiset ravitsemussuositukset 2014. Juvenes Oy, Helsinki 2014.

Wakefield AP, House JD, Ogbon MR, Weiler HA, Aukema HM. A diet with 35% of energy from protein leads to kidney damage in female Sprague-Dawley rats. *Br J Nutr.* 2011 Sep;106(5):656–63.

Welle S, Matthews DE, Campbell RG, Nair KS. Stimulation of protein turnover by carbohydrate overfeeding in men. *Am J Physiol.* 1989; 257: 413–7.

West DW, Burd NA, Coffey VG, Baker SK, Burke LM, Hawley JA, Moore DR, Stellingwerff T, Phillips SM. Rapid aminoacidemia enhances myofibrillar protein synthesis and anabolic intramuscular signaling responses after resistance exercise. *Am J Clin Nutr.* 2011; 94: 795–803.

Westerterp-Plantenga MS, Nieuwenhuizen A, Tomé D, Soenen S, Westerterp KR. Dietary protein, weight loss, and weight maintenance. *Annu Rev Nutr.* 2009;29:21–41.

Westerterp-Plantenga MS, Lemmens SG, Westerterp KR. Dietary protein – its role in satiety, energetics, weight loss and health. *Br J Nutr.* 2012 Aug;108 Suppl 2:S105–12.

Whitney EN, Rolfes SR. Protein: Amino Acids. Kirjassa: Whitney EN, Rolfes SR. Understanding Nutrition 7th ed. West publishing company, USA, 1996.

WHO Technical Report Series 935. Protein and Amino Acid Requirements in Human Nutrition: report of a joint FAO/WHO/UNU expert consultation. Report of a Joint WHO/FAO/UNU Expert Consultation. 2011.

Wilkinson SB, Tarnopolsky MA, MacDonald MJ, Macdonald JR, Armstrong D, Phillips SM. Consumption of fluid skim milk promotes greater muscle protein accretion following resistance exercise than an isonitrogenous and isoenergetic soy protein beverage. *Am J Clin Nutr.* 2007; 85: 1031–1040

Wilson JM, Marin PJ, Rhea MR, Wilson SM, Loenneke JP, Anderson JC. Concurrent training: a meta-analysis examining interference of aerobic and resistance exercises. *J Strength Cond Res.* 2012 Aug;26(8):2293–307

Witard OC, Jackman SR, Kies AK, Jeukendrup AE, Tipton KD. Effect of increased dietary protein on tolerance to intensified training. *Med Sci Sports Exerc.* 2011 Apr;43(4):598–607

Witard OC, Turner JE, Jackman SR, Kies AK, Jeukendrup AE, Bosch JA, Tipton KD. High dietary protein restores overreaching induced impairments in leukocyte trafficking and reduces the incidence of upper respiratory tract infection in elite cyclists. *Brain Behav Immun.* 2014 Jul;39:211–9. (a)

Witard OC, Jackman SR, Breen L, Smith K, Selby A, Tipton KD. Myofibrillar muscle protein synthesis rates subsequent to a meal in response to increasing doses of whey protein at rest and after resistance exercise. *Am J Clin Nutr.* 2014 Jan;99(1):86–95.

Wycherley TP, Moran LJ, Clifton PM, Noakes M, Brinkworth GD. Effects of energy-restricted high-protein, low-fat compared with standard-protein, low-fat diets: a meta-analysis of randomized controlled trials. *Am J Clin Nutr.* 2012 Dec;96(6):1281–98.

Yang D, Liu Z, Yang H, Jue Y. Acute effects of high-protein versus normal-protein isocaloric meals on satiety and ghrelin. *Eur J Nutr.* 2014 Mar;53(2):493–500

Zajac A, Poprzecki S, Maszczyk A, Czuba M, Michalczyk M, Zydek G. The effects of a ketogenic diet on exercise metabolism and physical performance in off-road cyclists. *Nutrients.* 2014 Jun 27;6(7):2493–508.

Lähdeluettelo luku 7

Alfenas RC, Mattes RD. Effect of fat sources on satiety. *Obes Res.* 2003 Feb;11(2):183–7.

Artero EG, Lee DC, Lavie CJ, España-Romero V, Sui X, Church TS, Blair SN. Effects of muscular strength on cardiovascular risk factors and prognosis. *J Cardiopulm Rehabil Prev.* 2012 Nov-Dec;32(6):351–8.

Assunção ML, Ferreira HS, dos Santos AF, Cabral CR Jr, Florêncio TM. Effects of dietary coconut oil on the biochemical and anthropometric profiles of women presenting abdominal obesity. *Lipids.* 2009 Jul;44(7):593–601.

Astrup A. The role of dietary fat in obesity. *Semin Vasc Med.* 2005 Feb;5(1):40–7. Review.

Astrup A. Yogurt and dairy product consumption to prevent cardiometabolic diseases: epidemiologic and experimental studies. *Am J Clin Nutr.* 2014 May;99(5 Suppl):1235S-42S.

Baxheinrich A, Stratmann B, Lee-Barkey YH, Tschoepe D, Wahrburg U. Effects of a rapeseed oil-enriched hypoenergetic diet with a high content of -linolenic acid on body weight and cardiovascular risk profile in patients with the metabolic syndrome. *Br J Nutr.* 2012 Aug;108(4):682–91.

Bazzano LA. Effects of soluble dietary fiber on low-density lipoprotein cholesterol and coronary heart disease risk. *Curr Atheroscler Rep.* 2008 Dec;10(6):473–7. Review.

Beermann C, Hartung J. Physiological properties of milk ingredients released by fermentation. *Food Funct.* 2013 Feb;4(2):185–99

Bell EA, Castellanos VH, Pelkman CL, Thorwart ML, Rolls BJ. Energy density of foods affects energy intake in normal-weight women. *Am J Clin Nutr.* 1998 Mar;67(3):412–20.

Bender N, Portmann M, Heg Z, Hofmann K, Zwahlen M, Egger M. Fish or n3-PUFA intake and body composition: a systematic review and meta-analysis. *Obes Rev.* 2014 May 29.

Bendsen NT, Christensen R, Bartels EM, Astrup A. Consumption of industrial and ruminant trans fatty acids and risk of coronary heart disease: a systematic review and meta-analysis of cohort studies. *Eur J Clin Nutr.* 2011 Jul;65(7):773–83.

Bhardwaj S, Passi SJ, Misra A. Overview of trans fatty acids: biochemistry and health effects. *Diabetes Metab Syndr.* 2011 Jul-Sep;5(3):161–4.

Black KE, Skidmore PM, Brown RC. Energy intakes of ultraendurance cyclists during competition, an observational study. *Int J Sport Nutr Exerc Metab.* 2012 Feb;22(1):19–23.

Blundell JE, MacDiarmid JI. Fat as a risk factor for overconsumption: satiation, satiety, and patterns of eating. *J Am Diet Assoc.* 1997 Jul;97(7 Suppl):S63–9. Review.

Blundell JE, Stubbs RJ, Golding C, Croden F, Alam R, Whybrow S, Le Noury J, Lawton CL. Resistance and susceptibility to weight gain: individual variability in response to a high-fat diet. *Physiol Behav.* 2005 Dec 15;86(5):614–22. Epub 2005 Oct 12. Review.

Bourre JM. Roles of unsaturated fatty acids (especially omega-3 fatty acids) in the brain at various ages and during ageing. *J Nutr Health Aging.* 2004;8(3):163–74. Review. (a)

Bourre JM. Dietary omega-3 Fatty acids and psychiatry: mood, behaviour, stress, depression, dementia and aging. *J Nutr Health Aging.* 2005;9(1):31–8. (b)

Bray GA, Paeratakul S, Popkin BM. Dietary fat and obesity: a review of animal, clinical and epidemiological studies. *Physiol Behav.* 2004 Dec 30;83(4):549–55.

Brouns F, Beckers E. Is the gut an athletic organ? Digestion, absorption and exercise. *Sports Med.* 1993 Apr;15(4):242–57.

Brouwer IA, Wanders AJ, Katan MB. Effect of animal and industrial trans fatty acids on HDL and LDL cholesterol levels in humans--a quantitative review. *PLoS One.* 2010 Mar 2;5(3):e9434.

Brouwer IA, Wanders AJ, Katan MB. Trans fatty acids and cardiovascular health: research completed? *Eur J Clin Nutr.* 2013 May;67(5):541–7.

Brown MJ, Ferruzzi MG, Nguyen ML, Cooper DA, Eldridge AL, Schwartz SJ, White WS. Carotenoid bioavailability is higher from salads ingested with full-fat than with fat-reduced salad dressings as measured with electrochemical detection. *Am J Clin Nutr.* 2004 Aug;80(2):396–403.

Buckley JD, Howe PR. Long-chain omega-3 polyunsaturated fatty acids may be beneficial for reducing obesity-a review. *Nutrients.* 2010 Dec;2(12):1212–30.

Burke LM. Energy needs of athletes. *Can J Appl Physiol.* 2001;26 Suppl:S202–19. Review.

Burke LM, Collier GR, Beasley SK, Davis PG, Fricker PA, Heeley P, Walder K, Hargreaves M. Effect of coingestion of fat and protein with carbohydrate feedings on muscle glycogen storage. *J Appl Physiol* (1985). 1995 Jun;78(6):2187–92.

Burke LM, Hawley JA. Effects of short-term fat adaptation on metabolism and performance of prolonged exercise. *Med Sci Sports Exerc.* 2002 Sep;34(9):1492–8.

Burke LM, Hawley JA, Angus DJ, Cox GR, Clark S, Cummings NK, Desbrow B, and Hargreaves M. Adaptations to short-term high-fat diet persist during exercise despite high carbohydrate availability. *Med Sci Sports Exerc* 34: 83–91, 2002.

Burke LM, Kiens B. “Fat adaptation” for athletic performance: the nail in the coffin? *J Appl Physiol* (1985). 2006 Jan;100(1):7–8.

Casas-Agustench P, López-Uriarte P, Bulló M, Ros E, Gómez-Flores A, Salas-Salvadó J. Acute effects of three high-fat meals with different fat saturations on energy expenditure, substrate oxidation and satiety. *Clin Nutr.* 2009 Feb;28(1):39–45*

Chapman CD, Benedict C, Brooks SJ, Schiöth HB. Lifestyle determinants of the drive to eat: a meta-analysis. Am J Clin Nutr. 2012 Sep;96(3):492–7.

Chatterton DE, Nguyen DN, Bering SB, Sangild PT. Anti-inflammatory mechanisms of bioactive milk proteins in the intestine of newborns. Int J Biochem Cell Biol. 2013 Aug;45(8):1730–47

Choi Y, Song S, Song Y, Lee JE. Consumption of red and processed meat and esophageal cancer risk: meta-analysis. World J Gastroenterol. 2013 Feb 21;19(7):1020–9.

Clarke R, Frost C, Collins R, Appleby P, Peto R. Dietary lipids and blood cholesterol: quantitative meta-analysis of metabolic ward studies. BMJ. 1997 Jan 11;314(7074):112–7.

Clegg ME. Medium-chain triglycerides are advantageous in promoting weight loss although not beneficial to exercise performance. Int J Food Sci Nutr. 2010 Nov;61(7):653–79.

Clifton PM. Palm oil and LDL cholesterol. Am J Clin Nutr. 2011 Dec;94(6):1392–3.

Codex Alimentarius Commision. Amended. (2003). Codex Standard for Named Vegetable Oils. Codex Stan 210, p. 13

Collier G, McLean A, O'Dea K. Effect of co-ingestion of fat on the metabolic responses to slowly and rapidly absorbed carbohydrates. Diabetologia. 1984 Jan;26(1):50–4.

Cooney MT, Dudina A, De Bacquer D, Wilhelmsen L, Sans S, Menotti A, De Backer G, Jousilahti P, Keil U ym. HDL cholesterol protects against cardiovascular disease in both genders, at all ages and at all levels of risk. Atherosclerosis. 2009 Oct;206(2):611–6.

Darzi J, Frost GS, Robertson MD. Do SCFA have a role in appetite regulation? Proc Nutr Soc. 2011 Feb;70(1):119–28.

Décombaz J, Grathwohl D, Pollien P, Schmitt JA, Borrani F, Lecoultrre V. Effect of short-duration lipid supplementation on fat oxidation during exercise and cycling performance. Appl Physiol Nutr Metab. 2013 Jul;38(7):766–72

de Oliveira Otto MC, Mozaffarian D, Kromhout D, Bertoni AG, Sibley CT, Jacobs DR Jr, Nettleton JA. Dietary intake of saturated fat by food source and incident cardiovascular disease: the Multi-Ethnic Study of Atherosclerosis. Am J Clin Nutr. 2012 Aug;96(2):397–404.

de Oliveira EP, Burini RC, Jeukendrup A. Gastrointestinal complaints during exercise: prevalence, etiology, and nutritional recommendations. Sports Med. 2014 May;44 Suppl 1:S79–85.

de Souza RJ, Bray GA, Carey VJ, Hall KD, LeBoff MS, Loria CM, Laranjo NM, Sacks FM, Smith SR. Effects of 4 weight-loss diets differing in fat, protein, and carbohydrate on fat mass, lean mass, visceral adipose tissue, and hepatic fat: results from the POUNDS LOST trial. Am J Clin Nutr. 2012 Mar;95(3):614–25.

Donahoo W, Wyatt HR, Kriehn J, Stuht J, Dong F, Hosokawa P, Grunwald GK, Johnson SL, Peters JC, Hill JO. Dietary fat increases energy intake across the range of typical consumption in the United States. Obesity (Silver Spring). 2008 Jan;16(1):64–9.

Donnelly JE, Sullivan DK, Smith BK, Jacobsen DJ, Washburn RA, Johnson SL, Hill JO, Mayo MS ym. Alteration of dietary fat intake to prevent weight gain: Jayhawk Observed Eating Trial. Obesity (Silver Spring). 2008 Jan;16(1):107–12

Dorgan JF, Judd JT, Longcope C, Brown C, Schatzkin A, Clevidence BA, Campbell WS, Nair PP, Franz C ym. Effects of dietary fat and fiber on plasma and urine androgens and estrogens in men: a controlled feeding study. Am J Clin Nutr. 1996 Dec;64(6):850–5.

Durstine JL, Grandjean PW, Cox CA, Thompson PD. Lipids, lipoproteins, and exercise. J Cardiopulm Rehabil. 2002 Nov-Dec;22(6):385–98.

EFSA Panel on Dietetic Products, Nutrition, and Allergies (NDA) Scientific Opinion on Dietary Reference Values for fats, including saturated fatty acids, polyunsaturated fatty acids, monounsaturated fatty acids, trans fatty acids, and cholesterol EFSA Journal 2010;8(3):1461

Elliot TA, Cree MG, Sanford AP, Wolfe RR, Tipton KD. Milk ingestion stimulates net muscle protein synthesis following resistance exercise. Med Sci Sports Exerc. 2006 Apr;38(4):667–74.

Ello-Martin JA, Ledikwe JH, Rolls BJ. The influence of food portion size and energy density on energy intake: implications for weight management. Am J Clin Nutr. 2005 Jul;82(1 Suppl):236S-241S. Review.

European Food Safety Authority. Scientific Opinion on Dietary Reference Values for fats, including saturated fatty acids, polyunsaturated fatty acids, monounsaturated fatty acids, trans fatty acids, and cholesterol. EFSA Journal 2010; 8: 1461.

Fattore E, Bosetti C, Brighenti F, Agostoni C, Fattore G. Palm oil and blood lipid-related markers of cardiovascular disease: a systematic review and meta-analysis of dietary intervention trials. Am J Clin Nutr. 2014 Apr 9;99(6):1331–1350.

Field AE, Willett WC, Lissner L, Colditz GA. Dietary fat and weight gain among women in the Nurses' Health Study. Obesity (Silver Spring). 2007 Apr;15(4):967–76.

Finravinto 2012 -tutkimus. Helldán A, Raulio S, Kosola M, Tapanainen H, Ovaskainen M-L, Virtanen S. THL Raportti 16/2013. Juvenes Print – Suomen Yliopistopaino Oy. Tampere 2013.

Fleming J, Sharman MJ, Avery NG, Love DM, Gómez AL, Scheett TP, Kraemer WJ, Volek JS. Endurance capacity and high-intensity exercise performance responses to a high fat diet. Int J Sport Nutr Exerc Metab. 2003 Dec;13(4):466–78.

Flint A, Helt B, Raben A, Toubro S, Astrup A. Effects of different dietary fat types on postprandial appetite and energy expenditure. Obes Res 2003;11:1449–55.

Flores-Mateo G, Rojas-Rueda D, Basora J, Ros E, Salas-Salvadó J. Nut intake and adiposity: meta-analysis of clinical trials. Am J Clin Nutr. 2013 Jun;97(6):1346–55.

Fogelholm M, Anderssen S, Gunnarsdottir I, Lahti-Koski M. Dietary macronutrients and food consumption as determinants of long-term weight change in adult populations: a systematic literature review. Food Nutr Res. 2012;56.

Forouhi NG, Sharp SJ, Du H, van der A DL, Halkjaer J, Schulze MB, Tjønneland A, Overvad K ym. Dietary fat intake and subsequent weight change in adults: results from the European Prospective Investigation into Cancer and Nutrition cohorts. Am J Clin Nutr. 2009 Dec;90(6):1632–41.

Fouillet H, Gaudichon C, Mariotti F, Bos C, Huneau JF, Tomé D. Energy nutrients modulate the splanchnic sequestration of dietary nitrogen in humans: a compartmental analysis. Am J Physiol Endocrinol Metab. 2001 Aug;281(2):E248–60.

Gardner CD, Fortmann SP, Krauss RM. Association of small low-density lipoprotein particles with the incidence of coronary artery disease in men and women. JAMA. 1996;276:875–81

Gargallo Fernández M, Quiles Izquierdo J, Basulto Marset J ym. FESNAD-SEEDO consensus group. Evidence-based nutritional recommendations for the prevention and treatment of overweight and obesity in adults (FESNAD-SEEDO consensus document). The role of diet in obesity prevention (II/III). Nutr Hosp. 2012 May-Jun;27(3):800–32.

Gebauer SK, Chardigny JM, Jakobsen MU, Lamarche B, Lock AL, Proctor SD, Baer DJ. Effects of ruminant trans fatty acids on cardiovascular disease and cancer: a comprehensive review of epidemiological, clinical, and mechanistic studies. Adv Nutr. 2011 Jul;2(4):332–54.

Gerlach KE, Burton HW, Dorn JM, Leddy JJ, Horvath PJ. Fat intake and injury in female runners. J Int Soc Sports Nutr. 2008 Jan 3;5:1. doi: 10.1186/1550-2783-5-1.

Goedecke JH, Christie C, Wilson G, Dennis SC, Noakes TD, Hopkins WG, and Lambert EV. Metabolic adaptations to a high-fat diet in endurance cyclists. Metabolism 48: 1509–1517, 1999.

Goedecke JH, Clark VR, Noakes TD, Lambert EV. The effects of medium-chain triacylglycerol and carbohydrate ingestion on ultra-endurance exercise performance. Int J Sport Nutr Exerc Metab. 2005 Feb;15(1):15–27.

Gow ML, Ho M, Burrows TL, Baur LA, Stewart L, Hutchesson MJ, Cowell CT, Collins CE, Garnett SP. Impact of dietary macronutrient distribution on BMI and cardiometabolic outcomes in overweight and obese children and adolescents: a systematic review. Nutr Rev. 2014 Jun 11.

Greene NP, Martin SE, Crouse SF. Acute exercise and training alter blood lipid and lipoprotein profiles differently in overweight and obese men and women. Obesity (Silver Spring). 2012 Aug;20(8):1618–27.

- Hall KD.** What is the Required Energy Deficit per unit Weight Loss? *Int J Obes (Lond)* 2008; 32: 573–576.
- Hammond VJ, O'Donnell VB.** Esterified eicosanoids: generation, characterization and function. *Biochim Biophys Acta*. 2012 Oct;1818(10):2403–12
- Han JR, Deng B, Sun J, Chen CG, Corkey BE, Kirkland JL, Ma J, Guo W.** Effects of dietary medium-chain triglyceride on weight loss and insulin sensitivity in a group of moderately overweight free-living type 2 diabetic Chinese subjects. *Metabolism*. 2007 Jul;56(7):985–91.
- Hargreaves M, Hawley JA, Jeukendrup A.** Pre-exercise carbohydrate and fat ingestion: effects on metabolism and performance. *J Sports Sci*. 2004 Jan;22(1):31–8.
- Harvard 2005.** Making sense of cholesterol tests. <http://www.health.harvard.edu/fhg/updates/update0205c.shtml>
- Hauner H, Bechthold A, Boeing H, Brönstrup A, Buyken A, Leschik-Bonnet E ym. ; German Nutrition Society.** Evidence-based guideline of the German Nutrition Society: carbohydrate intake and prevention of nutrition-related diseases. *Ann Nutr Metab*. 2012;60 Suppl 1:1–58
- Havemann L, West SJ, Goedecke JH, Macdonald IA, St Clair Gibson A, Noakes TD, Lambert EV.** Fat adaptation followed by carbohydrate loading compromises high-intensity sprint performance. *J Appl Physiol (1985)*. 2006 Jan;100(1):194–202.
- Hawley JA, Burke LM, Angus DJ, Fallon KE, Martin DT, Febbraio MA.** Effect of altering substrate availability on metabolism and performance during intense exercise. *Br J Nutr*. 2000 Dec;84(6):829–38.
- Hayes KC, Pronczuk A.** Replacing trans fat: the argument for palm oil with a cautionary note on interesterification. *J Am Coll Nutr*. 2010 Jun;29(3 Suppl):253S-284S. Review.
- Hjerpsted J, Leedo E, Tholstrup T.** Cheese intake in large amounts lowers LDL-cholesterol concentrations compared with butter intake of equal fat content. *Am J Clin Nutr*. 2011 Dec;94(6):1479–84.
- Hooper L, Summerbell CD, Thompson R, Sills D, Roberts FG, Moore HJ, Davey Smith G.** Reduced or modified dietary fat for preventing cardiovascular disease. *Cochrane Database Syst Rev*. 2012 May 16;5:CD002137. (a)
- Hooper L, Abdelhamid A, Moore HJ, Douthwaite W, Skeaff CM, Summerbell CD.** Effect of reducing total fat intake on body weight: systematic review and meta-analysis of randomised controlled trials and cohort studies. *BMJ*. 2012 Dec 6;345:e7666. (b)
- Horvath PJ, Eagen CK, Ryer-Calvin SD, Pendergast DR.** The effects of varying dietary fat on the nutrient intake in male and female runners. *J Am Coll Nutr*. 2000 Feb;19(1):42–51.
- Hu FB.** Are refined carbohydrates worse than saturated fat? *Am J Clin Nutr*. 2010 Jun;91(6):1541–2.
- Hurst S, Zainal Z, Caterson B, Hughes CE, Harwood JL.** Dietary fatty acids and arthritis. *Prostaglandins Leukot Essent Fatty Acids*. 2010 Apr-Jun;82(4–6):315–8.
- Huth PJ, Park KM.** Influence of dairy product and milk fat consumption on cardiovascular disease risk: a review of the evidence. *Adv Nutr*. 2012 May 1;3(3):266–85.
- Hämäläinen EK, Adlercreutz H, Puska P, Pietinen P.** Decrease of serum total and free testosterone during a low-fat high-fibre diet. *J Steroid Biochem*. 1983 Mar;18(3):369–70.
- Høstmark AT, Tomten SE.** The Oslo health study: cheese intake was negatively associated with the metabolic syndrome. *J Am Coll Nutr*. 2011 Jun;30(3):182–90.
- IOC.** Nutrition Working Group of the International Olympic Committee. Nutrition for Athletes. A practical guide to eating for health and performance. Revised and updated 2012.
- Jackson CL, Hu FB.** Long-term associations of nut consumption with body weight and obesity. *Am J Clin Nutr*. 2014 Jun 4.
- Jakobsen MU, O'Reilly EJ, Heitmann BL, Pereira MA, Bälter K, Fraser GE, Goldbourt U, Hallmans G, ym.** Major types of dietary fat and risk of coronary heart disease: a pooled analysis of 11 cohort studies *Am J Clin Nutr*. 2009 May;89(5):1425–32.

Jakobsen MU, Dethlefsen C, Joensen AM, Stegger J, Tjønneland A, Schmidt EB, Overvad K. Intake of carbohydrates compared with intake of saturated fatty acids and risk of myocardial infarction: importance of the glycemic index. *Am J Clin Nutr.* 2010 Jun;91(6):1764–8.

Jakulj F, Zernicke K, Bacon SL, van Wielingen LE, Key BL, West SG, Campbell TS. A high-fat meal increases cardiovascular reactivity to psychological stress in healthy young adults. *J Nutr.* 2007 Apr;137(4):935–9.

Jeukendrup AE, Aldred S. Fat supplementation, health, and endurance performance. *Nutrition.* 2004 Jul-Aug;20(7–8):678–88.

Johnson NA, Stannard SR, Thompson MW. Muscle triglyceride and glycogen in endurance exercise: implications for performance. *Sports Med.* 2004;34(3):151–64.

Jones PJ, Jew S, AbuMweis S. The effect of dietary oleic, linoleic, and linolenic acids on fat oxidation and energy expenditure in healthy men. *Metabolism.* 2008 Sep;57(9):1198–203.

Josse AR, Kendall CW, Augustin LS, Ellis PR, Jenkins DJ. Almonds and postprandial glycemia--a dose-response study. *Metabolism.* 2007 Mar;56(3):400–4.

Júdice PB, Silva AM, Magalhães JP, Matias CN, Sardinha LB. Sedentary behaviour and adiposity in elite athletes. *J Sports Sci.* 2014 Jun 10:1–8.

Kanerva N, Kaartinen NE, Rissanen H, Knek P, Eriksson JG, Sääksjärvi K, Sundvall J, Männistö S. Associations of the Baltic Sea diet with cardiometabolic risk factors – a meta-analysis of three Finnish studies. *Br J Nutr.* 2014 May 27:1–11.

Kathiresan S, Otvos JD, Sullivan LM, et al. Increased small low-density lipoprotein particle number: a prominent feature of the metabolic syndrome in the Framingham Heart Study. *Circulation.* 2006;113:20–29.

Keith SW, Redden DT, Katzmarzyk PT, Boggiano MM, Hanlon EC, Benca RM, Ruden D, Pietrobelli A ym. Putative contributors to the secular increase in obesity: exploring the roads less traveled. *Int J Obes (Lond).* 2006 Nov;30(11):1585–94.

Kelley GA, Kelley KS. Impact of progressive resistance training on lipids and lipoproteins in adults: a meta-analysis of randomized controlled trials. *Prev Med.* 2009 Jan;48(1):9–19.

Kelley GA, Kelley KS, Roberts S, Haskell W. Comparison of aerobic exercise, diet or both on lipids and lipoproteins in adults: a meta-analysis of randomized controlled trials. *Clin Nutr.* 2012 Apr;31(2):156–67.

Kendall CW, Esfahani A, Josse AR, Augustin LS, Vidgen E, Jenkins DJ. The glycemic effect of nut-enriched meals in healthy and diabetic subjects. *Nutr Metab Cardiovasc Dis.* 2011 Jun;21 Suppl 1:S34–9.

Kien CL, Bunn JY, Ugrasbul F. Increasing dietary palmitic acid decreases fat oxidation and daily energy expenditure. *Am J Clin Nutr.* 2005 Aug;82(2):320–6.

Kien CL, Bunn JY, Tompkins CL, Dumas JA, Crain KI, Ebenstein DB, Koves TR, Muoio DM. Substituting dietary monounsaturated fat for saturated fat is associated with increased daily physical activity and resting energy expenditure and with changes in mood. *Am J Clin Nutr.* 2013 Apr;97(4):689–97.

King NA, Blundell JE. High-fat foods overcome the energy expenditure induced by high-intensity cycling or running. *Eur J Clin Nutr.* 1995 Feb;49(2):114–23.

Kinosian B, Glick H, Preiss L, Puder KL. Cholesterol and coronary heart disease: Predicting risks in men by changes in levels and ratios. *J Investig Med* 1995;43:443–50

Kodama S, Tanaka S, Saito K, et al. Effect of aerobic exercise training on serum levels of high-density lipoprotein cholesterol: a meta-analysis. *Arch Intern Med.* 2007;167:999–1008.

Koh-Banerjee P, Chu NF, Spiegelman D, Rosner B, Colditz G, Willett W, Rimm E. Prospective study of the association of changes in dietary intake, physical activity, alcohol consumption, and smoking with 9-y gain in waist circumference among 16 587 US men. *Am J Clin Nutr.* 2003 Oct;78(4):719–27.

Kratz M, Baars T, Guyenet S. The relationship between high-fat dairy consumption and obesity, cardiovascular, and metabolic disease. *Eur J Nutr.* 2013 Feb;52(1):1–24.

Krishnan S, Cooper JA. Effect of dietary fatty acid composition on substrate utilization and body weight maintenance in humans. *Eur J Nutr.* 2014 Apr;53(3):691–710.

Laake I, Pedersen JI, Selmer R, Kirkhus B, Lindman AS, Tverdal A, Veierød MB. A prospective study of intake of trans-fatty acids from ruminant fat, partially hydrogenated vegetable oils, and marine oils and mortality from CVD. *Br J Nutr.* 2012 Aug;108(4):743–54.

Lada AT, Rudel LL. Associations of low density lipoprotein particle composition with atherogenicity. *Curr Opin Lipidol.* 2004 Feb;15(1):19–24. Review.

Laitinen K, Gylling H. Dose-dependent LDL-cholesterol lowering effect by plant stanol ester consumption: clinical evidence. *Lipids Health Dis.* 2012 Oct 22;11:140.

Lambert EV, Goedecke JH. The role of dietary macronutrients in optimizing endurance performance. *Curr Sports Med Rep.* 2003 Aug;2(4):194–201.

Larson-Meyer DE, Borkhsenious ON, Gullett JC, Russell RR, Devries MC, Smith SR, Ravussin E. Effect of dietary fat on serum and intramyocellular lipids and running performance. *Med Sci Sports Exerc.* 2008 May;40(5):892–902.

Lawrence GD. Dietary fats and health: dietary recommendations in the context of scientific evidence. *Adv Nutr.* 2013 May 1;4(3):294–302.

Lawton CL, Delargy HJ, Brockman J, Smith FC, Blundell JE. The degree of saturation of fatty acids influences post-ingestive satiety. *Br J Nutr.* 2000;83:473–82.

Liau KM, Lee YY, Chen CK, Rasool AH. An open-label pilot study to assess the efficacy and safety of virgin coconut oil in reducing visceral adiposity. *ISRN Pharmacol.* 2011;2011:949686.

Loucks AB. Energy availability in athletes. IOC Consensus In Sports Nutrition Conference Lausanne, Switzerland 25th–27th October 2010

Lowery LM. Dietary Fat and Sports Nutrition: A Primer. *J Sports Sci Med.* 2004 Sep 1;3(3):106–117.

Magalhães B, Peleteiro B, Lunet N. Dietary patterns and colorectal cancer: systematic review and meta-analysis. *Eur J Cancer Prev.* 2012 Jan;21(1):15–23.

Manolopoulos KN, Karpe F, Frayn KN. Gluteofemoral body fat as a determinant of metabolic health. *Int J Obes (Lond).* 2010 Jun;34(6):949–59.

Marina AM, ym. Virgin coconut oil: emerging functional food oil. *Trends in Food Science & Technology* 20 (2009) 481–487

Marina AM, Man YB, Nazimah SA, Amin I. Antioxidant capacity and phenolic acids of virgin coconut oil. *Int J Food Sci Nutr.* 2009;60 Suppl 2:114–23.

Maughan RJ. Fundamentals of Sports Nutrition: Application to Sports Drinks. Kirjassa: Sports Drinks – Basic Science and Practical Aspects. Maughan RJ, Murray R (toim.). CRC Press. Boca Raton, USA, 2001.

Mensink RP, Zock PL, Kester AD, Katan MB. Effects of dietary fatty acids and carbohydrates on the ratio of serum total to HDL cholesterol and on serum lipids and apolipoproteins: a meta-analysis of 60 controlled trials. *Am J Clin Nutr.* 2003;77:1146–55

Mente A, de Koning L, Shannon HS, Anand SS. A systematic review of the evidence supporting a causal link between dietary factors and coronary heart disease. *Arch Intern Med.* 2009 Apr 13;169(7):659–69.

Micha R, Mozaffarian D. Trans fatty acids: effects on metabolic syndrome, heart disease and diabetes. *Nat Rev Endocrinol.* 2009 Jun;5(6):335–44.

Micha R, Mozaffarian D. Saturated fat and cardiometabolic risk factors, coronary heart disease, stroke, and diabetes: a fresh look at the evidence. *Lipids.* 2010 Oct;45(10):893–905.

Micha R, Michas G, Mozaffarian D. Unprocessed red and processed meats and risk of coronary artery disease and type 2 diabetes--an updated review of the evidence. *Curr Atheroscler Rep.* 2012 Dec;14(6):515–24.

Micha R, Michas G, Lajous M, Mozaffarian D. Processing of meats and cardiovascular risk: time to focus on preservatives. *BMC Med.* 2013 May 23;11:136.

Mickleborough TD, Head SK, Lindley MR. Exercise-induced asthma: nutritional management. *Curr Sports Med Rep.* 2011 Jul;10(4):197–202

Monnier L, Mas E, Ginet C, Michel F, Villon L, Cristol JP, Colette C. Activation of oxidative stress by acute glucose fluctuations compared with sustained chronic hyperglycemia in patients with type 2 diabetes. *JAMA*. 2006 Apr 12;295(14):1681–7.

Moreno Alcalde S, Ruiz-Roso B, Pérez-Olleros L, Belmonte Cortés S. Trans fatty acid content in foods marketed in the community of Madrid(Spain). *Nutr Hosp*. 2014 Jan 1;29(1):180–6.

Morgan J, Carey C, Lincoff A, Capuzzi D. High-density lipoprotein subfractions and risk of coronary artery disease. *Curr Atheroscler Rep*. 2004;6:359–365

Motard-Bélanger A, Charest A, Grenier G, Paquin P, Chouinard Y, Lemieux S, Couture P, Lamarche B. Study of the effect of trans fatty acids from ruminants on blood lipids and other risk factors for cardiovascular disease. *Am J Clin Nutr*. 2008 Mar;87(3):593–9.

Moussavi N, Gavino V, Receveur O. Could the quality of dietary fat, and not just its quantity, be related to risk of obesity? *Obesity (Silver Spring)*. 2008 Jan;16(1):7–15.

Mozaffarian D. Trans fatty acids – effects on systemic inflammation and endothelial function. *Atheroscler Suppl*. 2006 May;7(2):29–32. Epub 2006 May 18.

Mozaffarian D, Clarke R. Quantitative effects on cardiovascular risk factors and coronary heart disease risk of replacing partially hydrogenated vegetable oils with other fats and oils. *Eur J Clin Nutr*. 2009;63(Suppl 2):S22–S33.

Mozaffarian D, Aro A, Willett WC. Health effects of trans-fatty acids: experimental and observational evidence. *Eur J Clin Nutr*. 2009 May;63 Suppl 2:S5–21

Mozaffarian D, Micha R, Wallace S. Effects on coronary heart disease of increasing polyunsaturated fat in place of saturated fat: a systematic review and meta-analysis of randomized controlled trials. *PLoS Med*. 2010 Mar 23;7(3):e1000252. (a)

Mozaffarian D, Cao H, King IB, Lemaitre RN, Song X, Siscovick DS, Hotamisligil GS. Trans-palmitoleic acid, metabolic risk factors, and new-onset diabetes in U.S. adults: a cohort study. *Ann Intern Med*. 2010 Dec 21;153(12):790–9. (b)

Mozaffarian D, Hao T, Rimm EB, Willett WC, Hu FB. Changes in diet and lifestyle and long-term weight gain in women and men. *N Engl J Med*. 2011 Jun 23;364(25):2392–404

Mozaffarian D, de Oliveira Otto MC, Lemaitre RN, Fretts AM, Hotamisligil G, Tsai MY, Siscovick DS ym. trans-Palmitoleic acid, other dairy fat biomarkers, and incident diabetes: the Multi-Ethnic Study of Atherosclerosis (MESA). *Am J Clin Nutr*. 2013 Apr;97(4):854–61

Murakami I, Sakuragi T, Uemura H, Menda H, Shindo M, Tanaka H. Significant effect of a pre-exercise high-fat meal after a 3-day high-carbohydrate diet on endurance performance. *Nutrients*. 2012 Jul;4(7):625–37.

Nestel PJ, Chronopoulos A, Cehun M. Dairy fat in cheese raises LDL cholesterol less than that in butter in mildly hypercholesterolaemic subjects. *Eur J Clin Nutr*. 2005 Sep;59(9):1059–63.

Nevin KG, Rajamohan T. Beneficial effects of virgin coconut oil on lipid parameters and in vitro LDL oxidation. *Clin Biochem*. 2004 Sep;37(9):830–5.

Nyberg ST, Heikkilä K, Fransson EI, Alfredsson L, De Bacquer D, Bjorner JB, ym.; IPD-Work Consortium. Job strain in relation to body mass index: pooled analysis of 160 000 adults from 13 cohort studies. *J Intern Med*. 2012 Jul;272(1):65–73.

Ohlsson L. Dairy products and plasma cholesterol levels. *Food Nutr Res*. 2010 Aug 19;54.

O'Keefe JH, Gheewala NM, O'Keefe JO. Dietary strategies for improving post-prandial glucose, lipids, inflammation, and cardiovascular health. *J Am Coll Cardiol*. 2008 Jan 22;51(3):249–55.

Ortega JF, Fernández-Elías VE, Hamouti N, Mora-Rodriguez R. Increased blood cholesterol after a high saturated fat diet is prevented by aerobic exercise training. *Appl Physiol Nutr Metab*. 2013 Jan;38(1):42–8.

Pan A, Sun Q, Manson JE, Willett WC, Hu FB. Walnut consumption is associated with lower risk of type 2 diabetes in women. *J Nutr*. 2013 Apr;143(4):512–8.

- Parlesak A, Eckoldt J, Winkler K, Bode CJ, Schäfer C.** Intercorrelations of lipoprotein subfractions and their covariation with lifestyle factors in healthy men. *J Clin Biochem Nutr.* 2014 May;54(3):174–80.
- Pérez-Escamilla R, Obbagy JE, Altman JM, Essery EV, McGrane MM, Wong YP, Spahn JM, Williams CL.** Dietary energy density and body weight in adults and children: a systematic review. *J Acad Nutr Diet.* 2012 May;112(5):671–84.
- Pfeiffer B, Stellingwerff T, Zaltas E, Jeukendrup AE.** Oxidation of solid versus liquid CHO sources during exercise. *Med Sci Sports Exerc.* 2010 Nov;42(11):2030–7.
- Piers LS, Walker KZ, Stoney RM, Soares MJ, O'Dea K.** The influence of the type of dietary fat on postprandial fat oxidation rates: monounsaturated (olive oil) vs saturated fat (cream). *Int J Obes Relat Metab Disord.* 2002 Jun;26(6):814–21.
- Poppitt SD, Strik CM, MacGibbon AK, McArdle BH, Budgett SC, McGill AT.** Fatty acid chain length, postprandial satiety and food intake in lean men. *Physiol Behav.* 2010 Aug 4;101(1):161–7.
- Power O, Jakeman P, Fitzgerald RJ.** Antioxidative peptides: enzymatic production, in vitro and in vivo antioxidant activity and potential applications of milk-derived antioxidative peptides. *Amino Acids.* 2013 Mar;44(3):797–820.
- Prospective Studies Collaboration, Lewington S, Whitlock G, Clarke R, Sherliker P.** Blood cholesterol and vascular mortality by age, sex, and blood pressure: a meta-analysis of individual data from 61 prospective studies with 55,000 vascular deaths. *Lancet.* 2007 Dec 1;370(9602):1829–39. Review.
- Przybylski O, Aladedunye FA.** Formation of trans fats during food preparation. *Can J Diet Pract Res.* 2012 Summer;73(2):98–101.
- Rauch HG, Hawley JA, Woodey M, Noakes TD, Dennis SC.** Effects of ingesting a sports bar versus glucose polymer on substrate utilisation and ultra-endurance performance. *Int J Sports Med.* 1999 May;20(4):252–7.
- Rego Costa AC, Rosado EL, Soares-Mota M.** Influence of the dietary intake of medium chain triglycerides on body composition, energy expenditure and satiety: a systematic review. *Nutr Hosp.* 2012 Jan-Feb;27(1):103–8
- Rehrer NJ, van Kemenade M, Meester W, Brouns F, Saris WH.** Gastrointestinal complaints in relation to dietary intake in triathletes. *Int J Sport Nutr.* 1992 Mar;2(1):48–59.
- Risérus U.** Trans fatty acids and insulin resistance. *Atheroscler Suppl.* 2006 May;7(2):37–9. Epub 2006 May 18.
- Risérus U, Willett WC, Hu FB.** Dietary fats and prevention of type 2 diabetes. *Prog Lipid Res.* 2009 Jan;48(1):44–51.
- Ritvanen T, Putkonen T, Peltonen K.** A comparative study of the fatty acid composition of dairy products and margarines with reduced or substituted fat content. *Food and Nutrition Sciences, Vol. 3, 2012,* pp. 1189–1196
- Rodriguez-Leyva D, Dupasquier CM, McCullough R, Pierce GN.** The cardiovascular effects of flaxseed and its omega-3 fatty acid, alpha-linolenic acid. *Can J Cardiol.* 2010 Nov;26(9):489–96. Review.
- Roe M, Pinchen H, Church S, Elahi S, Walker M, Farron-Wilson M, Buttriss J, Finglas P.** Trans fatty acids in a range of UK processed foods. *Food Chem.* 2013 Oct 1;140(3):427–31.
- Rolls BJ.** The role of energy density in the overconsumption of fat. *J Nutr.* 2000 Feb;130(2S Suppl):268S–271S.
- Rolls BJ, Roe LS, Meengs JS.** Portion size can be used strategically to increase vegetable consumption in adults. *Am J Clin Nutr.* 2010 Apr;91(4):913–22.
- Rowlands DS, Hopkins WG.** Effect of high-fat, high-carbohydrate, and high-protein meals on metabolism and performance during endurance cycling. *Int J Sport Nutr Exerc Metab.* 2002 Sep;12(3):318–35.
- Roy BD, Tarnopolsky MA.** Influence of differing macronutrient intakes on muscle glycogen resynthesis after resistance exercise. *J Appl Physiol (1985).* 1998 Mar;84(3):890–6.
- Sacks FM, Katan M.** Randomized clinical trials on the effects of dietary fat and carbohydrate on plasma lipoproteins and cardiovascular disease. *Am J Med.* 2002 Dec 30;113 Suppl 9B:13S–24S.
- Sacks FM, Bray GA, Carey VJ, Smith SR, Ryan DH, Anton SD, McManus K, Champagne CM, Bishop LM ym.** Comparison of weight-loss diets with different compositions of fat, protein, and carbohydrates. *N Engl J Med.* 2009 Feb 26;360(9):859–73.

- Schrauwen-Hinderling VB, Hesselink MK, Schrauwen P, Kooi ME.** Intramyocellular lipid content in human skeletal muscle. *Obesity (Silver Spring)*. 2006 Mar;14(3):357–67. Review.
- Schwingshackl L, Strasser B, Hoffmann G.** Effects of monounsaturated fatty acids on cardiovascular risk factors: a systematic review and meta-analysis. *Ann Nutr Metab*. 2011;59(2–4):176–86
- Schwingshackl L, Hoffmann G.** Comparison of the long-term effects of high-fat v. low-fat diet consumption on cardiometabolic risk factors in subjects with abnormal glucose metabolism: a systematic review and meta-analysis. *Br J Nutr*. 2014 Jun;111(12):2047–58.
- Singh P, Somers VK, Romero-Corral A, Sert-Kuniyoshi FH, Pusalavidyasagar S, Davison DE, Jensen MD.** Effects of weight gain and weight loss on regional fat distribution. *Am J Clin Nutr*. 2012 Aug;96(2):229–33.
- Siri-Tarino PW.** Effects of diet on high-density lipoprotein cholesterol. *Curr Atheroscler Rep*. 2011 Dec;13(6):453–60
- Siri-Tarino PW, Sun Q, Hu FB, Krauss RM.** Meta-analysis of prospective cohort studies evaluating the association of saturated fat with cardiovascular disease. *Am J Clin Nutr*. 2010 Mar;91(3):535–46.
- Sjögren P, Rosell M, Skoglund-Andersson C, Zdravkovic S, Vessby B, de Faire U, Hamsten A, Hellenius ML ym.** Milk-derived fatty acids are associated with a more favorable LDL particle size distribution in healthy men. *J Nutr*. 2004 Jul;134(7):1729–35.
- Stender S, Astrup A, Dyerberg J.** Ruminant and industrially produced trans fatty acids: health aspects. *Food Nutr Res*. 2008;52.
- Stender S, Astrup A, Dyerberg J.** A trans European Union difference in the decline in trans fatty acids in popular foods: a market basket investigation. *BMJ Open*. 2012 Sep 17;2(5).
- Strik CM, Lithander FE, McGill AT, MacGibbon AK, McArdle BH, Poppitt SD.** No evidence of differential effects of SFA, MUFA or PUFA on post-ingestive satiety and energy intake: a randomised trial of fatty acid saturation. *Nutr J*. 2010 May 24;9:24.
- Stuemple KJ, Hoffman MD, Weschler LB, Rogers IR, Hew-Butler T.** Race diet of finishers and non-finishers in a 100 mile (161 km) mountain footrace. *J Am Coll Nutr*. 2011 Dec;30(6):529–35.
- Swinburn BA, Sacks G, Lo SK, Westerterp KR, Rush EC, Rosenbaum M, Luke A, Schoeller DA ym.** Estimating the changes in energy flux that characterize the rise in obesity prevalence. *Am J Clin Nutr*. 2009 Jun;89(6):1723–8.
- Tan SY, Mattes RD.** Appetitive, dietary and health effects of almonds consumed with meals or as snacks: a randomized, controlled trial. *Eur J Clin Nutr*. 2013 Nov;67(11):1205–14.
- Tchernof A, Després JP.** Pathophysiology of human visceral obesity: an update. *Physiol Rev*. 2013 Jan;93(1):359–404.
- Theuwissen E, Mensink RP.** Water-soluble dietary fibers and cardiovascular disease. *Physiol Behav*. 2008 May 23;94(2):285–92.
- Theuwissen E, Plat J, van der Kallen CJ, van Greevenbroek MM, Mensink RP.** Plant stanol supplementation decreases serum triacylglycerols in subjects with overt hypertriglyceridemia. *Lipids*. 2009 Dec;44(12):1131–40.
- Unlu NZ, Bohn T, Clinton SK, Schwartz SJ.** Carotenoid absorption from salad and salsa by humans is enhanced by the addition of avocado or avocado oil. *J Nutr*. 2005 Mar;135(3):431–6
- U.S. Department of Agriculture, U.S. Department of Health.** Dietary Guidelines for Americans 2010. www.dietaryguidelines.gov
- Vadivel V, Kunyanga CN, Biesalski HK.** Health benefits of nut consumption with special reference to body weight control. *Nutrition*. 2012 Nov-Dec;28(11–12):1089–97
- van Loon LJ, Koopman R, Stegen JH, Wagenmakers AJ, Keizer HA, Saris WH.** Intramyocellular lipids form an important substrate source during moderate intensity exercise in endurance-trained males in a fasted state. *J Physiol*. 2003 Dec 1;553(Pt 2):611–25.
- Vega-López S, Ausman LM, Jalbert SM, Erkkilä AT, Lichtenstein AH.** Palm and partially hydrogenated soybean oils adversely alter lipoprotein profiles compared with soybean and canola oils in moderately hyperlipidemic subjects. *Am J Clin Nutr*. 2006 Jul;84(1):54–62.

Venkatraman JT, Leddy J, Pendergast D. Dietary fats and immune status in athletes: clinical implications. *Med Sci Sports Exerc.* 2000 Jul;32(7 Suppl):S389–95. Review.

Venkatraman JT, Pendergast DR. Effect of dietary intake on immune function in athletes. *Sports Med.* 2002;32(5):323–37. Review.

Vesper HW, Kuiper HC, Mirel LB, Johnson CL, Pirkle JL. Levels of plasma trans-fatty acids in non-Hispanic white adults in the United States in 2000 and 2009. *JAMA.* 2012 Feb 8;307(6):562–3.

Villarino, B. J., Dy, L. M., & Lizada, C. C. (2007). Descriptive sensory evaluation of virgin coconut oil and refined, bleached and deo- dorized coconut oil. *LWT-Food Science and Technology*, 40, 193–199

Volek JS, Kraemer WJ, Bush JA, Incledon T, Boetes M. Testosterone and cortisol in relationship to dietary nutrients and resistance exercise. *J Appl Physiol (1985).* 1997 Jan;82(1):49–54.

Volek JS, Gómez AL, Love DM, Avery NG, Sharman MJ, Kraemer WJ. Effects of a high-fat diet on postabsorptive and postprandial testosterone responses to a fat-rich meal. *Metabolism.* 2001 Nov;50(11):1351–5.

VRN. Terveytä ruuasta – Suomalaiset ravitsemussuositukset 2014. Juvenes Oy, Helsinki 2014.

Walker TB, Parker MJ. Lessons from the War on Dietary Fat. *J Am Coll Nutr.* 2014 Jun 9:1–5.

Weigle DS, Breen PA, Matthys CC, Callahan HS, Meeuws KE, Burden VR, Purnell JQ. A high-protein diet induces sustained reductions in appetite, ad libitum caloric intake, and body weight despite compensatory changes in diurnal plasma leptin and ghrelin concentrations. *Am J Clin Nutr.* 2005 Jul;82(1):41–8.

Willett WC. Dietary fat plays a major role in obesity: no. *Obes Rev.* 2002 May;3(2):59–68. Review.

Willett W, Mozaffarian D. Ruminant or industrial sources of trans fatty acids: public health issue or food label skirmish? *Am J Clin Nutr.* 2008 Mar;87(3):515–6.

Wu JH, Micha R, Imamura F, Pan A, Biggs ML, Ajaz O, Djousse L, Hu FB, Mozaffarian D. Omega-3 fatty acids and incident type 2 diabetes: a systematic review and meta-analysis. *Br J Nutr.* 2012 Jun;107 Suppl 2:S214–27

Wu L, Ma D, Walton-Moss B, He Z. Effects of low-fat diet on serum lipids in premenopausal and postmenopausal women: a meta-analysis of randomized controlled trials. *Menopause.* 2014 Jan;21(1):89–99.

Wycherley TP, Moran LJ, Clifton PM, Noakes M, Brinkworth GD. Effects of energy-restricted high-protein, low-fat compared with standard-protein, low-fat diets: a meta-analysis of randomized controlled trials. *Am J Clin Nutr.* 2012 Dec;96(6):1281–98.

Yeo WK, Carey AL, Burke L, Spriet LL, Hawley JA. Fat adaptation in well-trained athletes: effects on cell metabolism. *Appl Physiol Nutr Metab.* 2011 Feb;36(1):12–22.

Zehnder M, Christ ER, Ith M, Acheson KJ, Pouteau E, Kreis R, Trepp R, Diem P, Boesch C, Décombaraz J. Intramyocellular lipid stores increase markedly in athletes after 1.5 days lipid supplementation and are utilized during exercise in proportion to their content. *Eur J Appl Physiol.* 2006 Nov;98(4):341–54.

Lähdeluettelo luku 8

Adams WM, Ferraro EM, Huggins RA, Casa DJ. Influence of body mass loss on changes in heart rate during exercise in the heat: a systematic review. *J Strength Cond Res.* 2014 Aug;28(8):2380–9.

Adan A. Cognitive performance and dehydration. *J Am Coll Nutr.* 2012 Apr;31(2):71–8.

Adolph EF, Brown AH, Goddard DR, et al. Physiology of man in the desert. New York: Interscience, 1947:16–33, 226–241, 326–42.

Ali A, Gardiner R, Foskett A, Gant N. Fluid balance, thermoregulation and sprint and passing skill performance in female soccer players. *Scand J Med Sci Sports.* 2011 Jun;21(3):437–45.

American College of Sports Medicine, Sawka MN, Burke LM, Eichner ER, Maughan RJ, Montain SJ, Stachenfeld NS. American College of Sports Medicine position stand. Exercise and fluid replacement. *Med Sci Sports Exerc.* 2007 Feb;39(2):377–90.

Anastasiou CA, Kavouras SA, Arnaoutis G, Gioxari A, Kollia M, Botoula E, Sidossis LS. Sodium replacement and plasma sodium drop during exercise in the heat when fluid intake matches fluid loss. *J Athl Train.* 2009 Mar-Apr;44(2):117–23.

Armstrong LE, Soto JA, Hacker FT Jr, Casa DJ, Kavouras SA, Maresh CM. Urinary indices during dehydration, exercise, and rehydration. *Int J Sport Nutr.* 1998 Dec;8(4):345–55.

Armstrong LE, Whittlesey MJ, Casa DJ, Elliott TA, Kavouras SA, Keith NR, Maresh CM. No effect of 5% hypohydration on running economy of competitive runners at 23 degrees C. *Med Sci Sports Exerc.* 2006 Oct;38(10):1762–9.

Armstrong LE, Ganio MS, Casa DJ, Lee EC, McDermott BP, Klau JF, Jimenez L, Le Bellego L, Chevillotte E, Lieberman HR. Mild dehydration affects mood in healthy young women. *J Nutr.* 2012 Feb;142(2):382–8.

Armstrong LE, Johnson EC, Ganio MS, Judelson DA, Vingren JL, Kupchak BR, Kunes LJ, Muñoz CX, McKenzie AL, Williamson KH. Effective body water and body mass changes during summer ultra-endurance road cycling. *J Sports Sci.* 2014 Jul 3:1–11. (a)

Armstrong LE, Johnson EC, Kunes LJ, Ganio MS, Judelson DA, Kupchak BR, Vingren JL, Munoz CX, Huggins RA, Hydren JR,ym. Drinking to Thirst Versus Drinking Ad Libitum During Road Cycling. *J Athl Train.* 2014 Aug 6. (b)

Arnaoutis G, Kavouras SA, Kotsis YP, Tsekouras YE, Makrillo M, Bardis CN. Ad libitum fluid intake does not prevent dehydration in suboptimally hydrated young soccer players during a training session of a summer camp. *Int J Sport Nutr Exerc Metab.* 2013 Jun;23(3):245–51.

Arnaoutis G, Kavouras SA, Angelopoulou A, Skouliki C, Bismikou S, Mourtakos S, Sidossis LS. FLUID BALANCE DURING TRAINING IN ELITE YOUNG ATHLETES OF DIFFERENT SPORTS. *J Strength Cond Res.* 2014 Feb 10.

Askew EW. Environmental and physical stress and nutrient requirements. *Am J Clin Nutr.* 1995 Mar;61(3 Suppl):631S-637S.

Baker LB, Dougherty KA, Chow M, Kenney WL. Progressive dehydration causes a progressive decline in basketball skill performance. *Med Sci Sports Exerc.* 2007 Jul;39(7):1114–23. (a)

Baker LB, Conroy DE, Kenney WL. Dehydration impairs vigilance-related attention in male basketball players. *Med Sci Sports Exerc.* 2007 Jun;39(6):976–83. (b)

Barnes MJ. Alcohol: impact on sports performance and recovery in male athletes. *Sports Med.* 2014 Jul;44(7):909–19.

Bassett DR Jr, Howley ET. Limiting factors for maximum oxygen uptake and determinants of endurance performance. *Med Sci Sports Exerc.* 2000 Jan;32(1):70–84.

Beis LY, Polyviou T, Malkova D, Pitsiladis YP. The effects of creatine and glycerol hyperhydration on running economy in well trained endurance runners. *J Int Soc Sports Nutr.* 2011 Dec 16;8(1):24.

Beis LY, Wright-Whyte M, Fudge B, Noakes T, Pitsiladis YP. Drinking behaviors of elite male runners during marathon competition. *Clin J Sport Med.* 2012 May;22(3):254–61.

Blank MC, Bedarf JR, Russ M, Grosch-Ott S, Thiele S, Unger JK. Total body Na(+)-depletion without hyponatraemia can trigger overtraining-like symptoms with sleeping disorders and increasing blood pressure: explorative case and literature study. *Med Hypotheses.* 2012 Dec;79(6):799–804.

Brade CJ, Dawson BT, Wallman KE. Effect of pre-cooling on repeat-sprint performance in seasonally acclimatised males during an outdoor simulated team-sport protocol in warm conditions. *J Sports Sci Med.* 2013 Sep 1;12(3):565–70. eCollection 2013.

Brouns F, Beckers E. Is the gut an athletic organ? Digestion, absorption and exercise. *Sports Med.* 1993;15:242–257.

Burke LM, Collier GR, Broad EM, Davis PG, Martin DT, Sanigorski AJ, Hargreaves M. Effect of alcohol intake on muscle glycogen storage after prolonged exercise. *J Appl Physiol* (1985). 2003 Sep;95(3):983–90.

Byrne C, Lee JK, Chew SA, Lim CL, Tan EY. Continuous thermoregulatory responses to mass-participation distance running in heat. *Med Sci Sports Exerc.* 2006 May;38(5):803–10.

Calbet JA, MacLean DA. Role of caloric content on gastric emptying in humans. *J Physiol.* 1997 Jan 15;498 (Pt 2):553–9.

Chen TI, Tsai PH, Lin JH, Lee NY, Liang MT. Effect of short-term heat acclimation on endurance time and skin blood flow in trained athletes. *Open Access J Sports Med.* 2013 Jun 18;4:161–70.

Cheuvront SN, Carter R 3rd, Castellani JW, Sawka MN. Hypohydration impairs endurance exercise performance in temperate but not cold air. *J Appl Physiol* (1985). 2005 Nov;99(5):1972–6.

Cheuvront SN, Haymes EM. Ad libitum fluid intakes and thermoregulatory responses of female distance runners in three environments. *J Sports Sci.* 2001 Nov;19(11):845–54.

Cheuvront SN, Carter R 3rd, Haymes EM, Sawka MN. No effect of moderate hypohydration or hyperthermia on anaerobic exercise performance. *Med Sci Sports Exerc.* 2006 Jun;38(6):1093–7.

Cheuvront SN, Kenefick RW, Montain SJ, Sawka MN. Mechanisms of aerobic performance impairment with heat stress and dehydration. *J Appl Physiol* (1985). 2010 Dec;109(6):1989–95.

Cheuvront SN, Kenefick RW, Ely BR, Harman EA, Castellani JW, Frykman PN, Nindl BC, Sawka MN. Hypohydration reduces vertical ground reaction impulse but not jump height. *Eur J Appl Physiol.* 2010 Aug;109(6):1163–70. (b)

Chinevere TD, Kenefick RW, Cheuvront SN, Lukaski HC, Sawka MN. Effect of heat acclimation on sweat minerals. *Med Sci Sports Exerc.* 2008 May;40(5):886–91.

Chlívková D, Knechtle B, Rosemann T, Zákovská A, Tomášková I, Shortall M, Tomášková I. Changes in foot volume, body composition, and hydration status in male and female 24-hour ultra-mountain bikers. *J Int Soc Sports Nutr.* 2014 Mar 24;11(1):12.

Clayton DJ, Evans GH, James LJ. Effect of drink carbohydrate content on postexercise gastric emptying, rehydration, and the calculation of net fluid balance. *Int J Sport Nutr Exerc Metab.* 2014 Feb;24(1):79–89.

Coles MG, Luetkemeier MJ. Sodium-facilitated hypervolemia, endurance performance, and thermoregulation. *Int J Sports Med.* 2005 Apr;26(3):182–7.

Cosgrove SD, Black KE. Sodium supplementation has no effect on endurance performance during a cycling time-trial in cool conditions: a randomised cross-over trial. *J Int Soc Sports Nutr.* 2013 Jun 3;10:30.

Cox GR, Broad EM, Riley MD, Burke LM. Body mass changes and voluntary fluid intakes of elite level water polo players and swimmers. *J Sci Med Sport.* 2002 Sep;5(3):183–93.

Cox GR, Clark SA, Cox AJ, et al. Daily training with high carbohydrate availability increases exogenous carbohydrate oxidation during endurance cycling. *J Appl Physiol.* 2010;109:126–134.

Coyle EF. Fluid and fuel intake during exercise. *J Sports Sci.* 2004 Jan;22(1):39–55. Review.

de Oliveira EP, Burini RC. The impact of physical exercise on the gastrointestinal tract. *Curr Opin Clin Nutr Metab Care.* 2009;12:533–538.

de Oliveira EP, Burini RC. Food-dependent, exercise-induced gastrointestinal distress. *J Int Soc Sports Nutr.* 2011;8:12

de Oliveira EP, Burini RC, Jeukendrup A. Gastrointestinal complaints during exercise: prevalence, etiology, and nutritional recommendations. *Sports Med.* 2014 May;44 Suppl 1:S79–85.

Devlin LH, Fraser SF, Barras NS, Hawley JA. Moderate levels of hypohydration impairs bowling accuracy but not bowling velocity in skilled cricket players. *J Sci Med Sport.* 2001 Jun;4(2):179–87.

Dion T, Savoie FA, Asselin A, Gariepy C, Goulet ED. Half-marathon running performance is not improved by a rate of fluid intake above that dictated by thirst sensation in trained distance runners. *Eur J Appl Physiol.* 2013 Dec;113(12):3011–20.

Dougherty KA, Baker LB, Chow M, Kenney WL. Two percent dehydration impairs and six percent carbohydrate drink improves boys basketball skills. *Med Sci Sports Exerc.* 2006 Sep;38(9):1650–8.

Duquette PP, Bissonnette P, Lapointe JY. Local osmotic gradients drive the water flux associated with Na(+) / glucose cotransport. *Proc Natl Acad Sci U S A.* 2001 Mar 27;98(7):3796–801.

Ely MR, Cheuvront SN, Roberts WO & Montain SJ. Impact of weather on marathon-running performance. *Med Sci Sports Exerc* 39, 487–493. (2007).

Ely BR, Ely MR, Cheuvront SN, Kenefick RW, Degroot DW, Montain SJ. Evidence against a 40 degrees C core temperature threshold for fatigue in humans. *J Appl Physiol* (1985). 2009 Nov;107(5):1519–25.

Ely BR, Cheuvront SN, Kenefick RW, Sawka MN. Aerobic performance is degraded, despite modest hyperthermia, in hot environments. *Med Sci Sports Exerc.* 2010 Jan;42(1):135–41.

Evans GH, Shirreffs SM, Maughan RJ. Postexercise rehydration in man: the effects of osmolality and carbohydrate content of ingested drinks. *Nutrition.* 2009 Sep;25(9):905–13.

Evans GH, Shirreffs SM, Maughan RJ. The effects of repeated ingestion of high and low glucose-electrolyte solutions on gastric emptying and blood 2H₂O concentration after an overnight fast. *Br J Nutr.* 2011 Dec;106(11):1732–9

Finravinto 2012 -tutkimus. Helldán A, Raulio S, Kosola M, Tapanainen H, Ovaskainen M-L, Virtanen S. THL Raportti 16/2013. Juvenes Print – Suomen Yliopistopaino Oy. Tampere 2013.

Fleming J, James LJ. Repeated familiarisation with hypohydration attenuates the performance decrement caused by hypohydration during treadmill running. *Appl Physiol Nutr Metab.* 2014 Feb;39(2):124–9.

Gabriel SE, Jaakkimainen L, Bombardier C. Risk for serious gastrointestinal complications related to use of nonsteroidal anti-inflammatory drugs. A meta-analysis. *Ann Intern Med.* 1991;115:787–796.

Galloway SD, Maughan RJ. Effects of ambient temperature on the capacity to perform prolonged cycle exercise in man. *Med Sci Sports Exerc.* 1997 Sep;29(9):1240–9.

Garrett AT, Goosens NG, Rehrer NJ, Patterson MJ, Harrison J, Sammut I, Cotter JD. Short-term heat acclimation is effective and may be enhanced rather than impaired by dehydration. *Am J Hum Biol.* 2014 May-Jun;26(3):311–20.

Garth AK, Burke LM. What do athletes drink during competitive sporting activities? *Sports Med.* 2013 Jul;43(7):539–64.

Gigou PY, Dion T, Asselin A, Berrigan F, Goulet ED. Pre-exercise hyperhydration-induced bodyweight gain does not alter prolonged treadmill running time-trial performance in warm ambient conditions. *Nutrients.* 2012 Aug;4(8):949–66.

Gisolfi CV, Summers RD, Schedl HP, Bleiler TL. Effect of sodium concentration in a carbohydrate-electrolyte solution on intestinal absorption. *Med Sci Sports Exerc.* 1995;27:1414–1420

Gisolfi CV, Lambert GP, Summers RW. Intestinal fluid absorption during exercise: role of sport drink osmolality and [Na⁺] *Med Sci Sports Exerc.* 2001;33:907–915.

Godek SF, Bartolozzi AR, Godek JJ. Sweat rate and fluid turnover in American football players compared with runners in a hot and humid environment. *Br J Sports Med.* 2005 Apr;39(4):205–11; discussion 205–11.

Godek SF, Peduzzi C, Burkholder R, Condon S, Dorshimer G, Bartolozzi AR. Sweat rates, sweat sodium concentrations, and sodium losses in 3 groups of professional football players. *J Athl Train.* 2010 Jul-Aug;45(4):364–71.

Gomes LH, Carneiro-Júnior MA, Marins JC. Thermoregulatory responses of children exercising in a hot environment. *Rev Paul Pediatr.* 2013 Jan-Mar;31(1):104–10.

González-Alonso J. Human thermoregulation and the cardiovascular system. *Exp Physiol.* 2012 Mar;97(3):340–6.

González-Alonso J, Calbet JA. Reductions in systemic and skeletal muscle blood flow and oxygen delivery limit maximal aerobic capacity in humans. *Circulation.* 2003 Feb 18;107(6):824–30.

Goulet ED. Effect of exercise-induced dehydration on time-trial exercise performance: a meta-analysis. *Br J Sports Med.* 2011 Nov;45(14):1149–56.

Goulet ED. Dehydration and endurance performance in competitive athletes. *Nutr Rev.* 2012 Nov;70 Suppl 2:S132–6.

Goulet ED. Effect of exercise-induced dehydration on endurance performance: evaluating the impact of exercise protocols on outcomes using a meta-analytic procedure. *Br J Sports Med.* 2013 Jul;47(11):679–86.

Hamouti N, Fernández-Elías VE, Ortega JF, Mora-Rodriguez R. Ingestion of sodium plus water improves cardiovascular function and performance during dehydrating cycling in the heat. *Scand J Med Sci Sports*. 2014 Jun;24(3):507–18.

Hew-Butler T, Collins M, Bosch A, Sharwood K, Wilson G, Armstrong M, Jennings C, Swart J, Noakes T. Maintenance of plasma volume and serum sodium concentration despite body weight loss in ironman triathletes. *Clin J Sport Med*. 2007 Mar;17(2):116–22.

Hew-Butler TD, Sharwood K, Collins M, Speedy D, Noakes T. Sodium supplementation is not required to maintain serum sodium concentrations during an Ironman triathlon. *Br J Sports Med*. 2006 Mar;40(3):255–9.

Hobson RM, Maughan RJ. Hydration status and the diuretic action of a small dose of alcohol. *Alcohol Alcohol*. 2010 Jul-Aug;45(4):366–73.

Hobson R, James L. The addition of whey protein to a carbohydrate-electrolyte drink does not influence post-exercise rehydration. *J Sports Sci*. 2014 Jul 17:1–8.

Hoffman MD, Hew-Butler T, Stuempfle KJ. Exercise-associated hyponatremia and hydration status in 161-km ultramarathoners. *Med Sci Sports Exerc*. 2013 Apr;45(4):784–91

Hosseini Lou A, Khamnei S, Zamanlu M. The effect of water temperature and voluntary drinking on the post rehydration sweating. *Int J Clin Exp Med*. 2013 Sep 1;6(8):683–7.

Houmard JA, Egan PC, Johns RA, Neufer PD, Chenier TC, Israel RG. Gastric emptying during 1 h of cycling and running at 75% VO_{2max}. *Med Sci Sports Exerc*. 1991 Mar;23(3):320–5.

Hunt JN, Smith JL, Jiang CL. Effect of meal volume and energy density on the gastric emptying of carbohydrates. *Gastroenterology*. 1985 Dec;89(6):1326–30.

Ihsan M, Landers G, Brearley M, Peeling P. Beneficial effects of ice ingestion as a precooling strategy on 40-km cycling time-trial performance. *Int J Sports Physiol Perform*. 2010;5:140–151.

International Marathon Medical Directors Association, Noakes T; IMMAD. Fluid replacement during marathon running. *Clin J Sport Med*. 2003 Sep;13(5):309–18.

IOC. Nutrition Working Group of the International Olympic Committee. Nutrition for Athletes. A practical guide to eating for health and performance. Revised and updated 2012.

Ismail I, Singh R, Sirisinghe RG. Rehydration with sodium-enriched coconut water after exercise-induced dehydration. *Southeast Asian J Trop Med Public Health*. 2007;13:769–785

James LJ, Gingell R, Evans GH. Whey protein addition to a carbohydrate-electrolyte rehydration solution ingested after exercise in the heat. *J Athl Train*. 2012 Jan-Feb;47(1):61–6.

James LJ, Evans GH, Madin J, Scott D, Stepney M, Harris R, Stone R, Clayton DJ. Effect of varying the concentrations of carbohydrate and milk protein in rehydration solutions ingested after exercise in the heat. *Br J Nutr*. 2013 Oct;110(7):1285–91.

James LJ, Mattin L, Aldiss P, Adebishi R, Hobson RM. Effect of whey protein isolate on rehydration after exercise. *Amino Acids*. 2014 May;46(5):1217–24.

Jéquier E, Constant F. Water as an essential nutrient: the physiological basis of hydration. *Eur J Clin Nutr*. 2010 Feb;64(2):115–23.

Jeukendrup AE, Vet-Joop K, Sturk A, et al. Relationship between gastro-intestinal complaints and endotoxaemia, cytokine release and the acute-phase reaction during and after a long-distance triathlon in highly trained men. *Clin Sci (Lond)* 2000;98:47–55.

Jeukendrup AE, Currell K, Clarke J, Cole J, Blannin AK. Effect of beverage glucose and sodium content on fluid delivery. *Nutr Metab (Lond)*. 2009 Feb 20;6:9.

Jeukendrup AE, Moseley L. Multiple transportable carbohydrates enhance gastric emptying and fluid delivery. *Scand J Med Sci Sports*. 2010 Feb;20(1):112–21

Jeukendrup AE, McLaughlin J. Carbohydrate ingestion during exercise: effects on performance, training adaptations and trainability of the gut. *Nestle Nutr Inst Workshop Ser*. 2011;69:1–12; discussion 13–7.

Jones EJ, Bishop PA, Green JM, Richardson MT. Effects of metered versus bolus water consumption on urine production and rehydration. *Int J Sport Nutr Exerc Metab.* 2010 Apr;20(2):139–44.

Jones PR, Barton C, Morrissey D, Maffulli N, Hemmings S. Pre-cooling for endurance exercise performance in the heat: a systematic review. *BMC Med.* 2012 Dec 18;10:166.

Judelson DA, Maresh CM, Anderson JM, Armstrong LE, Casa DJ, Kraemer WJ, Volek JS. Hydration and muscular performance: does fluid balance affect strength, power and high-intensity endurance? *Sports Med.* 2007;37(10):907–21. Review. (a)

Judelson DA, Maresh CM, Farrell MJ, Yamamoto LM, Armstrong LE, Kraemer WJ, Volek JS, Spiering BA, Casa DJ, Anderson JM. Effect of hydration state on strength, power, and resistance exercise performance. *Med Sci Sports Exerc.* 2007 Oct;39(10):1817–24. (b)

Judelson DA, Maresh CM, Yamamoto LM, Farrell MJ, Armstrong LE, Kraemer WJ, Volek JS, Spiering BA, Casa DJ, Anderson JM. Effect of hydration state on resistance exercise-induced endocrine markers of anabolism, catabolism, and metabolism. *J Appl Physiol* (1985). 2008 Sep;105(3):816–24.

Kao WF, Shyu CL, Yang XW, Hsu TF, Chen JJ, Kao WC, Polun-Chang, Huang YJ, Kuo FC, Huang CI, Lee CH. Athletic performance and serial weight changes during 12- and 24-hour ultra-marathons. *Clin J Sport Med.* 2008 Mar;18(2):155–8.

Khamnei S, Hosseinlou A, Zamanlu M. Water temperature, voluntary drinking and fluid balance in dehydrated taekwondo athletes. *J Sports Sci Med.* 2011 Dec 1;10(4):718–24.

Kipps C, Sharma S, Pedoe DT. The incidence of exercise-associated hyponatraemia in the london marathon. *Br J Sports Med.* 2011;10:14

Knechtle B, Knechtle P, Wirth A, Alexander Rüst C, Rosemann T. A faster running speed is associated with a greater body weight loss in 100-km ultra-marathoners. *J Sports Sci.* 2012;30(11):1131–40

Koehler K, Thevis M, Schaenzer W. Meta-analysis: Effects of glycerol administration on plasma volume, haemoglobin, and haematocrit. *Drug Test Anal.* 2013 Nov-Dec;5(11–12):896–9. doi: 10.1002/dta.1580.

Kovacs EM, Schmahl RM, Senden JM, Brouns F. Effect of high and low rates of fluid intake on post-exercise rehydration. *Int J Sport Nutr Exerc Metab.* 2002 Mar;12(1):14–23.

Lambert GP, Chang RT, Xia T, Summers RW, Gisolfi CV. Absorption from different intestinal segments during exercise. *J Appl Physiol* (1985). 1997 Jul;83(1):204–12.

Lambert GP, Lang J, Bull A, Eckerson J, Lanspa S, O'Brien J. Fluid tolerance while running: effect of repeated trials. *Int J Sports Med.* 2008 Nov;29(11):878–82.

Latzka WA, Sawka MN. Hyperhydration and glycerol: thermoregulatory effects during exercise in hot climates. *Can J Appl Physiol.* 2000 Dec;25(6):536–45. Review.

Lee JK, Shirreffs SM. The influence of drink temperature on thermoregulatory responses during prolonged exercise in a moderate environment. *J Sports Sci.* 2007 Jul;25(9):975–85.

Lee JK, Shirreffs SM, Maughan RJ. Cold drink ingestion improves exercise endurance capacity in the heat. *Med Sci Sports Exerc.* 2008 Sep;40(9):1637–44.

Lee JK, Nio AQ, Lim CL, Teo EY, Byrne C. Thermoregulation, pacing and fluid balance during mass participation distance running in a warm and humid environment. *Eur J Appl Physiol.* 2010 Jul;109(5):887–98.

Lee JB, Kim TW, Min YK, Yang HM. Long distance runners present upregulated sweating responses than sedentary counterparts. *PLoS One.* 2014 Apr 7;9(4):e93976.

Lee Jones B, O'Hara JP, Till K, Gerardus Joseph King RF. Dehydration and Hyponatremia in Professional Rugby Union Players; A Cohort Study observing English Premiership Rugby Union Players during Match Play, Field and Gym Training in Cool Environmental Conditions. *J Strength Cond Res.* 2014 Jul 15.

Leiper JB, Broad NP, Maughan RJ. Effect of intermittent high-intensity exercise on gastric emptying in man. *Med Sci Sports Exerc.* 2001 Aug;33(8):1270–8.

Levine L, Rose MS, Francesconi P, Neufer PD, Sawka MN. Fluid replacement during sustained activity in the heat: nutrient solution vs. water. *Aviat Space Environ Med.* 1991 Jun;62(6):559–64.

Lopez RM, Casa DJ, Jensen KA, DeMartini JK, Pagnotta KD, Ruiz RC, Roti MW, Stearns RL, Armstrong LE, Maresh CM. Examining the influence of hydration status on physiological responses and running speed during trail running in the heat with controlled exercise intensity. *J Strength Cond Res.* 2011 Nov;25(11):2944–54.

Luetkemeier MJ, Coles MG, Askew EW. Dietary sodium and plasma volume levels with exercise. *Sports Med.* 1997 May;23(5):279–86. Review.

MacLeod H, Sunderland C. Previous-day hypohydration impairs skill performance in elite female field hockey players. *Scand J Med Sci Sports.* 2012 Jun;22(3):430–8.

Maresh CM, Whittlesey MJ, Armstrong LE, Yamamoto LM, Judelson DA, Fish KE, Casa DJ, Kavouras SA, Castracane VD. Effect of hydration state on testosterone and cortisol responses to training-intensity exercise in collegiate runners. *Int J Sports Med.* 2006 Oct;27(10):765–70.

Marino F. Evidence for anticipatory regulation mediated by drink temperature during fixed intensity exercise in the heat. *Exp Physiol.* 2007 Mar;92(2):467–8; author reply 469

Marino FE, Mbambo Z, Kortekaas E, Wilson G, Lambert MI, Noakes TD, Dennis SC. Advantages of smaller body mass during distance running in warm, humid environments. *Pflugers Arch.* 2000 Dec;441(2–3):359–67.

Martens M.P., Watson J.C., Royland E.M., Beck N.C. Development of the athlete drinking scale. *Psychol. Addict. Behav.* 2005;19:158–164

Maughan RJ. Impact of mild dehydration on wellness and on exercise performance. *Eur J Clin Nutr.* 2003 Dec;57 Suppl 2:S19–23.

Maughan RJ, Owen JH, Shirreffs SM, Leiper JB. Post-exercise rehydration in man: effects of electrolyte addition to ingested fluids. *Eur J Appl Physiol Occup Physiol.* 1994;13:209–215

Maughan RJ, Leiper JB. Sodium intake and post-exercise rehydration in man. *Eur J Appl Physiol Occup Physiol.* 1995;71(4):311–9.

Maughan RJ, Leiper JB, Shirreffs SM. Restoration of fluid balance after exercise-induced dehydration: effects of food and fluid intake. *Eur J Appl Physiol Occup Physiol.* 1996;73(3–4):317–25. (b)

Maughan RJ, Leiper JB. Limitations to fluid replacement during exercise. *Can J Appl Physiol.* 1999 Apr;24(2):173–87. Review.

Maughan RJ, Griffin J. Caffeine ingestion and fluid balance: a review. *J Hum Nutr Diet.* 2003 Dec;16(6):411–20.

Maughan RJ, Shirreffs SM, Merson SJ, Horswill CA. Fluid and electrolyte balance in elite male football (soccer) players training in a cool environment. *J Sports Sci.* 2005 Jan;23(1):73–9.

Maughan RJ, Watson P, Evans GH, Broad N, Shirreffs SM. Water balance and salt losses in competitive football. *Int J Sport Nutr Exerc Metab.* 2007 Dec;17(6):583–94.

Maughan RJ, Dargavel LA, Hares R, Shirreffs SM. Water and salt balance of well-trained swimmers in training. *Int J Sport Nutr Exerc Metab.* 2009 Dec;19(6):598–606.

Maughan RJ, Otani H, Watson P. Influence of relative humidity on prolonged exercise capacity in a warm environment. *Eur J Appl Physiol.* 2012 Jun;112(6):2313–21.

Maughan RJ, Meyer NL. Hydration during intense exercise training. *Nestle Nutr Inst Workshop Ser.* 2013;76:25–37.

McGregor SJ, Nicholas CW, Lakomy HK, Williams C. The influence of intermittent high-intensity shuttle running and fluid ingestion on the performance of a soccer skill. *J Sports Sci.* 1999 Nov;17(11):895–903.

Mears SA, Shirreffs SM. The effects of high-intensity intermittent exercise compared with continuous exercise on voluntary water ingestion. *Int J Sport Nutr Exerc Metab.* 2013 Oct;23(5):488–97.

Mears SA, Shirreffs SM. Voluntary water intake during and following moderate exercise in the cold. *Int J Sport Nutr Exerc Metab.* 2014 Feb;24(1):47–58.

Merry TL, Ainslie PN, Walker R, Cotter JD. Fitness alters fluid regulatory but not behavioural responses to hypohydrated exercise. *Physiol Behav.* 2008 Oct 20;95(3):348–52.

Merry TL, Ainslie PN, Cotter JD. Effects of aerobic fitness on hypohydration-induced physiological strain and exercise impairment. *Acta Physiol (Oxf)*. 2010 Feb;198(2):179–90.

Meyer N. Nutrition for winter sports. IOC Consensus In Sports Nutrition Conference Lausanne, Switzerland 25th–27th October 2010

Minehan MR, Riley MD, Burke LM. Effect of flavor and awareness of kilojoule content of drinks on preference and fluid balance in team sports. *Int J Sport Nutr Exerc Metab*. 2002;12:81–92

Mitchell JB, Voss KW. The influence of volume on gastric emptying and fluid balance during prolonged exercise. *Med Sci Sports Exerc*. 1991 Mar;23(3):314–9. Review.

Mitchell JB, Grandjean PW, Pizza FX, Starling RD, Holtz RW. The effect of volume ingested on rehydration and gastric emptying following exercise-induced dehydration. *Med Sci Sports Exerc*. 1994 Sep;26(9):1135–43.

Moran DS, Montain SJ, Pandolf KB. Evaluation of different levels of hydration using a new physiological strain index. *Am J Physiol*. 1998 Sep;275(3 Pt 2):R854–60.

Mora-Rodriguez R, Hamouti N. Salt and fluid loading: effects on blood volume and exercise performance. *Med Sport Sci*. 2012;59:113–9.

Murray R. The effects of consuming carbohydrate-electrolyte beverages on gastric emptying and fluid absorption during and following exercise. *Sports Med*. 1987;4:322–351.

Mündel T, King J, Collacott E, Jones DA. Drink temperature influences fluid intake and endurance capacity in men during exercise in a hot, dry environment. *Exp Physiol*. 2006 Sep;91(5):925–33. Epub 2006 Jun 15.

Neufer PD, Young AJ, Sawka MN. Gastric emptying during exercise: effects of heat stress and hypohydration. *Eur J Appl Physiol Occup Physiol*. 1989;58:433–439.

Nielsen B. Heat acclimation--mechanisms of adaptation to exercise in the heat. *Int J Sports Med*. 1998 Jun;19 Suppl 2:S154–6.

Noakes TD. Fluid and electrolyte disturbances in heat illness. *Int J Sports Med*. 1998 Jun;19 Suppl 2:S146–9.

Noakes TD. Is drinking to thirst optimum? *Ann Nutr Metab*. 2010;57 Suppl 2:9–17.

Noakes TD. Time to move beyond a brainless exercise physiology: the evidence for complex regulation of human exercise performance. *Appl Physiol Nutr Metab*. 2011 Feb;36(1):23–35.

Noakes TD. Fatigue is a Brain-Derived Emotion that Regulates the Exercise Behavior to Ensure the Protection of Whole Body Homeostasis. *Front Physiol*. 2012 Apr 11;3:82.

Noakes TD, Sharwood K, Collins M, Perkins DR. The dipsomania of great distance: water intoxication in an Ironman triathlete. *Br J Sports Med*. 2004 Aug;38(4):E16.

Noakes TD, Sharwood K, Speedy D, Hew T, Reid S, Dugas J, Almond C, Wharam P, Weschler L. Three independent biological mechanisms cause exercise-associated hyponatremia: evidence from 2,135 weighed competitive athletic performances. *Proc Natl Acad Sci U S A*. 2005 Dec 20;102(51):18550–5.

Naharudin MN, Yusof A. Fatigue index and fatigue rate during an anaerobic performance under hypohydration. *PLoS One*. 2013 Oct 30;8(10):e77290.

Nybo L. Exercise and heat stress: cerebral challenges and consequences. *Prog Brain Res*. 2007;162:29–43. Review.

O'Brien C, Young AJ, Sawka MN. Bioelectrical impedance to estimate changes in hydration status. *Int J Sports Med*. 2002 Jul;23(5):361–6.

O'Brien WJ, Rowlands DS. Fructose-maltodextrin ratio in a carbohydrate-electrolyte solution differentially affects exogenous carbohydrate oxidation rate, gut comfort, and performance. *Am J Physiol Gastrointest Liver Physiol*. 2011 Jan;300(1):G181–9.

Osterberg KL, Horswill CA, Baker LB. Pregame urine specific gravity and fluid intake by National Basketball Association players during competition. *J Athl Train*. 2009 Jan-Feb;44(1):53–7.

Osterberg KL, Pallardy SE, Johnson RJ, Horswill CA. Carbohydrate exerts a mild influence on fluid retention following exercise-induced dehydration. *J Appl Physiol* (1985). 2010 Feb;108(2):245–50.

Owen JA, Kehoe SJ, Oliver SJ. Influence of fluid intake on soccer performance in a temperate environment. *J Sports Sci.* 2013;31(1):1–10.

Palmer MS, Spriet LL. Sweat rate, salt loss, and fluid intake during an intense on-ice practice in elite Canadian male junior hockey players. *Appl Physiol Nutr Metab.* 2008 Apr;33(2):263–71.

Palmer MS, Logan HM, Spriet LL. On-ice sweat rate, voluntary fluid intake, and sodium balance during practice in male junior ice hockey players drinking water or a carbohydrate-electrolyte solution. *Appl Physiol Nutr Metab.* 2010 Jun;35(3):328–35.

Peacock OJ, Stokes K, Thompson D. Initial hydration status, fluid balance, and psychological affect during recreational exercise in adults. *J Sports Sci.* 2011 Jun;29(9):897–904.

Peters HP, Bos M, Seebregts L, et al. Gastrointestinal symptoms in long-distance runners, cyclists, and triathletes: prevalence, medication, and etiology. *Am J Gastroenterol.* 1999;94:1570–81

Peters HP, Wiersma JW, Koerselman J, et al. The effect of a sports drink on gastroesophageal reflux during a run-bike-run test. *Int J Sports Med.* 2000;21:65–70

Pérez-Idárraga A, Aragón-Vargas LF. Postexercise rehydration: potassium-rich drinks versus water and a sports drink. *Appl Physiol Nutr Metab.* 2014 May 9:1–8.

Pfeiffer B, Cotterill A, Grathwohl D, Stellingwerff T, Jeukendrup AE. The effect of carbohydrate gels on gastrointestinal tolerance during a 16-km run. *Int J Sport Nutr Exerc Metab.* 2009 Oct;19(5):485–503.

Pfeiffer B, Stellingwerff T, Zaltas E, Jeukendrup AE. CHO oxidation from a CHO gel compared with a drink during exercise. *Med Sci Sports Exerc.* 2010 Nov;42(11):2038–45.5(11):1211–24. (a)

Pfeiffer B, Stellingwerff T, Zaltas E, Jeukendrup AE. Oxidation of solid versus liquid CHO sources during exercise. *Med Sci Sports Exerc.* 2010 Nov;42(11):2030–7.

Pfeiffer B, Stellingwerff T, Hodgson AB, Randell R, Pöttgen K, Res P, Jeukendrup AE. Nutritional intake and gastrointestinal problems during competitive endurance events. *Med Sci Sports Exerc.* 2012 Feb;44(2):344–51.

Quiterio AL, Silva AM, Minderico CS, Carnero EA, Fields DA, Sardinha LB. Total body water measurements in adolescent athletes: a comparison of six field methods with deuterium dilution. *J Strength Cond Res.* 2009 Jul;23(4):1225–37

Rehrer NJ, Beckers EJ, Brouns F, et al. Effects of dehydration on gastric emptying and gastrointestinal distress while running. *Med Sci Sports Exerc.* 1990;22:790–795.

Rehrer NJ. The maintenance of fluid balance during exercise. *Int J Sports Med.* 1994 Apr;15(3):122–5.

Ross ML, Garvican LA, Jeacocke NA, Laursen PB, Abbiss CR, Martin DT, Burke LM. Novel precooling strategy enhances time trial cycling in the heat. *Med Sci Sports Exerc.* 2011;43:123–133.

Rowland T. Fluid replacement requirements for child athletes. *Sports Med.* 2011 Apr 1;41(4):279–88.

Rudzki SJ, Hazard H, Collinson D. Gastrointestinal blood loss in triathletes: its etiology and relationship to sports anaemia. *Aust J Sci Med Sport.* 1995;27:3–8

Rüst CA, Knechtle B, Knechtle P, Wirth A, Rosemann T. Body mass change and ultraendurance performance: a decrease in body mass is associated with an increased running speed in male 100-km ultramarathoners. *J Strength Cond Res.* 2012 Jun;26(6):1505–16

Samborski P, Chmielarz-Czarnoci ska A, Grzymisławski M. Exercise-induced vomiting. *Prz Gastroenterol.* 2013;8(6):396–400

Saunders P.U., Pyne D.B., Telford R.D., Hawley J.A. Factors affecting running economy in trained distance runners. *Sports Med.* 2004;34:465–485

Sawka MN, Montain SJ, Latzka WA. Hydration effects on thermoregulation and performance in the heat. *Comp Biochem Physiol A Mol Integr Physiol.* 2001;128:679–690

Sawka MN, Cheuvront SN, Carter R 3rd. Human water needs. *Nutr Rev.* 2005 Jun;63(6 Pt 2):S30–9.

Sawka MN, Cheuvront SN, Kenefick RW. High skin temperature and hypohydration impair aerobic performance. *Exp Physiol.* 2012 Mar;97(3):327–32.

Seifert J, Harmon J, DeClercq P. Protein added to a sports drink improves fluid retention. *Int J Sport Nutr Exerc Metab.* 2006;13:420–429.

Sharp RL. Role of whole foods in promoting hydration after exercise in humans. *J Am Coll Nutr.* 2007 Oct;26(5 Suppl):592S-596S. Review.

Sharwood K, Collins M, Goedecke J, Wilson G, Noakes T. Weight changes, sodium levels, and performance in the South African Ironman Triathlon. *Clin J Sport Med.* 2002 Nov;12(6):391–9.

Sharwood KA, Collins M, Goedecke JH, Wilson G, Noakes TD. Weight changes, medical complications, and performance during an Ironman triathlon. *Br J Sports Med.* 2004 Dec;38(6):718–24.

Shephard RJ. Is it time to retire the ‘central governor’? *Sports Med.* 2009;39(9):709–21.

Shi X, Gisolfi CV. Fluid and carbohydrate replacement during intermittent exercise. *Sports Med.* 1998 Mar;25(3):157–72.

Shirreffs SM, Taylor AJ, Leiper JB, Maughan RJ. Post-exercise rehydration in man: effects of volume consumed and drink sodium content. *Med Sci Sports Exerc.* 1996 Oct;28(10):1260–71.

Shirreffs SM, Maughan RJ. Restoration of fluid balance after exercise-induced dehydration: effects of alcohol consumption. *J Appl Physiol (1985).* 1997 Oct;83(4):1152–8.

Shirreffs SM, Maughan RJ. The effect of alcohol on athletic performance. *Curr Sports Med Rep.* 2006 Jun;5(4):192–6.

Shirreffs SM. Fluid and electrolyte needs for training, competition and recovery. IOC Consensus In Sports Nutrition Conference Lausanne, Switzerland 25th–27th October 2010

Shirreffs SM, Aragon-Vargas LF, Chamorro M, Maughan RJ, Serratosa L, Zachwieja JJ. The sweating response of elite professional soccer players to training in the heat. *Int J Sports Med.* 2005 Mar;26(2):90–5.

Shirreffs SM, Aragon-Vargas LF, Keil M, Love TD, Phillips S. Rehydration after exercise in the heat: a comparison of 4 commonly used drinks. *Int J Sport Nutr Exerc Metab.* 2007 Jun;17(3):244–58. (a)

Shirreffs SM, Watson P, Maughan RJ. Milk as an effective post-exercise rehydration drink. *Br J Nutr.* 2007 Jul;98(1):173–80. (b)

Shirreffs SM, Sawka MN. Fluid and electrolyte needs for training, competition, and recovery. *J Sports Sci.* 2011;29 Suppl 1:S39–46.

Siegel R, Maté J, Brearley MB, Watson G, Nosaka K, Laursen PB. Ice slurry ingestion increases core temperature capacity and running time in the heat. *Med Sci Sports Exerc.* 2010;42:717–725.

Siegel R, Laursen PB. Keeping your cool: possible mechanisms for enhanced exercise performance in the heat with internal cooling methods. *Sports Med.* 2012 Feb 1;42(2):89–98.

Siegel R, Maté J, Watson G, Nosaka K, Laursen PB. Pre-cooling with ice slurry ingestion leads to similar run times to exhaustion in the heat as cold water immersion. *J Sports Sci.* 2012;30:155–165.

Silva AM, Júdice PB, Matias CN, Santos DA, Magalhães JP, St-Onge MP, Gonçalves EM, Armada-da-Silva P, Sardinha LB. Total body water and its compartments are not affected by ingesting a moderate dose of caffeine in healthy young adult males. *Appl Physiol Nutr Metab.* 2013 Jun;38(6):626–32.

Simrén M. Physical activity and the gastrointestinal tract. *Eur J Gastroenterol Hepatol.* 2002;14:1053–6

Sims ST, Rehrer NJ, Bell ML, Cotter JD. Preexercise sodium loading aids fluid balance and endurance for women exercising in the heat. *J Appl Physiol.* 2007;105:534–541. (a)

Sims ST, van Vliet L, Cotter JD, Rehrer NJ. Sodium loading aids fluid balance and reduces physiological strain of trained men exercising in the heat. *Medicine and Science in Sports and Exercise.* 2007;10:123–130. (b)

Speedy DB, Thompson J, Rodgers I, Collins M, Sharwood K. Oral salt supplementation during ultradistance exercise. *Clin J Sport Med.* 2002;10:279.

Stachenfeld NS. Acute effects of sodium ingestion on thirst and cardiovascular function. *Curr Sports Med Rep.* 2008 Jul-Aug;7(4 Suppl):S7–13.

Stachenfeld NS. The interrelationship of research in the laboratory and the field to assess hydration status and determine mechanisms involved in water regulation during physical activity. *Sports Med.* 2014 May;44 Suppl 1:S97–104.

Stellingwerff T. Contemporary nutrition approaches to optimize elite marathon performance. *Int J Sports Physiol Perform.* 2013 Sep;8(5):573–8. Epub 2013 Apr 9. Review.

Stellingwerff T, Jeukendrup AE. Don't forget the gut--it is an important athletic organ! *J Appl Physiol (1985).* 2011 Jan;110(1):278; discussion 294.

Stuempfle KJ, Hoffman MD, Hew-Butler T. Gastrointestinal distress in ultramarathoners is associated with race diet. *Int J Sport Nutr Exerc Metab.* 2013;23:103–109

Suzuki K, Hashimoto H, Oh T, Ishijima T, Mitsuda H, Peake JM, Sakamoto S, Muraoka I, Higuchi M. The effects of sports drink osmolality on fluid intake and immunoendocrine responses to cycling in hot conditions. *J Nutr Sci Vitaminol (Tokyo).* 2013;59(3):206–12.

Tai CY, Joy JM, Falcone PH, Carson LR, Mosman MM, Straight JL, Oury SL, Mendez C Jr, Loveridge NJ, Kim MP, Moon JR. An amino acid-electrolyte beverage may increase cellular rehydration relative to carbohydrate-electrolyte and flavored water beverages. *Nutr J.* 2014 May 26;13:47.

Tam N, Nolte HW, Noakes TD. Changes in total body water content during running races of 21.1 km and 56 km in athletes drinking ad libitum. *Clin J Sport Med.* 2011 May;21(3):218–25.

ter Steege RW, Van der Palen J, Kolkman JJ. Prevalence of gastrointestinal complaints in runners competing in a long-distance run: an internet-based observational study in 1281 subjects. *Scand J Gastroenterol.* 2008;43:1477–1482.

Twerenbold R, Knechtle B, Kakebeeke TH, Eser P, Müller G, von Arx P, Knecht H. Effects of different sodium concentrations in replacement fluids during prolonged exercise in women. *Br J Sports Med.* 2003 Aug;37(4):300–3; discussion 303.

van Rosendal SP, Osborne MA, Fassett RG, Coombes JS. Guidelines for glycerol use in hyperhydration and rehydration associated with exercise. *Sports Med.* 2010 Feb 1;40(2):113–29.

van Rosendal SP, Coombes JS. Glycerol use in hyperhydration and rehydration: scientific update. *Med Sport Sci.* 2012;59:104–12.

van Wijck K, Lenaerts K, Grootjans J, et al. Physiology and pathophysiology of splanchnic hypoperfusion and intestinal injury during exercise: strategies for evaluation and prevention. *Am J Physiol.* 2012;303:G155–G168 a)

van Wijck K, Lenaerts K, van Bijnen AA, et al. Aggravation of exercise-induced intestinal injury by Ibuprofen in athletes. *Med Sci Sports Exerc.* 2012;44:2257–2262. (b)

Vella LD, Cameron-Smith D. Alcohol, athletic performance and recovery. *Nutrients.* 2010 Aug;2(8):781–9.

Vrijens DM, Rehrer NJ. Sodium-free fluid ingestion decreases plasma sodium during exercise in the heat. *J Appl Physiol (1985).* 1999 Jun;86(6):1847–51.

VRN. Terveytä ruuasta – Suomalaiset ravitsemussuositukset 2014. Juvenes Oy, Helsinki 2014.

Wall BA, Watson G, Peiffer JJ, Abbiss CR, Siegel R, Laursen PB. Current hydration guidelines are erroneous: dehydration does not impair exercise performance in the heat. *Br J Sports Med.* 2013 Sep 20.

Watson P, Love TD, Maughan RJ, Shirreffs SM. A comparison of the effects of milk and a carbohydrate-electrolyte drink on the restoration of fluid balance and exercise capacity in a hot, humid environment. *Eur J Appl Physiol.* 2008 Nov;104(4):633–42.

Wegmann M, Faude O, Poppendieck W, Hecksteden A, Fröhlich M, Meyer T. Pre-cooling and sports performance: a meta-analytical review. *Sports Med.* 2012 Jul 1;42(7):545–64

Wright H, Collins M, Villiers RD, et al. Are splanchnic hemodynamics related to the development of gastrointestinal symptoms in ironman triathletes? A prospective cohort study. *Clin J Sport Med.* 2011;21:337–343.

Zouhal H, Groussard C, Minter G, Vincent S, Cretual A, Gratas-Delamarche A, Delamarche P, Noakes TD. Inverse relationship between percentage body weight change and finishing time in 643 forty-two-kilometre marathon runners. *Br J Sports Med.* 2011 Nov;45(14):1101–5.

Lähdeluettelo luku 9

- Affinito P, Tommaselli GA, DiCarlo C, Guida F, Nappi C.** Changes in bone mineral density and calcium metabolism in breastfeeding women: A one year follow-up study. *J Clin Endocrinol Metab* 1996;81:2314–8.
- Aissa Benhaddad A, Bouix D, Khaled S, Micallef JP, Mercier J, Bringer J, Brun JF.** Early hemorheologic aspects of overtraining in elite athletes. *Clin Hemorheol Microcirc.* 1999;20(2):117–25.
- Alessio HM.** Exercise-induced oxidative stress. *Med Sci Sports Exerc.* 1993 Feb;25(2):218–24.
- Alissa EM, Bahijri SM, Ferns GA.** The controversy surrounding selenium and cardiovascular disease: a review of the evidence. *Med Sci Monit.* 2003 Jan;9(1):RA9–18
- Anderson JB.** The important role of physical activity in skeletal development: how exercise may counter low calcium intake. *Am J Clin Nutr* 2000;71:1384–6.
- Anderson RA, Polansky MM, Bryden NA.** Stability and absorption of chromium and absorption of chromium histidinate complexes by humans. *Biol Trace Elem Res.* 2004 Dec;101(3):211–8.
- Argiratos V, Samman S.** The effect of calcium carbonate and calcium citrate on the absorption of zinc in healthy female subjects. *Eur J Clin Nutr.* 1994 Mar;48(3):198–204.
- Aruoma OI, Reilly T, MacLaren D, Halliwell B.** Iron, copper and zinc concentrations in human sweat and plasma; the effect of exercise. *Clin Chim Acta.* 1988 Sep 30;177(1):81–7.
- Aruoma OI.** Nutrition and health aspects of free radicals and antioxidants. *Food Chem Toxicol.* 1994 Jul;32(7):671–83.
- Ashe JR, Schofield FA, Gram MR.** The retention of calcium, iron, phosphorus, and magnesium during pregnancy; the adequacy of prenatal diets with and without supplementation. *Am J Clin Nutr* 1979;32:286–91.
- Atalay M, Laaksonen DE, Khanna S, Kaliste-Korhonen E, Hanninen O, Sen CK.** Vitamin E regulates changes in tissue antioxidants induced by fish oil and acute exercise. *Med Sci Sports Exerc.* 2000 Mar;32(3):601–7.
- Bahadori B, Wallner S, Schneider H, Wascher TC, Toplak H.** Effect of chromium yeast and chromium picolinate on body composition of obese, non-diabetic patients during and after a formula diet. *Acta Med Austriaca.* 1997;24(5):185–7.
- Bailey DM, Davies, B.** Acute mountain sickness; prophylactic benefits of antioxidant vitamin supplementation at high altitude. *High Alt Med Biol.* 2001 Spring;2(1):21–9.
- Baker DH.** Cupric oxide should not be used as a copper supplement for either animals or humans. *J Nutr.* 1999 Dec;129(12):2278–9.
- Baker LB, Stofan JR, Lukaski HC, Horswill CA.** Exercise-induced trace mineral element concentration in regional versus whole-body wash-down sweat. *Int J Sport Nutr Exerc Metab.* 2011 Jun;21(3):233–9.
- Bath SC, Button S, Rayman MP.** Iodine concentration of organic and conventional milk: implications for iodine intake. *Br J Nutr.* 2012 Apr;107(7):935–40.
- Bath SC, Rayman MP.** Iodine deficiency in the U.K.: an overlooked cause of impaired neurodevelopment? *Proc Nutr Soc.* 2013 May;72(2):226–35.
- Bazzarre TL, Scarpino A, Sigmon R, Marquart LF, Wu SM, Izurieta M.** Vitamin-mineral supplement use and nutritional status of athletes. *J Am Coll Nutr.* 1993 Apr;12(2):162–9.
- Beals KA.** Eating behaviors, nutritional status, and menstrual function in elite female adolescent volleyball players. *J Am Diet Assoc.* 2002 Sep;102(9):1293–6.
- Beard J, Tobin B.** Iron status and exercise. *Am J Clin Nutr.* 2000 Aug;72(2 Suppl):594S–7S
- Beitz R, Mensink GB, Henschel Y, Fischer B, Erbersdobler HF.** Dietary behaviour of German adults differing in levels of sport activity. *Public Health Nutr.* 2004 Feb;7(1):45–52.
- Beltz SD, Doering PL.** Efficacy of nutritional supplements used by athletes. *Clin Pharm.* 1993 Dec;12(12):900–8.
- Bergström L.** Nutrient losses and gains in the preparation of foods. *Uppsala: Livsmedelsverket,* 1994;32

Berkelhammer C, Bear RA. A clinical approach to common electrolyte problems: 4. Hypomagnesemia. *Can Med Assoc J.* 1985 Feb 15;132(4):360–8.

Bi L, Triadafilopoulos G. Exercise and gastrointestinal function and disease: an evidence-based review of risks and benefits. *Clin Gastroenterol Hepatol.* 2003 Sep;1(5):345–55.

Bjelakovic G, Nikolova D, Gluud C. Meta-regression analyses, meta-analyses, and trial sequential analyses of the effects of supplementation with beta-carotene, vitamin A, and vitamin E singly or in different combinations on all-cause mortality: do we have evidence for lack of harm? *PLoS One.* 2013 Sep 6;8(9):e74558.

Bjelakovic G, Nikolova D, Gluud C. Antioxidant supplements and mortality. *Curr Opin Clin Nutr Metab Care.* 2014 Jan;17(1):40–4.

Bielinski RW. Magnesium and exercise. *Rev Med Suisse.* 2006 Jul 26;2(74):1783–6

Black PN, Scragg R. Relationship between serum 25-hydroxyvitamin d and pulmonary function in the third national health and nutrition examination survey. *Chest.* 2005 Dec;128(6):3792–8.

Bloomer RJ, Goldfarb AH, McKenzie MJ, You T, Nguyen L. Effects of antioxidant therapy in women exposed to eccentric exercise. *Int J Sport Nutr Exerc Metab.* 2004 Aug;14(4):377–88.

Blumberg J, Block G. The Alpha-Tocopherol, Beta-Carotene Cancer Prevention Study in Finland. *Nutr Rev.* 1994 Jul;52(7):242–5.

Bonke D, Nickel B. Improvement of fine motoric movement control by elevated dosages of vitamin B1, B6, and B12 in target shooting. *Int J Vitam Nutr Res Suppl.* 1989;30:198–204

Borghi L, Schianchi T, Meschi T, Guerra A, Allegri F, Maggiore U, Novarini A. Comparison of two diets for the prevention of recurrent stones in idiopathic hypercalciuria. *New Engl J Med* 2002;346:77–84.

Braakhuis AJ, Hopkins WG, Lowe TE. Effects of dietary antioxidants on training and performance in female runners. *Eur J Sport Sci.* 2014;14(2):160–8.

Braun-Fahrländer C, von Mutius E. Can farm milk consumption prevent allergic diseases? *Clin Exp Allergy* 2011;41:29–35.

Brenneisen P, Steinbrenner H, Sies H. Selenium, oxidative stress, and health aspects. *Mol Aspects Med.* 2005 Aug-Oct;26(4–5):256–67

Brilla LR, Conte V. Effects of a Novel Zinc-Magnesium Formulation on Hormones and Strength. *Journal of Exercise Physiology online.* 2000; 3: 26–36.

Brink EJ, Beynen AC. Nutrition and magnesium absorption: a review. *Prog Food Nutr Sci.* 1992;16(2):125–62.

Brownlie T 4th, Utermohlen V, Hinton PS, Giordano C, Haas JD. Marginal iron deficiency without anemia impairs aerobic adaptation among previously untrained women. *Am J Clin Nutr.* 2002 Apr;75(4):734–42.

Brownlie T 4th, Utermohlen V, Hinton PS, Haas JD. Tissue iron deficiency without anemia impairs adaptation in endurance capacity after aerobic training in previously untrained women. *Am J Clin Nutr.* 2004 Mar;79(3):437–43.

Brun JF, Bouchahda C, Chaze D, Benhaddad AA, Micallef JP, Mercier J. The paradox of hematocrit in exercise physiology: which is the “normal” range from an hemorheologist’s viewpoint? *Clin Hemorheol Microcirc.* 2000;22(4):287–303.

Calder PC, Ahluwalia N, Brouns F, Buetler T, Clement K, Cunningham K, Esposito K, Jönsson LS ym. Dietary factors and low-grade inflammation in relation to overweight and obesity. *Br J Nutr.* 2011 Dec;106 Suppl 3:S5–78.

Caudarella R, Vescini F, Rizzoli E, Francucci CM. Salt intake, hypertension, and osteoporosis. *J Endocrinol Invest.* 2009;32(4 Suppl):15–20.

Cheng HH, Lai MH, Hou WC, Huang CL. Antioxidant effects of chromium supplementation with type 2 diabetes mellitus and euglycemic subjects. *J Agric Food Chem.* 2004 Mar 10;52(5):1385–9.

Clark LC, Combs GF Jr, Turnbull BW, Slate EH, Chalker DK, Chow J, Davis LS ym. Effects of selenium supplementation for cancer prevention in patients with carcinoma of the skin. A randomized controlled trial. *Nutritional Prevention of Cancer Study Group. JAMA.* 1996 Dec 25;276(24):1957–63

Clark M, Reed DB, Crouse SF, Armstrong RB. Pre- and post-season dietary intake, body composition, and performance indices of NCAA division I female soccer players. *Int J Sport Nutr Exerc Metab.* 2003 Sep;13(3):303–19.

Clarkson PM. Minerals: exercise performance and supplementation in athletes. *J Sports Sci.* 1991 Summer;9 Spec No:91–116

Clarkson PM, Thompson HS. Antioxidants: what role do they play in physical activity and health? *Am J Clin Nutr.* 2000 Aug;72(2 Suppl):637S-46S

Clayes W , Cardoen S, Daube G ym. Raw or heated cow milk consumption: Review of risks and benefits. *Food Control* 2013;31: 251–262.

Close GL, Ashton T, Cable T, Doran D, Holloway C, McArdle F, McLaren DP. Ascorbic acid supplementation does not attenuate post-exercise muscle soreness following muscle-damaging exercise but may delay the recovery process. *Br J Nutr.* 2006 May;95(5):976–81.

Close, G.L.; Russel, J.; Cobley, J.N.; Owens, D.J.; Wilson, G.; Gregson, W.; Fraser, W.D.; Morton, J.P. Assessment of vitamin D concentration in non-supplemented professional athletes and healthy adults during the winter months in the UK: Implications for skeletal muscle function. *J. Sports Sci.* 2013, 31, 344–353.

Coombes JS, Powers SK, Rowell B, Hamilton KL, Dodd SL, Shanelly RA, Sen CK, Packer L. Effects of vitamin E and alpha-lipoic acid on skeletal muscle contractile properties. *J Appl Physiol (1985).* 2001 Apr;90(4):1424–30.

Cordova A, Alvarez-Mon M. Behaviour of zinc in physical exercise: a special reference to immunity and fatigue. *Neurosci Biobehav Rev.* 1995 Fall;19(3):439–45

Cordova A, Navas FJ. Effect of training on zinc metabolism: changes in serum and sweat zinc concentrations in sportsmen. *Ann Nutr Metab.* 1998;42(5):274–82.

Couzy F, Lafargue P, Guezennecy CY. Zinc metabolism in the athlete: influence of training, nutrition and other factors. *Int J Sports Med.* 1990 Aug;11(4):263–6.

Crawford V, Scheckenbach R, Preuss HG. Effects of niacin-bound chromium supplementation on body composition in overweight African-American women. *Diabetes Obes Metab.* 1999 Nov;1(6):331–7.

Cupisti A, D'Alessandro C, Castrogiovanni S, Barale A, Morelli E. Nutrition survey in elite rhythmic gymnasts. *J Sports Med Phys Fitness.* 2000 Dec;40(4):350–5

Dam BV. Vitamins and sport. *Br J Sports Med.* 1978 Jun;12(2):74–9.

Davies KJ, Quintanilha AT, Brooks GA, Packer L. Free radicals and tissue damage produced by exercise. *Biochem Biophys Res Commun.* 1982 Aug 31;107(4):1198–205

Deakin V. Iron depletion in athletes. Kirjassa: Burke L, Deakin V (toim). Clinical sports nutrition 2nd ed. McGraw-Hill, Australia, 2002.

D'Elia L, Galletti F, Strazzullo P. Dietary salt intake and risk of gastric cancer. *Cancer Treat Res.* 2014;159:83–95.

de Oliveira Kde J, Donangelo CM, de Oliveira AV Jr, da Silveira CL, Koury JC. Effect of zinc supplementation on the antioxidant, copper, and iron status of physically active adolescents. *Cell Biochem Funct.* 2009 Apr;27(3):162–6.

DeRuisseau KC, Cheuvront SN, Haymes EM, Sharp RG. Sweat iron and zinc losses during prolonged exercise. *Int J Sport Nutr Exerc Metab.* 2002 Dec;12(4):428–37

Deuster PA, Day BA, Singh A, Douglass L, Moser-Veillon PB. Zinc status of highly trained women runners and untrained women. *Am J Clin Nutr.* 1989 Jun;49(6):1295–301.

Dietrich T, Nunn M, Dawson-Hughes B, Bischoff-Ferrari HA. Association between serum concentrations of 25-hydroxyvitamin D and gingival inflammation. *Am J Clin Nutr.* 2005 Sep;82(3):575–80.

Dillard CJ, Litov RE, Savin WM, Dumelin EE, Tappel AL. Effects of exercise, vitamin E, and ozone on pulmonary function and lipid peroxidation. *J Appl Physiol.* 1978 Dec;45(6):927–32.

Doetsch AM, Faber J, Lynnerup N, Wätjen I, Bliddal H, Danneskiold-Samsøe. The effects of calcium and vitamin D₃ supplementation on the healing of proximal humerus fracture: a randomized placebo-controlled study. *Calcif Tissue Int* 2004; 75:183–188.

Donaldsson CL, Hullet SB, Vogel JM, Hattner RS, Bayers JH, McMillan DE. Effect of prolonged bed rest on bone mineral. *Metabolism* 1970;19:1071–84

Donangelo CM, Woodhouse LR, King SM, Viteri FE, King JC. Supplemental zinc lowers measures of iron status in young women with low iron reserves. *J Nutr.* 2002 Jul;132(7):1860–4.

Douglas RM, Hemila H, D'Souza R, Chalker EB, Treacy B. Vitamin C for preventing and treating the common cold. *Cochrane Database Syst Rev.* 2004 Oct 18;(4):CD000980.

Douglas RM, Hemilä H. Vitamin C for preventing and treating the common cold. *PLoS Med.* 2005 Jun;2(6):e168; quiz e217. Epub 2005 Jun 28.

Dragan I, Dinu V, Mohora M, Cristea E, Ploesteanu E, Stroescu V. Studies regarding the antioxidant effects of selenium on top swimmers. *Rev Roum Physiol.* 1990 Jan-Mar;27(1):15–20.

Ducros V. Chromium metabolism. A literature review. *Biol Trace Elem Res.* 1992 Jan-Mar;32:65–77.

Evira Research Reports 2/2007 Intake of vitamin A, cadmium and lead via liver foods among Finnish women of fertile age – a quantitative risk assessment

Erdman JW, Jr, Balentine D, Arab L, Beecher G, Dwyer JT, Folts J, et al. Flavonoids and heart health: Proceedings of the ILSI North America Flavonoids Workshop, May 31-June 1, 2005, Washington, DC. *J Nutr.* 2007;137:718S–737

Faber M, Benade AJ. Mineral and vitamin intake in field athletes (discus-, hammer-, javelin-throwers and shotputters). *Int J Sports Med.* 1991 Jun;12(3):324–7.

Fallon KE, Sivy G, Sivy K, Dare A. Changes in haematological parameters and iron metabolism associated with a 1 600 kilometre ultramarathon. *Br J Sports Med.* 1999 Feb;33(1):27–31;

Fine KD, Santa Ana CA, Porter JL, Fordtran JS. Intestinal absorption of magnesium from food and supplements. *J Clin Invest.* 1991 Aug;88(2):396–402.

Finstad EW, Newhouse IJ, Lukaski HC, McAuliffe JE, Stewart CR. The effects of magnesium supplementation on exercise performance. *Med Sci Sports Exerc.* 2001 Mar;33(3):493–8.

Fisher AE, Naughton DP. Iron supplements: the quick fix with long-term consequences. *Nutr J.* 2004 Jan 16;3:2. **Fogelholm M.** Indicators of vitamin and mineral status in athletes' blood: a review. *Int J Sport Nutr.* 1995 Dec;5(4):267–84.

Fogelholm M. Vitamin, mineral and antioxidant needs of athletes. Kirjassa: Burke L, Deakin, V (toim.). Clinical sports nutrition 2nd ed. McGraw-Hill, 2002.

Fogelholm M, Tikkainen H, Naveri H, Harkonen M. High-carbohydrate diet for long distance runners--a practical view-point. *Br J Sports Med.* 1989 Jun;23(2):94–6.

Fogelholm M, Laakso J, Lehto J, Ruokonen I. Dietary intake and indicators of magnesium and zink status in male athletes. *Nutr Res* 1991;11:1111–8

Fogelholm M, Rehunen S, Gref CG, Laakso JT, Lehto J, Ruokonen I, Himberg JJ. Dietary intake and thiamin, iron, and zinc status in elite Nordic skiers during different training periods. *Int J Sport Nutr.* 1992 Dec;2(4):351–65.

Fogelholm GM, Koskinen R, Laakso J, Rankinen T, Ruokonen I. Gradual and rapid weight loss: effects on nutrition and performance in male athletes. *Med Sci Sports Exerc.* 1993 Mar;25(3):371–7.

Frank T, Kuhl M, Makowski B, Bitsch R, Jahreis G, Hubscher J. Does a 100-km walking affect indicators of vitamin status? *Int J Vitam Nutr Res.* 2000 Sep;70(5):238–50.

Fu L, Yun F, Oczak M, Wong BY, Vieth R, Cole DE. Common genetic variants of the vitamin D binding protein (DBP) predict differences in response of serum 25-hydroxyvitamin D [25(OH)D] to vitamin D supplementation. *Clin Biochem.* 2009 Jul;42(10–11):1174–7.

Gardner GW, Edgerton VR, Barnard RJ, Bernauer EM. Cardiorespiratory, hematological and physical performance responses of anemic subjects to iron treatment. *Am J Clin Nutr.* 1975 Sep;28(9):982–8.

Gleeson M, Lancaster GI, Bishop NC. Nutritional strategies to minimise exercise-induced immunosuppression in athletes. *Can J Appl Physiol.* 2001;26 Suppl:S23–35

Gleeson M. Nutritional support to maintain proper immune status during intense training. *Nestle Nutr Inst Workshop Ser.* 2013;75:85–97.

Gohil K, Packer L, de Lumen B, Brooks GA, Terblanche SE. Vitamin E deficiency and vitamin C supplements: exercise and mitochondrial oxidation. *J Appl Physiol.* 1986 Jun;60(6):1986–91.

Goldfarb AH, Bloomer RJ, McKenzie MJ. Combined antioxidant treatment effects on blood oxidative stress after eccentric exercise. *Med Sci Sports Exerc.* 2005 Feb;37(2):234–9.

Gollnick PD, Armstrong RB, Saltin B, Saubert CW 4th, Sembrowich WL, Shepherd RE. Effect of training on enzyme activity and fiber composition of human skeletal muscle. *J Appl Physiol.* 1973 Jan;34(1):107–11.

Gomez-Cabrera MC, Domenech E, Romagnoli M, Arduini A, Borras C, Pallardo FV, Sastre J, Viña J. Oral administration of vitamin C decreases muscle mitochondrial biogenesis and hampers training-induced adaptations in endurance performance. *Am J Clin Nutr.* 2008 Jan;87(1):142–9.

Grievink L, Jansen SM, van't Veer P, Brunekreef B. Acute effects of ozone on pulmonary function of cyclists receiving antioxidant supplements. *Occup Environ Med.* 1998 Jan;55(1):13–7.

Grievink L, Zijlstra AG, Ke X, Brunekreef B. Double-blind intervention trial on modulation of ozone effects on pulmonary function by antioxidant supplements. *Am J Epidemiol.* 1999 Feb 15;149(4):306–14.

Guilland JC, Penaranda T, Gallet C, Boggio V, Fuchs F, Klepping J. Vitamin status of young athletes including the effects of supplementation. *Med Sci Sports Exerc.* 1989 Aug;21(4):441–9.

Hallberg L, Rossander-Hulten L, Brune M, Gleerup A. Bioavailability in man of iron in human milk and cow's milk in relation to their calcium contents. *Pediatr Res.* 1992a May;31(5):524–7.

Hallberg L, Rossander-Hulten L, Brune M, Gleerup A. Calcium and iron absorption: mechanism of action and nutritional importance. *Eur J Clin Nutr.* 1992b May;46(5):317–27.

Handelman GJ, Machlin LJ, Fitch K, Weiter JJ, Dratz EA. Oral alpha-tocopherol supplements decrease plasma gamma-tocopherol levels in humans. *J Nutr.* 1985 Jun;115(6):807–13.

Haymes EM. Vitamin and mineral supplementation to athletes. *Int J Sport Nutr.* 1991 Jun;1(2):146–69.

Haymes EM, Lamanca JJ. Iron loss in runners during exercise. Implications and recommendations. *Sports Med.* 1989 May;7(5):277–85

Helldán A, Raulio S, Kosola M, Tapanainen H, Ovaskainen M-L, Virtanen S. Finraavinto 2012 –tutkimus. The National FINDIET 2012 Survey. THL Raportti 16/2013

Hemilä H. Vitamin C and common cold incidence: a review of studies with subjects under heavy physical stress. *Int J Sports Med.* 1996 Jul;17(5):379–83.

Hemila H, Virtamo J, Albane D, Kaprio J. Physical activity and the common cold in men administered vitamin E and beta-carotene. *Med Sci Sports Exerc.* 2003 Nov;35(11):1815–20

Henderson LM, Brewer GJ, Dressman JB, Swidan SZ, DuRoss DJ, Adair CH, Barnett JL, Berardi RR. Effect of intragastric pH on the absorption of oral zinc acetate and zinc oxide in young healthy volunteers. *JPEN J Parenter Enteral Nutr.* 1995 Sep-Oct;19(5):393–7.

Hennekens CH, Buring JE, Manson JE, Stampfer M, Rosner B, Cook NR, Belanger C, LaMotte F, Gaziano JM ym. Lack of effect of long-term supplementation with beta carotene on the incidence of malignant neoplasms and cardiovascular disease. *N Engl J Med.* 1996 May 2;334(18):1145–9.

Hickson RC. Skeletal muscle cytochrome c and myoglobin, endurance, and frequency of training. *J Appl Physiol.* 1981 Sep;51(3):746–9.

Hickson JF Jr, Schrader J, Trischler LC. Dietary intakes of female basketball and gymnastics athletes. *J Am Diet Assoc.* 1986 Feb;86(2):251–3.

Hirvonen T, Pietinen P, Virtanen M, Albane D, Virtamo J. Nutrient intake and use of beverages and the risk of kidney stones among male smokers. *Am J Epidemiol.* 1999;150:187–194.

Hoeger WW, Harris C, Long EM, Hopkins DR. Four-week supplementation with a natural dietary compound produces favorable changes in body composition. *Adv Ther.* 1998 Sep-Oct;15(5):305–14.

- Holick MF.** Environmental factors that affect the cutaneous production of vitamin D. *Am J Clin Nutr* 1995;61:638S-45S.
- Holick MF, Matsuoka LY, Wortsman J.** Age, vitamin D, and solar ultraviolet. *Lancet* 1989;2:1104–1105.
- Horvath PJ, Eagen CK, Ryer-Calvin SD, Pendergast DR.** The effects of varying dietary fat on the nutrient intake in male and female runners. *J Am Coll Nutr*. 2000 Feb;19(1):42–51.
- Howlett KF, Spriet LL, Hargreaves M.** Carbohydrate metabolism during exercise in females: effect of reduced fat availability. *Metabolism*. 2001 Apr;50(4):481–7.
- Hua Y, Clark S, Ren J, Sreejayan N.** Molecular mechanisms of chromium in alleviating insulin resistance. *J Nutr Biochem*. 2012 Apr;23(4):313–9.
- Hughes DA, Wright AJ, Finglas PM, Peerless AC, Bailey AL, Astley SB, Pinder AC, Southon S.** The effect of beta-carotene supplementation on the immune function of blood monocytes from healthy male nonsmokers. *J Lab Clin Med*. 1997 Mar;129(3):309–17.
- Hunt JR.** Bioavailability of iron, zinc, and other trace minerals from vegetarian diets. *Am J Clin Nutr*. 2003 Sep;78(3 Suppl):633S-639S
- Ito Y, Wakai K, Suzuki K, Ozasa K, Watanabe Y, Seki N, Ando M, Nishino Y, Kondo T, Ohno Y, Tamakoshi A.** Lung Cancer Mortality and Serum Levels of Carotenoids, Retinol, Tocopherols, and Folic Acid in Men and Women: a Case-Control Study Nested in the JACC Study. *J Epidemiol*. 2005;15 Suppl 2:S140–9.
- Itokawa Y, Kimura M, Nishino K, Mino M, Kitagawa M, Matsuoka M, Otsuka H.** Blood levels and urinary excretion of thiamin and riboflavin during oral administration of multivitamin tablets to healthy adults. *J Nutr Sci Vitaminol (Tokyo)*. 1992;Spec No:438–41.
- Jakulj F, Zernicke K, Bacon SL, van Wielingen LE, Key BL, West SG, Campbell TS.** A high-fat meal increases cardiovascular reactivity to psychological stress in healthy young adults. *J Nutr*. 2007 Apr;137(4):935–9.
- Jeejeebhoy KN, Chu RC, Marliss EB, Greenberg GR, Bruce-Robertson A.** Chromium deficiency, glucose intolerance, and neuropathy reversed by chromium supplementation, in a patient receiving long-term total parenteral nutrition. *Am J Clin Nutr*. 1977 Apr;30(4):531–8.
- Jiang Q, Elson-Schwab I, Courtemanche C, Ames BN.** gamma-tocopherol and its major metabolite, in contrast to alpha-tocopherol, inhibit cyclooxygenase activity in macrophages and epithelial cells. *Proc Natl Acad Sci U S A*. 2000 Oct 10;97(21):11494–9.
- Jones DP.** Redefining oxidative stress. *Antioxid Redox Signal*. 2006 Sep-Oct;8(9–10):1865–79.
- Jonnalagadda SS, Ziegler PJ, Nelson JA.** Food preferences, dieting behaviors, and body image perceptions of elite figure skaters. *Int J Sport Nutr Exerc Metab*. 2004 Oct;14(5):594–606.
- Kalkwarf HJ, Specker BL, Henbi JE, Vieira NE, Yerger AL.** Intestinal calcium absorption of women during lactation. *Am J Clin Nutr* 1996;63:526–31.
- Kalkwarf HJ, Specker BL, Bianchi DC, Ranz J, Ho M.** The effect of calcium supplementation on bone density during lactation and after weaning. *N Engl J Med* 1997;337:523–8.
- Kanter MM, Nolte LA, Holloszy JO.** Effects of an antioxidant vitamin mixture on lipid peroxidation at rest and postexercise. *J Appl Physiol*. 1993 Feb;74(2):965–9.
- Khaled S, Brun JF, Micallel JP, Bardet L, Cassanas G, Monnier JF, Orsetti A.** Serum zinc and blood rheology in sportsmen (football players). *Clin Hemorheol Microcirc*. 1997 Jan-Feb;17(1):47–58.
- Khaled S, Brun JF, Wagner A, Mercier J, Bringer J, Prefaut C.** Increased blood viscosity in iron-depleted elite athletes. *Clin Hemorheol Microcirc*. 1998 Jul;18(4):309–18.
- Khaled S, Brun JF, Cassanas G, Bardet L, Orsetti A.** Effects of zinc supplementation on blood rheology during exercise. *Clin Hemorheol Microcirc*. 1999;20(1):1–10.
- Kikukawa A, Kobayashi A.** Changes in urinary zinc and copper with strenuous physical exercise. *Aviat Space Environ Med*. 2002 Oct;73(10):991–5.
- Knek P, Kumpulainen J, Jarvinen R, Rissanen H, Heliovaara M, Reunanen A, Hakulinen T, Aromaa A.** Flavonoid intake and risk of chronic diseases. *Am J Clin Nutr*. 2002 Sep;76(3):560–8.

Knippl M, Mauri L, Bellushi R, Bana G. The action of thiamine on the production of lactic acid in cyclists. *Med Sports* 1986; 39(1):11.

Koehler K, Parr MK, Geyer H, Mester J, Schanzer W. Serum testosterone and urinary excretion of steroid hormone metabolites after administration of a high-dose zinc supplement. *Eur J Clin Nutr.* 2009 Jan;63(1):65–70.

Kozlovsky AS, Moser PB, Reiser S, Anderson RA. Effects of diets high in simple sugars on urinary chromium losses. *Metabolism.* 1986 Jun;35(6):515–8.

Kritchevsky SB. beta-Carotene, carotenoids and the prevention of coronary heart disease. *J Nutr.* 1999 Jan;129(1):5–8.

Lahti-Koski M, Valsta LM, Alftan G, Tapanainen H, Aro A. Iron status of adults in the capital area of Finland. *Eur J Nutr.* 2003 Oct;42(5):287–92.

Lamberg-Allardt C, Brustad M, Meyer HE, Steingrimsdottir L. Vitamin D – a systematic literature review for the 5th edition of the Nordic Nutrition Recommendations. *Food & Nutrition Research* 2013; 57: 22671.

Lappe YM. Calcium and vitamin d supplementation decreases incidence of stress fractures in female navy recruits. *J Bone Miner Res.* 23 (2008), pp. 741–749

LeBlanc A, Schneider W, Evans H, Engelbretson D, Krebs J. Bone mineral loss and recovery after 17 weeks of bed rest. *J Bone Miner Res.* 1990;5:843–50

Lee EH, Myung SK, Jeon YJ, Kim Y, Chang YJ, Ju W, Seo HG, Huh BY. Effects of selenium supplements on cancer prevention: meta-analysis of randomized controlled trials. *Nutr Cancer.* 2011 Nov;63(8):1185–95.

Leitzmann C. Vegetarian diets: what are the advantages? *Forum Nutr.* 2005;(57):147–56

Leonard SW, Leklem JE. Plasma B-6 vitamer changes following a 50-km ultra-marathon. *Int J Sport Nutr Exerc Metab.* 2000 Sep;10(3):302–14.

Lindberg JS, Zobitz MM, Poindexter JR, Pak CY. Magnesium bioavailability from magnesium citrate and magnesium oxide. *J Am Coll Nutr.* 1990 Feb;9(1):48–55.

Litoff D, Scherzer H, Harrison J. Effects of pantothenic acid supplementation on human exercise. *Med Sci Sports Exerc.* 1985; 17:287.

Loud KJ, Gordon CNM, Micheli LJ, Field AE. Correlates of stress fractures among preadolescent and adolescent girls. *Pediatrics.* 2005;115:399–406.

Lukaski HC. Magnesium, zinc, and chromium nutriture and physical activity. *Am J Clin Nutr.* 2000 Aug;72(2 Suppl):585S-93S

Lukaski HC. Vitamin and mineral status: effects on physical performance. *Nutrition.* 2004 Jul-Aug;20(7–8):632–44.

Machefer G, Groussard C, Zouhal H, Vincent S, Youssef H, Faure H, Malardé L, Gratas-Delamarche A. Nutritional and plasmatic antioxidant vitamins status of ultra endurance athletes. *J Am Coll Nutr.* 2007 Aug;26(4):311–6.

Manore MM. Vitamin B6 and exercise. *Int J Sport Nutr.* 1994 Jun;4(2):89–103.

Manore MM. Effect of physical activity on thiamine, riboflavin, and vitamin B-6 requirements. *Am J Clin Nutr.* 2000 Aug;72(2 Suppl):598S-606S

Manore MM, Leklem JE. Effect of carbohydrate and vitamin B6 on fuel substrates during exercise in women. *Med Sci Sports Exerc.* 1988 Jun;20(3):233–41.

Margaritis I, Tessier F, Prou E, Marconnet P, Marini JF. Effects of endurance training on skeletal muscle oxidative capacities with and without selenium supplementation. *J Trace Elem Med Biol.* 1997 Apr;11(1):37–43.

Martin BR, Davis S, Campbell WW, Weaver CM. Exercise and calcium supplementation: effects on calcium homeostasis in sportswomen. *Med Sci Sports Exerc.* 2007 Sep;39(9):1481–6.

Masharani U, Gjerde C, McCoy S, Maddux BA, Hessler D, Goldfine ID, Youngren JF. Chromium supplementation in non-obese non-diabetic subjects is associated with a decline in insulin sensitivity. *BMC Endocr Disord.* 2012 Nov 30;12:31.

Matter M, Stittfall T, Graves J, Myburgh K, Adams B, Jacobs P, Noakes TD. The effect of iron and folate therapy on maximal exercise performance in female marathon runners with iron and folate deficiency. *Clin Sci (Lond).* 1987 Apr;72(4):415–22.

Maughan RJ. Fundamentals of Sports Nutrition: Application to Sports Drinks. Kirjassa Sports Drinks – Basic Science an d Practical Aspects. Maughan RJ, Murray R (toim.) CRC Press, Boca Raton, Florida: 2001.

Maughan RJ, Leiper JB. Sodium intake and post-exercise rehydration in man. *Eur J Appl Physiol Occup Physiol.* 1995;71(4):311–9.

Maxwell SR, Jakeman P, Thomason H, Leguen C, Thorpe GH. Changes in plasma antioxidant status during eccentric exercise and the effect of vitamin supplementation. *Free Radic Res Commun.* 1993;19(3):191–202.

McAnulty S, McAnulty L, Nieman D, Morrow J, Dumke C, Henson D. Effect of NSAID on muscle injury and oxidative stress. *Int J Sports Med.* 2007 Nov;28(11):909–15.

McArdle WD, Katch FI, Katch VL. Vitamins, minerals and water. Kirjassa: Exercise Physiology 5th ed. Lippincott Williams & Wilkins, USA, 2001.

McBride JM, Kraemer WJ, Triplett-McBride T, Sebastianelli W. Effect of resistance exercise on free radical production. *Med Sci Sports Exerc.* 1998 Jan;30(1):67–72.

McGee SR. Muscle cramps. *Arch Intern Med.* 1990 Mar;150(3):511–8.

McGinley C, Shafat A, Donnelly AE. Does antioxidant vitamin supplementation protect against muscle damage? *Sports Med.* 2009;39(12):1011–32.

Meydani M. Vitamin E. *Lancet.* 1995 Jan 21;345(8943):170–5.

Meydani M, Evans WJ, Handelman G, Biddle L, Fielding RA, Meydani SN, Burrill J, Fiatarone MA, Blumberg JB, Cannon JG. Protective effect of vitamin E on exercise-induced oxidative damage in young and older adults. *Am J Physiol.* 1993 May;264(5 Pt 2):R992–8.

Micheletti A, Rossi R, Rufini S. Zinc status in athletes: relation to diet and exercise. *Sports Med.* 2001;31(8):577–82

Mickleborough TD, Lindley MR, Ray S. Dietary salt, airway inflammation, and diffusion capacity in exercise-induced asthma. *Med Sci Sports Exerc.* 2005 Jun;37(6):904–14.

Minetto MA, Holobar A, Botter A, Farina D. Origin and development of muscle cramps. *Exerc Sport Sci Rev.* 2013 Jan;41(1):3–10.

Monsen ER. Iron nutrition and absorption: dietary factors which impact iron bioavailability. *J Am Diet Assoc.* 1988 Jul;88(7):786–90

Mooren FC, Golf SW, Volker K. Effect of magnesium on granulocyte function and on the exercise induced inflammatory response. *Magnes Res.* 2003 Mar;16(1):49–58.

Moran DS, McClung JP, Kohen T, Lieberman HR. Vitamin d and physical performance. *Sports Med.* 2013 Jul;43(7):601–11.

Moretti C, Fabbri A, Gnessi L, Bonifacio V, Fraioli F, Isidori A. Pyridoxine (B6) suppresses the rise in prolactin and increases the rise in growth hormone induced by exercise. *N Engl J Med.* 1982 Aug 12;307(7):444–5.

Murphy NM, Carroll P. The effect of physical activity and its interaction with nutrition on bone health. *Proceedings of the Nutrition* 2003;62:829–838.

Mutanen M, Voutilainen E. Vitamiinit ja kivennäisaineet. Kirjassa: Aro A, Mutanen M, Uusitupa M (toim). Ravitsemustiede. Duodecim, Helsinki, 1999.

Murray R, Stofan JR. Formulating Carbohydrate-Electrolyte Drinks for Optimal Efficacy. Kirjassa Sports Drinks – Basic Science an d Practical Aspects. Maughan RJ, Murray R (toim.) CRC Press, Boca Raton, Florida: 2001, 206–207

Nachtigall D, Nielsen P, Fischer R, Engelhardt R, Gabbe EE. Iron deficiency in distance runners. A reinvestigation using Fe-labelling and non-invasive liver iron quantification. *Int J Sports Med.* 1996 Oct;17(7):473–9.

Navarro M, Wood RJ. Plasma changes in micronutrients following a multivitamin and mineral supplement in healthy adults. *J Am Coll Nutr.* 2003 Apr;22(2):124–32.

Newhouse IJ, Finstad EW. The effects of magnesium supplementation on exercise performance. *Clin J Sport Med.* 2000 Jul;10(3):195–200.

Nielsen P, Nachtigall D. Iron supplementation in athletes. Current recommendations. *Sports Med.* 1998 Oct;26(4):207–16.

Nieman DC, Butler JV, Pollett LM, Dietrich SJ, Lutz RD. Nutrient intake of marathon runners. *J Am Diet Assoc.* 1989 Sep;89(9):1273–8.

Nieves ym. Nutritional factors that influence change in bone density and stress fracture risk among young female cross-country runners. *PM R.* 2 (2010), pp. 740–750

Nikolaidis MG, Kerksick CM, Lamprecht M, McAnulty SR. Does vitamin C and E supplementation impair the favorable adaptations of regular exercise? *Oxid Med Cell Longev.* 2012;2012:707941.

Nissen SL, Sharp RL. Effect of dietary supplements on lean mass and strength gains with resistance exercise: a meta-analysis. *J Appl Physiol.* 2003 Feb;94(2):651–9.

Nuviala RJ, Lapieza MG, Bernal E. Magnesium, zinc, and copper status in women involved in different sports. *Int J Sport Nutr.* 1999 Sep;9(3):295–309.

Ogan D & Pritchett K. Vitamin D and the Athlete: Risks, Recommendations, and Benefits. *Nutrients* 2013, 5, 1856–1868

Ohno H, Yahata T, Sato Y, Yamamura K, Taniguchi noin Physical training and fasting erythrocyte activities of free radical scavenging enzyme systems in sedentary men. *Eur J Appl Physiol Occup Physiol.* 1988;57(2):173–6.

Onakpoya I, Posadzki P, Ernst E. Chromium supplementation in overweight and obesity: a systematic review and meta-analysis of randomized clinical trials. *Obes Rev.* 2013 Jun;14(6):496–507.

Osteoporoosi: Käypä hoito-suositus, 2014

Outila TA, Kärkkäinen MU, Seppänen RH, Lamberg-Allardt CJ. Dietary intake of vitamin D in premenopausal, healthy vegans was insufficient to maintain concentrations of serum 25-hydroxyvitamin D and intact parathyroid hormone within normal ranges during the winter in Finland. *J Am Diet Assoc* 2000;100:434–441

Paajanen ym. No difference in symptoms during challenges with homogenized and unhomogenized cow's milk in subjects with subjective hypersensitivity to homogenized milk. *J Dairy Res* 2003;70(2):175–9.

Peterlik M, Cross HS. Vitamin D and calcium deficits predispose for multiple chronic diseases. *Eur J Clin Invest* 2005; 35:290–304.

Peternelj TT, Coombes JS. Antioxidant supplementation during exercise training: beneficial or detrimental? *Sports Med.* 2011 Dec 1;41(12):1043–69.

Peters EM, Cambell A, Pawley L. Vitamin A fails to increase resistance to upper respiratory tract infection in distance runners. *S Afr J Sports Med* 1992;7:3–7

Peters EM, Goetzsche JM, Grobbelaar B, Noakes TD. Vitamin C supplementation reduces the incidence of postrace symptoms of upper-respiratory-tract infection in ultramarathon runners. *Am J Clin Nutr.* 1993 Feb;57(2):170–4.

Porea TJ, Belmont JW, Mahoney DH Jr. Zinc-induced anemia and neutropenia in an adolescent. *J Pediatr.* 2000 May;136(5):688–90.

Powers SK. Antioxidant supplementaton for athletes: sense **or nonsense?** IOC Consensus In Sports Nutrition Conference Lausanne, Switzerland 25th–27th October 2010.

Powers SK, Ji LL, Leeuwenburgh C. Exercise training-induced alterations in skeletal muscle antioxidant capacity: a brief review. *Med Sci Sports Exerc.* 1999 Jul;31(7):987–97.

Powers SK, Jackson MJ. Exercise-induced oxidative stress: cellular mechanisms and impact on muscle force production. *Physiol Rev.* 2008 Oct;88(4):1243–76.

Puhl JL, Runyan WS, Kruse SJ. Erythrocyte changes during training in high school women cross-country runners. Res Q Exerc Sport. 1981 Dec;52(4):484–94

Rankinen T, Lyytikainen S, Vanninen E, Penttila I, Rauramaa R, Uusitupa M. Nutritional status of the Finnish elite ski jumpers. Med Sci Sports Exerc. 1998 Nov;30(11):1592–7.

Razavi, M.Z.; Nazarali, P.; Hanachi, P. The effect of an exercise programme and consumption of vitamin D on performance and respiratory indicators in patients with asthma. Sport Sci. Health 2011, 6, 89–92.

Reid MB. Invited Review: redox modulation of skeletal muscle contraction: what we know and what we don't. J Appl Physiol (1985). 2001 Feb;90(2):724–31. Review.

Rimm EB, Stampfer MJ, Ascherio A, Giovannucci E, Colditz GA, Willett WC. Vitamin E consumption and the risk of coronary heart disease in men. N Engl J Med. 1993 May 20;328(20):1450–6.

Ristow M, Zarse K, Oberbach A, Klöting N, Birringer M, Kiehntopf M, Stumvoll M, Kahn CR, Blüher M. Antioxidants prevent health-promoting effects of physical exercise in humans. Proc Natl Acad Sci U S A. 2009 May 26;106(21):8665–70.

Robertson JD, Maughan RJ, Duthie GG, Morrice PC. Increased blood antioxidant systems of runners in response to training load. Clin Sci (Lond). 1991 Jun;80(6):611–8.

Rodenberg RE, Gustafson S. Iron as an ergogenic aid: ironclad evidence? Curr Sports Med Rep. 2007 Jul;6(4):258–64.

Rokitzki L, Berg A, Keul J. Blood and serum status of water- and fat-soluble vitamins in athletes and non-athletes. Int J Vitam Nutr Res Suppl. 1989;30:192–7.

Rokitzki L, Sagredos AN, Reuss F, Buchner M, Keul J. Acute changes in vitamin B6 status in endurance athletes before and after a marathon. Int J Sport Nutr. 1994a Jun;4(2):154–65.

Rokitzki L, Sagredos AN, Reuss F, Cufi D, Keul J. Assessment of vitamin B6 status of strength and speedpower athletes. J Am Coll Nutr. 1994b Feb;13(1):87–94.

Rokitzki L, Sagredos A, Keck E, Sauer B, Keul J. Assessment of vitamin B2 status in performance athletes of various types of sports. J Nutr Sci Vitaminol (Tokyo). 1994c Feb;40(1):11–22

Rokitzki L, Logemann E, Sagredos AN, Murphy M, Wetzel-Roth W, Keul J. Lipid peroxidation and antioxidative vitamins under extreme endurance stress. Acta Physiol Scand. 1994 Jun;151(2):149–58.

Roodenburg AJ, Leenen R, van het Hof KH, Weststrate JA, Tijburg LB. Amount of fat in the diet affects bioavailability of lutein esters but not of alpha-carotene, beta-carotene, and vitamin E in humans. Am J Clin Nutr. 2000 May;71(5):1187–93.

Rudzki SJ, Hazard H, Collinson D. Gastrointestinal blood loss in triathletes: its etiology and relationship to sports anaemia. Aust J Sci Med Sport. 1995 Mar;27(1):3–8.

Salzman MB, Smith EM, Koo C. Excessive oral zinc supplementation. J Pediatr Hematol Oncol. 2002 Oct;24(7):582–4.

Saris NE, Mervaala E, Karppanen H, Khawaja JA, Lewenstein A. Magnesium. An update on physiological, clinical and analytical aspects. Clin Chim Acta. 2000 Apr;294(1–2):1–26

Schrauzer GN. Nutritional selenium supplements: product types, quality, and safety. J Am Coll Nutr. 2001 Feb;20(1):1–4.

Schroder H, Navarro E, Mora J, Galiano D, Tramullas A. Effects of alpha-tocopherol, beta-carotene and ascorbic acid on oxidative, hormonal and enzymatic exercise stress markers in habitual training activity of professional basketball players. Eur J Nutr. 2001 Aug;40(4):178–84.

Schwartz AE, Vanagunas A, Kamel PL. Endoscopy to evaluate gastrointestinal bleeding in marathon runners. Ann Intern Med. 1990 Oct 15;113(8):632–3.

Schwellnus MP, Derman EW, Noakes TD. Aetiology of skeletal muscle ‘cramps’ during exercise: a novel hypothesis. J Sports Sci. 1997 Jun;15(3):277–85

Schwellnus MP, Nicol J, Laubscher R, Noakes TD. Serum electrolyte concentrations and hydration status are not associated with exercise associated muscle cramping (EAMC) in distance runners. *Br J Sports Med.* 2004 Aug;38(4):488–92.

Schwellnus MP. Muscle cramping in the marathon : aetiology and risk factors. *Sports Med.* 2007;37(4–5):364–7.

Schwellnus MP. Cause of exercise associated muscle cramps (EAMC)--altered neuromuscular control, dehydration or electrolyte depletion? *Br J Sports Med.* 2009 Jun;43(6):401–8.

Schwellnus MP, Drew N, Collins M. Increased running speed and previous cramps rather than dehydration or serum sodium changes predict exercise-associated muscle cramping: a prospective cohort study in 210 Ironman triathletes. *Br J Sports Med.* 2011 Jun;45(8):650–6

Sen CK. Antioxidants in exercise nutrition. *Sports Med.* 2001;31(13):891–908

Sen CK, Atalay M, Agren J, Laaksonen DE, Roy S, Hanninen O. Fish oil and vitamin E supplementation in oxidative stress at rest and after physical exercise. *J Appl Physiol.* 1997 Jul;83(1):189–95.

Shafat A, Butler P, Jensen RL, Donnelly AE. Effects of dietary supplementation with vitamins C and E on muscle function during and after eccentric contractions in humans. *Eur J Appl Physiol.* 2004 Oct;93(1–2):196–202

Shang G, Collins M, Schwellnus MP. Factors associated with a self-reported history of exercise-associated muscle cramps in Ironman triathletes: a case-control study. *Clin J Sport Med.* 2011 May;21(3):204–10.

Shaskey DJ, Green GA. Sports haematology. *Sports Med.* 2000 Jan;29(1):27–38.

Shils ME, Olson JA, Shike M (toim.). Modern nutrition in health and disease. 8th ed. Williams & Wilkins, USA, 1994.

Simon-Schnass IM. Nutrition at high altitude. *J Nutr.* 1992 Mar;122(3 Suppl):778–81

Smith JA, Gray AB, Pyne DB, Baker MS, Telford RD, Weidemann MJ. Moderate exercise triggers both priming and activation of neutrophil subpopulations. *Am J Physiol.* 1996 Apr;270(4 Pt 2):R838–45.

Smith JK, Grisham MB, Granger DN, Korthuis RJ. Free radical defense mechanisms and neutrophil infiltration in postischemic skeletal muscle. *Am J Physiol.* 1989 Mar;256(3 Pt 2):H789–93.

Smyth PP, Duntas LH. Iodine uptake and loss – can frequent strenuous exercise induce iodine deficiency? *Horm Metab Res.* 2005 Sep;37(9):555–8.

Specker BL, Vieira NE, O'Brian KO, Ho ML, Heubi JE, Abrams SA, Yerger AL. Calcium kinetics in lactating women with low and high calcium intakes. *Am J Clin Nutr.* 1994;59:593–9.

Srinath Reddy K, Katan MB. Diet, nutrition and the prevention of hypertension and cardiovascular diseases. *Public Health Nutr.* 2004 Feb;7(1A):167–86.

Steen SN, Mayer K, Brownell KD, Wadden TA. Dietary intake of female collegiate heavyweight rowers. *Int J Sport Nutr.* 1995 Sep;5(3):225–31.

Sulzer NU, Schwellnus MP, Noakes TD. Serum electrolytes in Ironman triathletes with exercise-associated muscle cramping. *Med Sci Sports Exerc.* 2005 Jul;37(7):1081–5.

Suter PM, Sierro C, Vetter W. Nutritional factors in the control of blood pressure and hypertension. *Nutr Clin Care.* 2002 Jan-Feb;5(1):9–19

Svilaas A, Sakhi AK, Andersen LF, Svilaas T, Strom EC, Jacobs DR Jr, Ose L, Blomhoff R. Intakes of antioxidants in coffee, wine, and vegetables are correlated with plasma carotenoids in humans. *J Nutr.* 2004 Mar;134(3):562–7.

Tangpricha V, Koutkia P, Rieke SM, Chen TC, Perez AA, Holick MF. Fortification of orange juice with vitamin D: a novel approach for enhancing vitamin D nutritional health. *Am J Clin Nutr.* 2003 Jun;77(6):1478–83.

Telford RD, Catchpole EA, Deakin V, McLeay AC, Plank AW. The effect of 7 to 8 months of vitamin/mineral supplementation on the vitamin and mineral status of athletes. *Int J Sport Nutr.* 1992 Jun;2(2):123–34.

Tenforde YM. Evaluating the relationship of calcium and vitamin D in the prevention of stress fracture injuries in the young athlete: a review of the literature. *PM R.* 2010 Oct;2(10):945–9.

- Tessier F, Margaritis I, Richard MJ, Moynot C, Marconnet P.** Selenium and training effects on the glutathione system and aerobic performance. *Med Sci Sports Exerc.* 1995 Mar;27(3):390–6.
- Theriault A, Chao JT, Wang Q, Gapor A, Adeli K.** Tocotrienol: a review of its therapeutic potential. *Clin Biochem.* 1999 Jul;32(5):309–19.
- Thompson D, Williams C, Kingsley M, Nicholas CW, Lakomy HK, McArdle F, Jackson MJ.** Muscle soreness and damage parameters after prolonged intermittent shuttle-running following acute vitamin C supplementation. *Int J Sports Med.* 2001a Jan;22(1):68–75.
- Thompson D, Williams C, McGregor SJ, Nicholas CW, McArdle F, Jackson MJ, Powell JR.** Prolonged vitamin C supplementation and recovery from demanding exercise. *Int J Sport Nutr Exerc Metab.* 2001b Dec;11(4):466–81.
- Tian H, Guo X, Wang X, He Z, Sun R, Ge S, Zhang Z.** Chromium picolinate supplementation for overweight or obese adults. *Cochrane Database Syst Rev.* 2013 Nov 29;11:CD010063.
- Trang HM, Cole DEC, Rubin LA, Pierratos A, Siu S, Vieth R.** Evidence that vitamin D₃ increases serum 25-hydroxy vitamin D more effectively than does vitamin D₂. *Am J Clin Nutr* 1998;68:854–858.
- Tuntawiroon M, Sritongkul N, Brune M, Rossander-Hulten L, Pleehachinda R, Suwanik R ym.** Dose-dependent inhibitory effect of phenolic compounds in foods on nonheme-iron absorption in men. *Am J Clin Nutr.* 1991 Feb;53(2):554–7.
- Tilikainen R, Pöllänen K, Lahti-Koski M, Borg P.** Suomalaisten nuorten urheilijoiden ravinnonsaanti. *Urheilijan ravinto* 2001/1:45–49
- Valk EE, Hornstra G.** Relationship between vitamin E requirement and polyunsaturated fatty acid intake in man: a review. *Int J Vitam Nutr Res.* 2000 Mar;70(2):31–42.
- van der Beek EJ.** Vitamins and endurance training. Food for running or faddish claims? *Sports Med.* 1985 May-Jun;2(3):175–97.
- van der Beek EJ.** Vitamin supplementation and physical exercise performance. *J Sports Sci.* 1991 Summer;9 Spec No:77–90
- van der Beek EJ, van Dokkum W, Schrijver J, Wesstra A, Kistemaker C, Hermus RJ.** Controlled vitamin C restriction and physical performance in volunteers. *J Am Coll Nutr.* 1990 Aug;9(4):332–9.
- van der Beek EJ, van Dokkum W, Wedel M, Schrijver J, van den Berg H.** Thiamin, riboflavin and vitamin B6: impact of restricted intake on physical performance in man. *J Am Coll Nutr.* 1994 Dec;13(6):629–40.
- van der Velde RY, Brouwers JR, Geusens PP, Lems WF, van den Bergh JP.** Calcium and vitamin D supplementation: state of the art for daily practice. *Food Nutr Res.* 2014 Aug 7;58.
- van Het Hof KH, West CE, Weststrate JA, Hautvast JG.** Dietary factors that affect the bioavailability of carotenoids. *J Nutr.* 2000 Mar;130(3):503–6.
- Van Loan MD, Sutherland B, Lowe NM, Turnlund JR, King JC.** The effects of zinc depletion on peak force and total work of knee and shoulder extensor and flexor muscles. *Int J Sport Nutr.* 1999 Jun;9(2):125–35.
- Viitala P, Newhouse IJ.** Vitamin E supplementation, exercise and lipid peroxidation in human participants. *Eur J Appl Physiol.* 2004 Oct;93(1–2):108–15.
- Vincent JB.** The potential value and toxicity of chromium picolinate as a nutritional supplement, weight loss agent and muscle development agent. *Sports Med.* 2003;33(3):213–30.
- Volpe SL.** Magnesium in disease prevention and overall health. *Adv Nutr.* 2013 May 1;4(3):378S–83S.
- VRN.** Valtion ravitsemusneuvottelukunta. *Terveytä ruuasta – Suomalaiset ravitsemussuositukset* 2014. Juvenes Oy, Helsinki 2014.
- Välimäki V-V, Alftan H, Lehmuskallio E, Löyttyniemi E, Sahi T, Suominen H, Välimäki MJ.** Risk factors for clinical stress fractures in male military recruits: A prospective cohort study. *Bone* 2005;37:267–273.
- Walker AF, Marakis G, Christie S, Byng M.** Mg citrate found more bioavailable than other Mg preparations in a randomised, double-blind study. *Magnes Res.* 2003 Sep;16(3):183–91.

Watson TA, Blake RJ, Callister R, Garg ML. Antioxidant-restricted diet reduces plasma nonesterified fatty acids in trained athletes. *Lipids*. 2005 Apr;40(4):433–5.

Weber P, Bendich A, Machlin LJ. Vitamin E and human health: rationale for determining recommended intake levels. *Nutrition*. 1997 May;13(5):450–60

Weaver CM, Rajaram S. Exercise and iron status. *J Nutr*. 1992 Mar;122(3 Suppl):782–7.

Weaver CM, Williams RP, Heaney R. Choices for achieving adequate dietary calcium with a vegetarian diet. *Am J Clin Nutr* 1999;70(suppl):543S-8S.

Webster MJ. Physiological and performance responses to supplementation with thiamin and pantothenic acid derivatives. *Eur J Appl Physiol Occup Physiol*. 1998 May;77(6):486–91.

Webster MJ, Scheett TP, Doyle MR, Branz M. The effect of a thiamin derivative on exercise performance. *Eur J Appl Physiol Occup Physiol*. 1997;75(6):520–4.

Weight LM, Noakes TD, Labadarios D, Graves J, Jacobs P, Berman PA. Vitamin and mineral status of trained athletes including the effects of supplementation. *Am J Clin Nutr*. 1988 Feb;47(2):186–91.

Weller E, Bachert P, Meinck HM, Friedmann B, Bartsch P, Mairbaurl H. Lack of effect of oral Mg-supplementation on Mg in serum, blood cells, and calf muscle. *Med Sci Sports Exerc*. 1998 Nov;30(11):1584–91.

Whitney EN, Rolfes SR. Understanding Nutrition 7th ed. West Publishing Company, USA, 1996.

WHO. Safe vitamin A dosage during pregnancy and lactation. Recommendations and report from a consultation. Micronutrient series. Geneva: World Health Organization, 1998. (WHO.NUT/98.4.)

Wilborn CD, Kerksick CM, Campbell BI, Taylor LW, Marcello BM, Rasmussen CJ, Greenwood MC, Almada A, Kreider RB. Effects of Zinc Magnesium Aspartate (ZMA) Supplementation on Training Adaptations and Markers of Anabolism and Catabolism. *J Int Soc Sports Nutr*. 2004 Dec 31;1(2):12–20.

Willis WT, Brooks GA, Henderson SA, Dallman PR. Effects of iron deficiency and training on mitochondrial enzymes in skeletal muscle. *J Appl Physiol*. 1987 Jun;62(6):2442–6.

Willis, K.S.; Smith, D.T.; Broughton, K.S.; Larson-Meyer, D.E. Vitamin D status and biomarkers of inflammation in runners. *Open Access J. Sports Med*. 2012, 3, 35–42.

Zheng JJ, Mason JB, Rosenberg IH, Wood RJ. Measurement of zinc bioavailability from beef and a ready-to-eat high-fiber breakfast cereal in humans: application of a whole-gut lavage technique. *Am J Clin Nutr*. 1993 Dec;58(6):902–7.

Ziegler P, Sharp R, Hughes V, Evans W, Khoo CS. Nutritional status of teenage female competitive figure skaters. *J Am Diet Assoc*. 2002 Mar;102(3):374–9.

Zierath J, Kaiserauer S, Snyder AC. Dietary patterns of amenorrheic and regularly menstruating runners. *Med Sci Sports Exerc* 196;18(suppl):S55–6

Zimmermann MB. Iodine deficiency. *Endocr Rev*. 2009 Jun;30(4):376–408.

Zoller H, Vogel W. Iron supplementation in athletes--first do no harm. *Nutrition*. 2004 Jul-Aug;20(7–8):615–9.

Lähdeluettelo luku 10

Abel T, Knechtle B, Perret C, Eser P, von Arx P, Knecht H. Influence of chronic supplementation of arginine aspartate in endurance athletes on performance and substrate metabolism – a randomized, double-blind, placebo-controlled study. *Int J Sports Med*. 2005 Jun;26(5):344–9.

Alvares TS, Conte CA, Paschoalin VM, Silva JT, Meirelles Cde M, Bhambhani YN, Gomes PS. Acute l-arginine supplementation increases muscle blood volume but not strength performance. *Appl Physiol Nutr Metab*. 2012 Feb;37(1):115–26.

American College of Sport Medicine. Round Table, the physiological and health effects of oral creatine supplementation. *Med Sci Sports Exc*. 2000;32:706–717.

Antonio J, Sanders MS, van Gammere D. The effects of bovine colostrum supplementation on body composition and exercise performance in active men and women. *Nutrition*. 2001 Mar;17(3):243–7.

Antonio J, Sanders MS, Kalman D, Woodgate D, Street C. The effects of high-dose glutamine ingestion on weightlifting performance. *J Strength Cond Res*. 2002 Feb;16(1):157–60.

Antonio J, Ciccone V. The effects of pre versus post workout supplementation of creatine monohydrate on body composition and strength. *J Int Soc Sports Nutr*. 2013 Aug 6;10(1):36.

Anomasiri W, Sanguanrungsingkul S, Saichandee P. Low dose creatine supplementation enhances sprint phase of 400 meters swimming performance. *J Med Assoc Thai*. 2004 Sep;87 Suppl 2:S228–32.

Applegate E. Effective nutritional ergogenic aids. *Int J Sport Nutr*. 1999 Jun;9(2):229–39.

Armstrong LE. Caffeine, body fluid-electrolyte balance, and exercise performance. *Int J Sport Nutr Exerc Metab*. 2002 Jun;12(2):189–206.

Armstrong LE, Pumerantz AC, Roti MW, Judelson DA, Watson G, Dias JC, Sokmen B, Casa DJ ym. Fluid, electrolyte, and renal indices of hydration during 11 days of controlled caffeine consumption. *Int J Sport Nutr Exerc Metab*. 2005 Jun;15(3):252–65.

Astorino TA, Roberson DW. Efficacy of acute caffeine ingestion for short-term high-intensity exercise performance: a systematic review. *J Strength Cond Res*. 2010 Jan;24(1):257–65.

Avelar-Escobar G, Méndez-Navarro J, Ortiz-Olvera NX, Castellanos G, Ramos R, Gallardo-Cabrera VE, Vargas-Alemán Jde J, Díaz de León O, Rodríguez EV, Dehesa-Violante M. Hepatotoxicity associated with dietary energy supplements: use and abuse by young athletes. *Ann Hepatol*. 2012 Jul-Aug;11(4):564–9.

Balage M, Dardevet D. Long-term effects of leucine supplementation on body composition. *Curr Opin Clin Nutr Metab Care*. 2010; 13: 265–70.

Balsom PD, Soderlund K, Ekblom B. Creatine in humans with special reference to creatine supplementation. *Sports Med*. 1994 Oct;18(4):268–80.

Baume N, Mahler N, Kamber M, Mangin P, Saugy M. Research of stimulants and anabolic steroids in dietary supplements. *Scand J Med Sci Sports*. 2006 Feb;16(1):41–8.

Bazzucchi I, Felici F, Sacchetti M. Effect of short-term creatine supplementation on neuromuscular function. *Med Sci Sports Exerc*. 2009 Oct;41(10):1934–41.

Bell DG, McLellan TM. Exercise endurance 1, 3, and 6 h after caffeine ingestion in caffeine users and nonusers. *J Appl Physiol*. 2002 Oct;93(4):1227–34.

Bescós R, Sureda A, Tur JA, Pons A. The effect of nitric-oxide-related supplements on human performance. *Sports Med*. 2012 Feb 1;42(2):99–117.

Bigelman KA, Fan EH, Chapman DP, Freese EC, Trilk JL, Cureton KJ. Effects of six weeks of quercetin supplementation on physical performance in ROTC cadets. *Mil Med*. 2010 Oct;175(10):791–8.

Birnbaum LJ, Herbst JD. Physiologic effects of caffeine on cross-country runners. *J Strength Cond Res*. 2004 Aug;18(3):463–5.

Bizzarini E, De Angelis L. Is the use of oral creatine supplementation safe? *J Sports Med Phys Fitness*. 2004 Dec;44(4):411–6.

Branch JD. Effect of creatine supplementation on body composition and performance: a meta-analysis. *Int J Sport Nutr Exerc Metab*. 2003 Jun;13(2):198–226.

Brass EP. Carnitine and sports medicine: use or abuse? *Ann N Y Acad Sci*. 2004 Nov;1033:67–78.

Brien S, Prescott P, Lewith G. Meta-analysis of the related nutritional supplements dimethyl sulfoxide and methylsulfonylmethane in the treatment of osteoarthritis of the knee. *Evid Based Complement Alternat Med*. 2011;2011:528403.

Brinkworth GD, Buckley JD, Bourdon PC, Gulbin JP, David A. Oral bovine colostrum supplementation enhances buffer capacity but not rowing performance in elite female rowers. *Int J Sport Nutr Exerc Metab*. 2002 Sep;12(3):349–65.

Brinkworth GD, Buckley JD. Bovine colostrum supplementation does not affect plasma buffer capacity or haemoglobin content in elite female rowers. *Eur J Appl Physiol.* 2004 Mar;91(2–3):353–6. Epub 2003 Dec 18.

Brinkworth GD, Buckley JD, Slavotinek JP, Kurmis AP. Effect of bovine colostrum supplementation on the composition of resistance trained and untrained limbs in healthy young men. *Eur J Appl Physiol.* 2004 Jan;91(1):53–60.

Bruce CR, Anderson ME, Fraser SF, Stepto NK, Klein R, Hopkins WG, Hawley JA. Enhancement of 2000-m rowing performance after caffeine ingestion. *Med Sci Sports Exerc.* 2000 Nov;32(11):1958–63.

Burke L, Desbrow B, Minehan M.. Dietary supplements and nutritional ergogenic aids. In: *Clinical Sports Nutrition* (2nd ed.), edited by L. Burke and V. Deakin. Sydney: McGraw-Hill, 2000, p. 455–528.

Burke DG, Chilibeck PD, Parise G, Candow DG, Mahoney D, Tarnopolsky M. Effect of creatine and weight training on muscle creatine and performance in vegetarians. *Med Sci Sports Exerc.* 2003 Nov;35(11):1946–55.

Burke DG, Candow DG, Chilibeck PD, MacNeil LG, Roy BD, Tarnopolsky MA, Ziegenfuss T. Effect of creatine supplementation and resistance-exercise training on muscle insulin-like growth factor in young adults. *Int J Sport Nutr Exerc Metab.* 2008 Aug;18(4):389–98.

Burke LM. Practical considerations for bicarbonate loading and sports performance. *Nestle Nutr Inst Workshop Ser.* 2013;75:15–26.

Candow DG, Chilibeck PD, Burke D, Daviseon KS, Smith-Palmer T. Effect of glutamine supplementation combined with resistance training in young adults. *Eur J Appl Physiol.* 2001 Dec;86(2):142–9.

Casuso RA, Martínez-López EJ, Nordsborg NB, Hita-Contreras F, Martínez-Romero R, Cañuelo A, Martínez-Amat A. Oral quercetin supplementation hampers skeletal muscle adaptations in response to exercise training. *Scand J Med Sci Sports.* 2013 Oct 14.

Ceremuzynski L, Chamiec T, Herbaczynska-Cedro K. Effect of supplemental oral L-arginine on exercise capacity in patients with stable angina pectoris. *Am J Cardiol.* 1997 Aug 1;80(3):331–3.

Cermak NM, Gibala MJ, van Loon LJ. Nitrate supplementation's improvement of 10-km time-trial performance in trained cyclists. *Int J Sport Nutr Exerc Metab.* 2012;22:64–71.

Chromiak J Antonio J. Use of amino acids as growth hormone-releasing agents by athletes. *Nutrition.* 2002 Jul-Aug;18(7–8):657–61.

Chwalbinska-Moneta J. Effect of creatine supplementation on aerobic performance and anaerobic capacity in elite rowers in the course of endurance training. *Int J Sport Nutr Exerc Metab.* 2003 Jun;13(2):173–83.

Cole KJ, Costill DL, Starling RD, Goodpaster BH, Trappe SW, Fink WJ. Effect of caffeine ingestion on perception of effort and subsequent work production. *Int J Sport Nutr.* 1996 Mar;6(1):14–23

Collomp K, Ahmaidi S, Chatard JC, Audran M, Prefaut C. Benefits of caffeine ingestion on sprint performance in trained and untrained swimmers. *Eur J Appl Physiol Occup Physiol.* 1992;64(4):377–80.

Colombani P, Wenk C, Kunz I, Krahnenbuhl S, Kuhnt M, Arnold M, Frey-Rindova P, Frey W, Langhans W. Effects of L-carnitine supplementation on physical performance and energy metabolism of endurance-trained athletes: a double-blind crossover field study. *Eur J Appl Physiol Occup Physiol.* 1996;73(5):434–9.

Colombani PC, Bitzi R, Frey-Rindova P, Frey W, Arnold M, Langhans W, Wenk C. Chronic arginine aspartate supplementation in runners reduces total plasma amino acid level at rest and during a marathon run. *Eur J Nutr.* 1999 Dec;38(6):263–70.

Cook CJ, Crewther BT, Kilduff LP, Drawer S, Gaviglio CM. Skill execution and sleep deprivation: effects of acute caffeine or creatine supplementation – a randomized placebo-controlled trial. *J Int Soc Sports Nutr.* 2011 Feb 16;8:2.

Coombes JS, Conacher M, Austen SK, Marshall PA. Dose effects of oral bovine colostrum on physical work capacity in cyclists. *Med Sci Sports Exerc.* 2002 Jul;34(7):1184–8.

Cooper R, Naclerio F, Allgrove J, Jimenez A. Creatine supplementation with specific view to exercise/sports performance: an update. *J Int Soc Sports Nutr.* 2012; 9: 33.

- Cox G, Mujika I, Tumilty D, Burke L.** Acute creatine supplementation and performance during a field test simulating match play in elite female soccer players. *Int J Sport Nutr Exerc Metab.* 2002 Mar;12(1):33–46.
- Dalbo VJ, Roberts MD, Stout JR, Kerksick CM.** Putting to rest the myth of creatine supplementation leading to muscle cramps and dehydration. *Br J Sports Med.* 2008; 42: 567–73.
- Daneshvar P, Hariri M, Ghiasvand R, Askari G, Darvishi L, Mashhadi NS, Khosravi-Boroujeni H.** Effect of eight weeks of quercetin supplementation on exercise performance, muscle damage and body muscle in male badminton players. *Int J Prev Med.* 2013 Apr;4(Suppl 1):S53–7.
- Davis JM, Murphy EA, Carmichael MD, Davis B.** Quercetin increases brain and muscle mitochondrial biogenesis and exercise tolerance. *Am J Physiol Regul Integr Comp Physiol.* 2009 Apr;296(4):R1071–7.
- Davison G.** Bovine colostrum and immune function after exercise. *Med Sport Sci.* 2012;59:62–9.
- Decombaz J, Gmuender B, Sierro G, Cerretelli P.** Muscle carnitine after strenuous endurance exercise. *J Appl Physiol.* 1992 Feb;72(2):423–7.
- Decombaz J, Deriaz O, Acheson K, Gmuender B, Jequier E.** Effect of L-carnitine on submaximal exercise metabolism after depletion of muscle glycogen. *Med Sci Sports Exerc.* 1993 Jun;25(6):733–40.
- Denadai BS, Denadai ML.** Effects of caffeine on time to exhaustion in exercise performed below and above the anaerobic threshold. *Braz J Med Biol Res.* 1998 Apr;31(4):581–5.
- de Salles Painelli V, Saunders B, Sale C, Harris RC, Solis MY, Roschel H, Gualano B, Artioli GG, Lancha AH Jr.** Influence of training status on high-intensity intermittent performance in response to -alanine supplementation. *Amino Acids.* 2014 May;46(5):1207–15.
- DiRienzo DB.** Effect of probiotics on biomarkers of cardiovascular disease: implications for heart-healthy diets. *Nutr Rev.* 2014 Jan;72(1):18–29.
- Doherty M, Smith PM, Davison RC, Hughes MG.** Caffeine is ergogenic after supplementation of oral creatine monohydrate. *Med Sci Sports Exerc.* 2002 Nov;34(11):1785–92.
- Doherty M, Smith P, Hughes M, Davison R.** Caffeine lowers perceptual response and increases power output during high-intensity cycling. *J Sports Sci.* 2004 Jul;22(7):637–43.
- Doherty M, Smith PM.** Effects of caffeine ingestion on rating of perceived exertion during and after exercise: a meta-analysis. *Scand J Med Sci Sports.* 2005 Apr;15(2):69–78.
- Donaldson SKB, Hermansen L.** Differential direct effects of H⁺ and Ca²⁺-activated force of skinned fibres from the soleus, cardiac, adductor magnus muscle of rabbits. *Pflugers Arch.* 1978;376:55–65.
- Dong JY, Qin LQ, Zhang Z, Zhao Y, Wang J, Arigoni F, Zhang W.** Effect of oral L-arginine supplementation on blood pressure: a meta-analysis of randomized, double-blind, placebo-controlled trials. *Am Heart J.* 2011 Dec;162(6):959–65.
- Dong JY, Szeto IM, Makinen K, Gao Q, Wang J, Qin LQ, Zhao Y.** Effect of probiotic fermented milk on blood pressure: a meta-analysis of randomised controlled trials. *Br J Nutr.* 2013 Oct;110(7):1188–94.
- Earnest CP, Almada AL, Mitchell TL.** Effects of creatine monohydrate ingestion upon intermediate length anaerobic treadmill running to exhaustion. *J Strength Cond Res* 1997;4:234–238
- Engelhardt M, Neumann G, Berbalk A, Reuter I.** Creatine supplementation in endurance sports. *Med Sci Sports Exerc.* 1998 Jul;30(7):1123–9.
- Eto B, Peres G, Le Moel G.** Effects of an ingested glutamate arginine salt on ammonemia during and after long lasting cycling. *Arch Int Physiol Biochim Biophys.* 1994 May-Jun;102(3):161–2.
- Ganio MS, Klau JF, Casa DJ, Armstrong LE, Maresh CM.** Effect of caffeine on sport-specific endurance performance: a systematic review. *J Strength Cond Res.* 2009 Jan;23(1):315–24.
- Ganio MS, Armstrong LE, Johnson EC, Klau JF, Ballard KD, Michniak-Kohn B, Kaushik D, Maresh CM.** Effect of quercetin supplementation on maximal oxygen uptake in men and women. *J Sports Sci.* 2010 Jan;28(2):201–8.
- Geyer H, Parr MK, Mareck U, Reinhart U, Schrader Y, Schänzer W.** Analysis of non-hormonal nutritional supplements for anabolic-androgenic steroids – results of an international study. *Int J Sports Med.* 2004 Feb;25(2):124–9.

Gleeson M. Interrelationship between physical activity and branched-chain amino acids. *J Nutr.* 2005 Jun;135(6 Suppl):1591S-5S.

Gleeson M. Nutritional support to maintain proper immune status during intense training. *Nestle Nutr Inst Workshop Ser.* 2013;75:85–97.

Gleeson M, Bishop NC, Oliveira M, Tauler P. Daily probiotic's (*Lactobacillus casei* Shirota) reduction of infection incidence in athletes. *Int J Sport Nutr Exerc Metab.* 2011 Feb;21(1):55–64.

Gleeson M, Williams C. Intense exercise training and immune function. *Nestle Nutr Inst Workshop Ser.* 2013;76:39–50.

Goldfinch J, Mc Naughton L, Davies P. Induced metabolic alkalosis and its effects on 400-m racing time. *Eur J Appl Physiol Occup Physiol.* 1988;57(1):45–8.

Gomez-Cabrera MC, Domenech E, Romagnoli M, Arduini A, Borras C, Pallardo FV, Sastre J, Viña J. Oral administration of vitamin C decreases muscle mitochondrial biogenesis and hampers training-induced adaptations in endurance performance. *Am J Clin Nutr.* 2008 Jan;87(1):142–9.

Graham TE, Spriet LL. Performance and metabolic responses to a high caffeine dose during prolonged exercise. *J Appl Physiol.* 1991 Dec;71(6):2292–8.

Graham TE, Hibbert E, Sathasivam P. Metabolic and exercise endurance effects of coffee and caffeine ingestion. *J Appl Physiol.* 1998 Sep;85(3):883–9.

Greenhaff PL, Bodin K, Soderlund K, Hultman E. Effect of oral creatine supplementation on skeletal muscle phosphocreatine resynthesis. *Am J Physiol.* 1994 May;266(5 Pt 1):E725–30.

Greenwood M, Kreider RB, Greenwood L, Byars A. Cramping and Injury Incidence in Collegiate Football Players Are Reduced by Creatine Supplementation. *J Athl Train.* 2003 Sep;38(3):216–219.

Greer BK, Jones BT. Acute arginine supplementation fails to improve muscle endurance or affect blood pressure responses to resistance training. *J Strength Cond Res.* 2011 Jul;25(7):1789–94.

Hao Q, Lu Z, Dong BR, Huang CQ, Wu T. Probiotics for preventing acute upper respiratory tract infections. *Cochrane Database Syst Rev.* 2011 Sep 7;(9):CD006895.

Harris RC, Edwards RHT, Hultman E, Nordesjo LO, Nylin B. The time course of phosphorylcreatine resynthesis during recovery of the quadriceps muscle in man. *Pflugers Arch.* 1976;367:137–142.

Harris RC, Soderlund K, Hultman E. Elevation of creatine in resting and exercised muscle of normal subjects by creatine supplementation. *Clin Sci (Lond).* 1992 Sep;83(3):367–74.

Harris RC, Stellingwerff T. Effect of -alanine supplementation on high-intensity exercise performance. *Nestle Nutr Inst Workshop Ser.* 2013;76:61–71

Harwood M, Danielewska-Nikiel B, Borzelleca JF, Flamm GW, Williams GM, Lines TC. A critical review of the data related to the safety of quercetin and lack of evidence of in vivo toxicity, including lack of genotoxic/carcinogenic properties. *Food Chem Toxicol.* 2007;45(11):2179–2205.

Haywood BA, Black KE, Baker D, McGarvey J, Healey P, Brown RC. Probiotic supplementation reduces the duration and incidence of infections but not severity in elite rugby union players. *J Sci Med Sport.* 2014 Jul;17(4):356–60.

Heikkinen A, Alaranta A, Helenius I, Vasankari T. Use of dietary supplements in Olympic athletes is decreasing: a follow-up study between 2002 and 2009. *J Int Soc Sports Nutr.* 2011 Feb 4;8(1):1.

Heinonen OJ. Carnitine and physical exercise. *Sports Med.* 1996 Aug;22(2):109–32.

Heinz SA, Henson DA, Austin MD, Jin F, Nieman DC. Quercetin supplementation and upper respiratory tract infection: A randomized community clinical trial. *Pharmacol Res.* 2010 Sep;62(3):237–42.

Henrotin Y, Chevalier X, Herrero-Beaumont G, McAlindon T, Mobasher A, Pavelka K, Schön C, Weinans H, Biesalski H; Participants at the Hohenheim Consensus Conference in August 29th 2011. Physiological effects of oral glucosamine on joint health: current status and consensus on future research priorities. *BMC Res Notes.* 2013 Mar 26;6:115.

Hickner RC, Dyck DJ, Sklar J, Hatley H, Byrd P. Effect of 28 days of creatine ingestion on muscle metabolism and performance of a simulated cycling road race. *J Int Soc Sports Nutr.* 2010 Jul;7:26.

Hobson RM, Saunders B, Ball G, Harris RC, Sale C. Effects of -alanine supplementation on exercise performance: a meta-analysis. *Amino Acids.* 2012 Jul;43(1):25–37.

Hobson RM, Harris RC, Martin D, Smith P, Macklin B, Gualano B, Sale C. Effect of Beta-Alanine With and Without Sodium Bicarbonate on 2,000-m Rowing Performance. *Int J Sport Nutr Exerc Metab.* 2013 Oct;23(5):480–7.

Hofman Z, Smeets R, Verlaan G, Lugt R, Verstappen PA. The effect of bovine colostrum supplementation on exercise performance in elite field hockey players. *Int J Sport Nutr Exerc Metab.* 2002 Dec;12(4):461–9.

Hoon MW, Johnson NA, Chapman PG, Burke LM. The effect of nitrate supplementation on exercise performance in healthy individuals: a systematic review and meta-analysis. *Int J Sport Nutr Exerc Metab.* 2013 Oct;23(5):522–32.

Huang A, Owen K. Role of supplementary L-carnitine in exercise and exercise recovery. *Med Sport Sci.* 2012;59:135–42.

Huertas R, Campos Y, Diaz E, Esteban J, Vechietti L, Montanari G, D'Iddio S, Corsi M, Arenas J. Respiratory chain enzymes in muscle of endurance athletes: effect of L-carnitine. *Biochem Biophys Res Commun.* 1992 Oct 15;188(1):102–7.

Hultman E, Soderlund K, Timmons JA, Cederblad G, Greenhaff PL. Muscle creatine loading in men. *J Appl Physiol.* 1996 Jul;81(1):232–7.

Hursel R, Westerterp-Plantenga MS. Green tea catechin plus caffeine supplementation to a high-protein diet has no additional effect on body weight maintenance after weight loss. *Am J Clin Nutr.* 2009; 89: 822–30.

Ibanez J, Pullinen T, Gorostiaga E, Postigo A, Mero A. Blood lactate and ammonia in short-term anaerobic work following induced alkalosis. *J Sports Med Phys Fitness.* 1995 Sep;35(3):187–93.

ISSN; Buford T, Kreider R, Stout J, Greenwood M, Campbell B, Spano M, Ziegenfuss T ym. International Society of Sports Nutrition position stand: creatine supplementation and exercise. *J Int Soc Sports Nutr.* 2007;4:6.

ISSN; Goldstein ER, Ziegenfuss T, Kalman D, Kreider R, Campbell B, Wilborn C, Taylor L ym. International society of sports nutrition position stand: caffeine and performance. *J Int Soc Sports Nutr.* 2010 Jan 27;7(1):5.

Jagim AR, Oliver JM, Sanchez A, Galvan E, Fluckey J, Riechman S, Greenwood M, Kelly K, ym. A buffered form of creatine does not promote greater changes in muscle creatine content, body composition, or training adaptations than creatine monohydrate. *J Int Soc Sports Nutr.* 2012; 9:43.

Jones AM, Atter T, Georg KP. Oral creatine supplementation improves multiple sprint performance in elite ice-hockey players. *J Sports Med Phys Fitness.* 1999 Sep;39(3):189–96.

Jones AM, Bailey SJ, Vanhatalo A, Fulford J, Gilchrist M, Benjamin N, Winyard PG. Reply to Lundberg, Larsen, and Weitzberg. *J Appl Physiol* (1985). 2011 Aug;111(2):619

Jones AM, Vanhatalo A, Bailey SJ. Influence of dietary nitrate supplementation on exercise tolerance and performance. *Nestle Nutr Inst Workshop Ser.* 2013;75:27–40.

Jones AM. Dietary nitrate supplementation and exercise performance. *Sports Med.* 2014 May;44 Suppl 1:S35–45.

Jówko E, Ostaszewski P, Jank M, Sacharuk J, Zieniewicz A, Wilczak J, Nissen S. Creatine and beta-hydroxy-beta-methylbutyrate (HMB) additively increase lean body mass and muscle strength during a weight-training program. *Nutrition.* 2001 Jul-Aug;17(7–8):558–66

Juhn MS, Tarnopolsky M. Potential side effects of oral creatine supplementation: a critical review. *Clin J Sport Med.* 1998 Oct;8(4):298–304.

Jäger R, Metzger J, Lautmann K, Shushakov V, Purpura M, Geiss KR, Maassen N. The effects of creatine pyruvate and creatine citrate on performance during high intensity exercise. *J Int Soc Sports Nutr.* 2008 Feb 13;5:4

Kalmar JM, Cafarelli E. Caffeine: a valuable tool to study central fatigue in humans? *Exerc Sport Sci Rev.* 2004 Oct;32(4):143–7

Kang E-J, Kim SY, Hwang I-H, et al. (2013) The effect of probiotics on prevention of common cold: a meta-analysis of randomized controlled trial studies. *Korean J Fam Med* 34, 2–10

Kekkonen RA, Vasankari TJ, Vuorimaa T, Haahtela T, Julkunen I, Korpela R. The effect of probiotics on respiratory infections and gastrointestinal symptoms during training in marathon runners. *Int J Sport Nutr Exerc Metab.* 2007 Aug;17(4):352–63.

Kelly GS. Bovine colostrums: a review of clinical uses. *Altern Med Rev.* 2003 Nov;8(4):378–94.

Killer SC, Blannin AK, Jeukendrup AE. No evidence of dehydration with moderate daily coffee intake: a counterbalanced cross-over study in a free-living population. *PLoS One.* 2014 Jan 9;9(1):e84154.

Kim HJ, Kim CK, Carpentier A, Poortmans JR. Studies on the safety of creatine supplementation. *Amino Acids.* 2011 May;40(5):1409–18.

King S, Glanville J, Sanders ME, Fitzgerald A, Varley D. Effectiveness of probiotics on the duration of illness in healthy children and adults who develop common acute respiratory infectious conditions: a systematic review and meta-analysis. *Br J Nutr.* 2014 Jul 14;112(1):41–54.

Knitter AE, Panton L, Rathmacher JA, Petersen A, Sharp R. Effects of beta-hydroxy-beta-methylbutyrate on muscle damage after a prolonged run. *J Appl Physiol.* 2000 Oct;89(4):1340–4.

Koopman R, Wagenmakers AJM, Manders RJ, Zorenc AHG, Senden JMG, Gorselink J, Kreizer HA, van Loon LJC. Combined ingestion of protein and free leucine with carbohydrate increases postexercise muscle protein synthesis in vivo in male subjects. *Am J Physiol Endocrinol Metab.* 2005 Apr;288(4):E645–53.

Kreider RB. Dietary supplements and the promotion of muscle growth with resistance exercise. *Sports Med.* 1999 Feb;27(2):97–110.

Kreider RB, Ferreira M, Wilson M, Almada AL. Effects of calcium beta-hydroxy-beta-methylbutyrate (HMB) supplementation during resistance-training on markers of catabolism, body composition and strength. *Int J Sports Med.* 1999 Nov;20(8):503–9.

Kreider RB, Ferreira MP, Greenwood M, Wilson M, Almada AL. Effects of conjugated linoleic acid supplementation during resistance training on body composition, bone density, strength, and selected hematological markers. *J Strength Cond Res.* 2002 Aug;16(3):325–34.

Kreider RB. Effects of creatine supplementation on performance and training adaptations. *Mol Cell Biochem.* 2003 Feb;244(1–2):89–94

Kreider RB, Wilborn CD, Taylor L, Campbell B, Almada AL, Collins R, Cooke M, Earnest CP, Greenwood M, Kalman DS, Kerksick CM, ym.. ISSN exercise & sport nutrition review: research & recommendations. *J Int Soc Sports Nutr.* 2010; 7: 7.

Kressler J, Millard-Stafford M, Warren GL. Quercetin and endurance exercise capacity: a systematic review and meta-analysis. *Med Sci Sports Exerc.* 2011 Dec;43(12):2396–404.

Kuipers H, van Breda E, Verlaan G, Smeets R. Effects of oral bovine colostrum supplementation on serum insulin-like growth factor-I levels. *Nutrition.* 2002 Jul-Aug;18(7–8):566–7.

Lambert EV, Goedecke JH, Bluett K, Heggie K, Claassen A, Rae DE, West S, Dugas J, Dugas L, Meltzeri S, Charlton K, Mohede I. Conjugated linoleic acid versus high-oleic acid sunflower oil: effects on energy metabolism, glucose tolerance, blood lipids, appetite and body composition in regularly exercising individuals. *Br J Nutr.* 2007 May;97(5):1001–11.

Lamprecht M, Frauwallner A. Exercise, intestinal barrier dysfunction and probiotic supplementation. *Med Sport Sci.* 2012;59:47–56.

Laurent D, Schneider KE, Prusaczyk WK, Franklin C, Vogel SM, Krssak M, Petersen KF, Goforth HW ym. Effects of caffeine on muscle glycogen utilization and the neuroendocrine axis during exercise. *J Clin Endocrinol Metab.* 2000 Jun;85(6):2170–5

Lee CL, Lin JC, Cheng CF. Effect of caffeine ingestion after creatine supplementation on intermittent high-intensity sprint performance. *Eur J Appl Physiol.* 2011 Aug;111(8):1669–77

Linderman J, Fahey TD. Sodium bicarbonate ingestion and exercise performance. An update. *Sports Med.* 1991 Feb;11(2):71–7.

Linderman JK, Gosselink KL. The effects of sodium bicarbonate ingestion on exercise performance. *Sports Med.* 1994 Aug;18(2):75–80

Lopez RM, Casa DJ, McDermott BP, Ganio MS, Armstrong LE, Maresh CM. Does creatine supplementation hinder exercise heat tolerance or hydration status? A systematic review with meta-analyses. *J Athl Train.* 2009 Mar-Apr;44(2):215–23.

Lundberg JO, Larsen FJ, Weitzberg E. Supplementation with nitrate and nitrite salts in exercise: a word of caution. *J Appl Physiol* (1985). 2011 Aug;111(2):616–7.

MacRae HS, Mefferd KM. Dietary antioxidant supplementation combined with quercetin improves cycling time trial performance. *Int J Sport Nutr Exerc Metab.* 2006;16:405–19.

Madsen K, MacLean DA, Kiens B, Christensen D. Effects of glucose, glucose plus branched-chain amino acids, or placebo on bike performance over 100 km. *J Appl Physiol.* 1996 Dec;81(6):2644–50.

Marconi C, Sassi G, Carpinelli A, Cerretelli P. Effects of L-carnitine loading on the aerobic and anaerobic performance of endurance athletes. *Eur J Appl Physiol Occup Physiol.* 1985;54(2):131–5.

Martello S, Felli M, Chiarotti M. Survey of nutritional supplements for selected illegal anabolic steroids and ephedrine using LC-MS/MS and GC-MS methods, respectively. *Food Addit Contam.* 2007 Mar;24(3):258–65.

Matson LG, Tran ZV. Effects of sodium bicarbonate ingestion on anaerobic performance: a meta-analytic review. *Int J Sport Nutr.* 1993 Mar;3(1):2–28.

Matthews DE. Observations of branched-chain amino acid administration in humans. *J Nutr.* 2005 Jun;135(6 Suppl):1580S-4S.

Maughan RJ. Creatine supplementation and exercise performance. *Int J Sport Nutr.* 1995 Jun;5(2):94–101

Maughan RJ. Risks and rewards of dietary supplement use by athletes. IOC Consensus In Sports Nutrition Conference Lausanne, Switzerland 25th–27th October 2010.

Metzl JD, Small E, Levine SR, Gershel JC. Creatine use among young athletes. *Pediatrics.* 2001;108:421–425.

McKenna MJ, Medved I, Goodman CA, Brown MJ, Bjorksten AR, Murphy KT, Petersen AC, Sostaric S, Gong X. N-acetylcysteine attenuates the decline in muscle Na⁺,K⁺-pump activity and delays fatigue during prolonged exercise in humans. *J Physiol.* 2006 Oct 1;576(Pt 1):279–88.

Medved I, Brown MJ, Bjorksten AR, Murphy KT, Petersen AC, Sostaric S, Gong X, McKenna MJ. N-acetylcysteine enhances muscle cysteine and glutathione availability and attenuates fatigue during prolonged exercise in endurance-trained individuals. *J Appl Physiol* (1985). 2004 Oct;97(4):1477–85.

Moreira A, Kekkonen RA, Delgado L, Fonseca J, Korpela R, Haahtela T. Nutritional modulation of exercise-induced immunodepression in athletes: a systematic review and meta-analysis. *Eur J Clin Nutr.* 2007 Apr;61(4):443–60.

Motl RW, O'Connor PJ, Dishman RK. Effect of caffeine on perceptions of leg muscle pain during moderate intensity cycling exercise. *J Pain.* 2003 Aug;4(6):316–21

Mujika I, Padilla S, Ibanez J, Izquierdo M, Gorostiaga E. Creatine supplementation and sprint performance in soccer players. *Med Sci Sports Exerc.* 2000 Feb;32(2):518–25.

Newsholme EA, Blomstrand E. Tryptophan, 5-hydroxytryptamine and a possible explanation for central fatigue. *Adv Exp Med Biol.* 1995;384:315–20.

Nieman DC, Henson DA, Gross SJ, Jenkins DP, Davis JM, Murphy EA, Carmichael MD, Dumke CL, Utter AC, McAnulty SR, McAnulty LS, Mayer EP. Quercetin reduces illness but not immune perturbations after intensive exercise. *Med Sci Sports Exerc.* 2007 Sep;39(9):1561–9.

Nieman DC, Williams AS, Shanely RA, Jin F, McAnulty SR, Triplett NT, Austin MD, Henson DA. Quercetin's influence on exercise performance and muscle mitochondrial biogenesis. *Med Sci Sports Exerc.* 2010 Feb;42(2):338–45.

Nikolaidis MG, Kerksick CM, Lamprecht M, McAnulty SR. Does vitamin C and E supplementation impair the favorable adaptations of regular exercise? *Oxid Med Cell Longev.* 2012;2012:707941.

Nissen SL ja Sharp RL. Effect of dietary supplements on lean mass and strength gains with resistance exercise: a meta-analysis. *J Appl Physiol.* 2003 Feb;94(2):651–9.

Norris LE, Collene AL, Asp ML, Hsu JC, Liu LF, Richardson JR, Li D, Bell D, Osei K, Jackson RD, Belury MA. Comparison of dietary conjugated linoleic acid with safflower oil on body composition in obese postmenopausal women with type 2 diabetes mellitus. *Am J Clin Nutr.* 2009 Sep;90(3):468–76.

O'Connor DM, Crowe MJ. Effects of beta-hydroxy-beta-methylbutyrate and creatine monohydrate supplementation on the aerobic and anaerobic capacity of highly trained athletes. *J Sports Med Phys Fitness.* 2003 Mar;43(1):64–8.

Onakpoya IJ, Posadzki PP, Watson LK, Davies LA, Ernst E. The efficacy of long-term conjugated linoleic acid (CLA) supplementation on body composition in overweight and obese individuals: a systematic review and meta-analysis of randomized clinical trials. *Eur J Nutr.* 2012 Mar;51(2):127–34.

Oöpik V, Saaremets I, Medijainen L, Karelson K, Janson T, Timpmann S. Effects of sodium citrate ingestion before exercise on endurance performance in well trained college runners. *Br J Sports Med.* 2003 Dec;37(6):485–9.

Ostojic SM. Creatine supplementation in young soccer players. *Int J Sport Nutr Exerc Metab.* 2004 Feb;14(1):95–103.

Pasiakos SM, McClung HL, McClung JP, Margolis LM, Andersen NE, Cloutier GJ, Pikosky MA, Rood JC, Fielding RA, Young AJ. Leucine-enriched essential amino acid supplementation during moderate steady state exercise enhances postexercise muscle protein synthesis. *Am J Clin Nutr.* 2011; 94: 809–18.

Pasman WJ, van Baak MA, Jeukendrup AE, de Haan A. The effect of different dosages of caffeine on endurance performance time. *Int J Sports Med.* 1995 May;16(4):225–30.

Peternejl TT, Coombes JS. Antioxidant supplementation during exercise training: beneficial or detrimental? *Sports Med.* 2011 Dec 1;41(12):1043–69.

Peeters BM, Lantz CD, Mayhew JL. Effects of oral creatine monohydrate and creatine phosphate supplementation on maximal strength indices, body composition and blood pressure. *J Strength Cond Res* 1998;13:3–9.

Petersen AC, McKenna MJ, Medved I, Murphy KT, Brown MJ, Della Gatta P, Cameron-Smith D. Infusion with the antioxidant N-acetylcysteine attenuates early adaptive responses to exercise in human skeletal muscle. *Acta Physiol (Oxf).* 2012 Mar;204(3):382–92.

Persky AM ja Brazeau GA. Clinical pharmacology of the dietary supplement creatine monohydrate. *Pharmacol Rev.* 2001 Jun;53(2):161–76.

Poortmans JR, Francaux M. Adverse effects of creatine supplementation: fact or fiction? *Sports Med.* 2000 Sep;30(3):155–70.

Portal S, Zadik Z, Rabinowitz J, Pilz-Burstein R, Adler-Portal D, Meckel Y, Cooper DM, Eliakim A, Nemet D. The effect of HMB supplementation on body composition, fitness, hormonal and inflammatory mediators in elite adolescent volleyball players: a prospective randomized, double-blind, placebo-controlled study. *Eur J Appl Physiol.* 2011; 111: 2261–9.

Powers S. Antioxidant supplementaton for athletes: sense **or nonsense?** IOC Consensus In Sports Nutrition Conference Lausanne, Switzerland 25th–27th October 2010.

Potteiger JA, Nickel GL, Webster MJ, Haub MD, Palmer RJ. Sodium citrate ingestion enhances 30 km cycling performance. *Int J Sports Med.* 1996 Jan;17(1):7–11.

Preen D, Dawson B, Goodman C, Beilby J, Ching S. Creatine supplementation: a comparison of loading and maintenance protocols on creatine uptake by human skeletal muscle. *Int J Sport Nutr Exerc Metab.* 2003 Mar;13(1):97–111.

Price M, Moss P. The effects of work:rest duration on physiological and perceptual responses during intermittent exercise and performance. *J Sports Sci.* 2007;25:1613–1621.

Ranchordas MK, Blomstrand E, Calder PC, Burke LM, Stear SJ, Castell LM. Br J Sports Med 2011; 45: 830–831. A–Z of nutritional supplements: dietary supplements, sports nutrition foods and ergogenic aids for health and performance—Part 23.

Ragle RL, Sawitzke AD. Nutraceuticals in the management of osteoarthritis : a critical review. *Drugs Aging.* 2012 Sep;29(9):717–31.

Rainer L ja Heiss CJ. Conjugated linoleic acid: health implications and effects on body composition. *J Am Diet Assoc.* 2004 Jun;104(6):963–8,

Rawson ES, Volek JS. Effects of creatine supplementation and resistance training on muscle strength and weightlifting performance. *J Strength Cond Res.* 2003 Nov;17(4):822–31.

Reid MB, Stokić DS, Koch SM, Khawli FA, Leis AA. N-acetylcysteine inhibits muscle fatigue in humans. *J Clin Invest.* 1994 Dec;94(6):2468–74.

Richy F, Bruyere O, Ethgen O, Cucherat M, Henrotin Y, Reginster JY. Structural and symptomatic efficacy of glucosamine and chondroitin in knee osteoarthritis: a comprehensive meta-analysis. *Arch Intern Med.* 2003 Jul 14;163(13):1514–22.

Risérus U, Vessby B, Arner P, Zethelius B. Supplementation with trans10cis12-conjugated linoleic acid induces hyperproinsulinaemia in obese men: close association with impaired insulin sensitivity. *Diabetologia.* 2004; 47: 1016–9.

Salas-Salvadó J, Márquez-Sandoval F, Bulló M. Conjugated linoleic acid intake in humans: a systematic review focusing on its effect on body composition, glucose, and lipid metabolism. *Crit Rev Food Sci Nutr.* 2006; 46: 479–88.

Saunders B, Sale C, Harris RC, Sunderland C. Sodium bicarbonate and high-intensity-cycling capacity: variability in responses. *Int J Sports Physiol Perform.* 2014 Jul;9(4):627–32

Scholten SD, Sergeev IN. Long-term quercetin supplementation reduces lipid peroxidation but does not improve performance in endurance runners. *Open Access J Sports Med.* 2013 Mar 12;4:53–61.

Schubert MM, Astorino TA. A systematic review of the efficacy of ergogenic aids for improving running performance. *J Strength Cond Res.* 2013 Jun;27(6):1699–707.

Selsby JT, Beckett KD, Kern M, Devor ST. Swim performance following creatine supplementation in Division III athletes. *J Strength Cond Res.* 2003 Aug;17(3):421–4.

Schabot EJ, Wilson G, Noakes TD. Dose-related elevations in venous pH with citrate ingestion do not alter 40-km cycling time-trial performance. *Eur J Appl Physiol.* 2000 Nov;83(4 -5):320–7.

Schilling B, Stone M, Utter A, Kearney J, Johnson M, Coglianese R, Smith L, O'Bryant H, Fry A, Starks M. et al. Creatine supplementation and health variables: a retrospective study. *Med Sci Sports Exerc.* 2001;33:183–188.

Schubert MM, Astorino TA. A systematic review of the efficacy of ergogenic aids for improving running performance. *J Strength Cond Res.* 2013 Jun;27(6):1699–707.

Shave R, Whyte G, Siemann A, Doggart L. The effects of sodium citrate ingestion on 3,000-meter time-trial performance. *J Strength Cond Res.* 2001 May;15(2):230–4.

Shimomura Y, Inaguma A, Watanabe S, Yamamoto Y, Muramatsu Y, Bajotto G, Sato J, ym. Branched-chain amino acid supplementation before squat exercise and delayed-onset muscle soreness. *Int J Sport Nutr Exerc Metab.* 2010; 20: 236–44.

Shimomura Y, Yamamoto Y, Bajotto G, Sato J, Murakami T, Shimomura N, Kobayashi H, Mawatari K. Nutraceutical effects of branched-chain amino acids on skeletal muscle. *J Nutr.* 2006 Feb;136(2):529S–532S.

Shing CM, Jenkins DG, Stevenson L, Coombes JS. The influence of bovine colostrum supplementation on exercise performance in highly trained cyclists. *Br J Sports Med.* 2006 Sep;40(9):797–801

Shing CM, Hunter DC, Stevenson LM. Bovine colostrum supplementation and exercise performance: potential mechanisms. *Sports Med.* 2009;39(12):1033–54.

Siervo M, Lara J, Ogbonwan I, Mathers JC. Inorganic nitrate and beetroot juice supplementation reduces blood pressure in adults: a systematic review and meta-analysis. *J Nutr.* 2013 Jun;143(6):818–26.

Skare OC, Skadberg, Wisnes AR. Creatine supplementation improves sprint performance in male sprinters. *Scand J Med Sci Sports.* 2001 Apr;11(2):96–102.

Slater GJ, Jenkins D. Beta-hydroxy-beta-methylbutyrate (HMB) supplementation and the promotion of muscle growth and strength. *Sports Med.* 2000 Aug;30(2):105–16.

Slattery KM, Dascombe B, Wallace LK, Bentley DJ, Coutts AJ. Effect of N-acetylcysteine on Cycling Performance after Intensified Training. *Med Sci Sports Exerc.* 2014 Jun;46(6):1114–23.

Smith JC, Stephens DP, Hall EL, Jackson AW, Earnest CP. Effect of oral creatine ingestion on parameters of the work rate-time relationship and time to exhaustion in high-intensity cycling. *Eur J Appl Physiol Occup Physiol.* 1998 Mar;77(4):360–5.

Sneddon AA, Tsofliou F, Fyfe CL, Matheson I, Jackson DM, Horgan G, Winzell MS, Wahle KW, Ahren B, Williams LM. Effect of a conjugated linoleic acid and omega-3 fatty acid mixture on body composition and adiponectin. *Obesity (Silver Spring).* 2008; 16: 1019–24.

Spriet LL, MacLean DA, Dyck DJ, Hultman E, Cederblad G, Graham TE. Caffeine ingestion and muscle metabolism during prolonged exercise in humans. *Am J Physiol.* 1992 Jun;262(6 Pt 1):E891–8

Struder HK, Hollmann W, Platen P, Donike M, Gotzmann A, Weber K. Influence of paroxetine, branched-chain amino acids and tyrosine on neuroendocrine system responses and fatigue in humans. *Horm Metab Res.* 1998 Apr;30(4):188–94.

Stuessi C, Hofer P, Meier C, Boutellier U. L-L-Carnitine and the recovery from exhaustive endurance exercise: a randomised, double-blind, placebo-controlled trial. *Eur J Appl Physiol.* 2005 Sep 29;1–5

Syrotuik DG, Bell GJ. Acute creatine monohydrate supplementation: a descriptive physiological profile of responders vs. nonresponders. *J Strength Cond Res.* 2004 Aug;18(3):610–7.

Tang JE, Lysecki PJ, Manolakos JJ, MacDonald MJ, Tarnopolsky MA, Phillips SM. Bolus arginine supplementation affects neither muscle blood flow nor muscle protein synthesis in young men at rest or after resistance exercise. *J Nutr.* 2011 Feb;141(2):195–200.

Tarnopolsky MA. Caffeine and endurance performance. *Sports Med.* 1994 Aug;18(2):109–25

Tarnopolsky M, Cupido C. Caffeine potentiates low frequency skeletal muscle force in habitual and nonhabitual caffeine consumers. *J Appl Physiol.* 2000 Nov;89(5):1719–24.

Terjung RL, Clarkson P, Eichner ER, Greenhaff P, Hespel PJ, Istael RG, Kraemer WJ, Meyer RA, Spriet LL, ym. American College of Sports Medicine roundtable. The physiological and health effects of oral creatine supplementation. *Med Sci Sports Exerc.* 2000 Mar;32(3):706–17.

Terpstra AHM. Effect of conjugated linoleic acid on body composition and plasma lipids in humans: an overview of the literature. *Am J Clin Nutr.* 2004 Mar;79(3):352–61.

Tipton KD, Rasmussen BB, Miller SL, Wolf SE, Owens-Stovall SK, Petroni BE, Wolfe RR. Timing of amino acid-carbohydrate ingestion alters anabolic response of muscle to resistance exercise. *Am J Physiol Endocrinol Metab.* 2001 Aug;281(2):E197–206.

Trewin AJ, Petersen AC, Billaut F, McQuade LR, McInerney BV, Stepto NK. N-acetylcysteine alters substrate metabolism during high-intensity cycle exercise in well-trained humans. *Appl Physiol Nutr Metab.* 2013 Dec;38(12):1217–27.

Unnithan VB, Veehof SH, Vella CA, Kern M. Is there a physiologic basis for creatine use in children and adolescents? *J Strength Cond Res.* 2001;15:524–528.

Vandenbergh K, Gillis N, Van Leemputte M, Van Hecke P, Vanstapel F, Hespel P. Caffeine counteracts the ergogenic action of muscle creatine loading. *J Appl Physiol.* 1996 Feb;80(2):452–7.

Vandenbergh K, Goris M, Van Hecke P, Van Leemputte M, Vangerven L, Hespel P. Long-term creatine intake is beneficial to muscle performance during resistance training. *J Appl Physiol.* 1997; 83: 2055–63.

van Gammeren D. Endurance performance. Kirjassa: Antonio J, Stout JR (toim): Sports Supplements. Lippincott Williams & Wilkins 2001.

Vanhatalo A, Bailey SJ, Blackwell JR, et al. Acute and chronic effects of dietary nitrate supplementation on blood pressure and the physiological responses to moderate-intensity and incremental exercise. *Am J Physiol.* 2010;299:1121–1131.

van Loon LJ, Oosterlaar AM, Hartgens F, Hesselink MK, Snow RJ, Wagenmakers AJ. Effects of creatine loading and prolonged creatine supplementation on body composition, fuel selection, sprint and endurance performance in humans. *Clin Sci (Lond).* 2003 Feb;104(2):153–62.

Van Montfoort MC, Van Dieren L, Hopkins WG, Shearman JP. Effects of ingestion of bicarbonate, citrate, lactate, and chloride on sprint running. *Med Sci Sports Exerc.* 2004 Jul;36(7):1239–43.

van Nieuwenhoven MA, Brummer RM, Brouns F. Gastrointestinal function during exercise: comparison of water, sports drink, and sports drink with caffeine. *J Appl Physiol.* 2000 Sep;89(3):1079–85.

van Nieuwenhoven MA, Brouns F, Kovacs EM. The effect of two sports drinks and water on GI complaints and performance during an 18-km run. *Int J Sports Med.* 2005 May;26(4):281–5.

van Someren K, Fulcher K, McCarthy J, Moore J, Horgan G, Langford R. An investigation into the effects of sodium citrate ingestion on high-intensity exercise performance. *Int J Sport Nutr.* 1998 Dec;8(4):356–63.

Voevodin M, Sinclair A, Gibson R, Makrides M. The effect of CLA on body composition in humans: systematic review and meta-analysis. *Asia Pac J Clin Nutr.* 2005;14 Suppl:S55.

Vukovich MD, Costill DL, Fink WJ. Carnitine supplementation: effect on muscle carnitine and glycogen content during exercise. *Med Sci Sports Exerc.* 1994 Sep;26(9):1122–9.

Wald A, Back C, Bayless TM. Effect of caffeine on the human small intestine. *Gastroenterology.* 1976 Nov;71(5):738–42.

Watson P, Judkins C, Houghton E, Russell C, Maughan RJ. Urinary nandrolone metabolite detection after ingestion of a nandrolone precursor. *Med Sci Sports Exerc.* 2009 Apr;41(4):766–72.

Wilkes D, Gledhill N, Smyth R. Effect of acute induced metabolic alkalosis on 800-m racing time. *Med Sci Sports Exerc.* 1983;15(4):277–80.

Wilkinson DJ, Hossain T, Hill DS, Phillips BE, Crossland H, Williams J, Loughna P, Churchward-Venne TA, Breen L, Phillips SM, Etheridge T, Rathmacher JA, Smith K, Szewczyk NJ, Atherton PJ. Effects of leucine and its metabolite β -hydroxy- β -methylbutyrate on human skeletal muscle protein metabolism. *J Physiol.* 2013 Jun 1;591(Pt 11):2911–23.

Williams MH. Facts and fallacies of purported ergogenic amino acid supplements. *Clin Sports Med.* 1999 Jul;18(3):633–49

Williams M. Dietary supplements and sport performance: amino acids. *J Int Soc Sports Nutr* 2005;2(2):63–67.

Williamson G, Manach C. Bioavailability and bioefficacy of polyphenols in humans. II. Review of 93 intervention studies. *Am J Clin Nutr.* 2005;81:243S–55

Wilson JM, Fitschen PJ, Campbell B, Wilson GJ, Zanchi N, Taylor L, Wilborn C, Kalman DS, Stout JR, Hoffman JR, Ziegenfuss TN, Lopez HL, Kreider RB, Smith-Ryan AE, Antonio J. International Society of Sports Nutrition Position Stand: beta-hydroxy-beta-methylbutyrate (HMB). *J Int Soc Sports Nutr.* 2013 Feb 2;10(1):6.

Wilson JM, Lowery RP, Joy JM, Walters JA, Baier SM, Fuller JC Jr, Stout JR, Norton LE, Sikorski EM, Wilson SM, Duncan NM, Zanchi NE, Rathmacher J. β -Hydroxy- β -methylbutyrate free acid reduces markers of exercise-induced muscle damage and improves recovery in resistance-trained men. *Br J Nutr.* 2013 Aug 28;110(3):538–44.

West NP, Pyne DB, Cripps AW, Hopkins WG, Eskesen DC, Jairath A, Christophersen CT, Conlon MA, Fricker PA. Lactobacillus fermentum (PCC®) supplementation and gastrointestinal and respiratory-tract illness symptoms: a randomised control trial in athletes. *Nutr J.* 2011 Apr 11;10:30.

West NP, Pyne DB, Peake JM, Cripps AW. Probiotics, immunity and exercise: a review. *Exerc Immunol Rev.* 2009;15:107–26. Review.

Zajac A, Poprzeczk S, Zebrowska A, Chalimoniuk M, Langfort J. Arginine and ornithine supplementation increases growth hormone and insulin-like growth factor-1 serum levels after heavy-resistance exercise in strength-trained athletes. *J Strength Cond Res.* 2010 Apr;24(4):1082–90.

Zhu Q, Yue X, Tian QY, Saren G, Wu MH, Zhang Y, Liu TT. Effect of L-arginine supplementation on blood pressure in pregnant women: a meta-analysis of placebo-controlled trials. *Hypertens Pregnancy.* 2013;32(1):32–41.

Ziegenfuss T, Gales D, Felix S, ym. Performance benefits following a five day creatine loading procedure persist for at least four weeks. *Med Sci Sports Exerc* 1998;30(Suppl):265